


AUTHOR INDEX

Animal Feed Science and Technology, volumes 1-25



A

- Abate, A., Kayongo-Male, H. and Karue, C.N.
Dry matter, protein, energy and fibre intake by dairy heifers grazing a Rhodes grass (*Chloris gayana*) pasture 6(1981)15
- Abate, A. and Pfeffer, E.
Changes in nutrient intake and performance by goats fed coffee pulp-based diets followed by a commercial concentrate 14(1986)1
- Abate, A., see Kategile, J.A. 20(1988)97
- Abate, A.N. and Gomez, M.
Substitution of finger millet (*Eleusine coracana*) and bulrush millet (*Pennisetum typhoides*) for maize in broiler feeds 10(1983/1984)291
- Abd El Ati, N., see Abel, H.J. 19(1988)111
- Abd El Hafiz, G., see Abel, H.J. 19(1988)111
- Abd El-Razek, M., see Abel, H.J. 19(1988)111
- Abdel-Raheim, F.M., see El-Tayeb, A.E. 22(1988)333
- Abdullah, N. and Hutagalung, R.I.
Rumen fermentation, urease activity and performance of cattle given palm kernel cake-based diet 20(1988)79
- Abdullah, N., see Ho, Y.W. 22(1988)161
- Abel, H.J., Abd El Ati, N., Abd El Hafiz, G., Abd El-Razek, M., El-Din, T. and El Shazly, K.
Investigations on the nutritional value for ruminants of various fats 19(1988)111
- Abidin, Z. and Kempton, T.J.
Effects of treatment of barley straw with anhydrous ammonia and supplementation with heat-treated protein meals on feed intake and liveweight performance of growing lambs 6(1981)145
- Abouheif, M.A., Basmaeil, S., Metwally, H. and Masoud, S.
Chemical preparation of NaOH-keratin hydrolysate for improving the nutritive value of wheat straw 13(1985)215
- Abrams, S.M.
Sources of error in predicting digestible dry matter from the acid-detergent fiber content of forages 21(1988)205
- Acamovic, T., see D'Mello, J.P.F. 7(1982)247
- Adams, K.L. and Jensen, A.H.
Effect of processing on the utilization by young pigs of the fat in soya beans and sunflower seeds 12(1984/1985)267
- Adams, K.L. and Jensen, A.H.
Effect of dietary protein and fat levels on the utilization of the fat in sunflower seeds by the young pig 13(1985)159
- Adams, K.L. and Jensen, A.H.
High-fat maize in diets for pigs and sows 17(1987)201
- Adamson, A.H., see Ibbotson, C.F. 10(1983/1984)223
- Adamson, A.H., see Givens, D.I. 19(1988)173
- Adamson, A.H., see Givens, D.I. 24(1989)27
- Adamu, A.M., Russell, J.R. and Trenkle, A.
Effects of ruminal degradability of the protein supplement on the utilization of maize-stover silage by growing beef cattle 20(1988)241
- Adamu, A.M., Russell, J.R., McGilliard, A.D. and Trenkle, A.
Effects of added dietary urea on the utilization of maize stover silage by growing beef cattle 22(1988)227
- Adebowale, E.A.
Organic waste ash as possible source of alkali for animal feed treatment 13(1985)237
- Adegbola, A.A., see Smith, O.B. 13(1985)249
- Adegbola, A.A., see Smith, O.B. 20(1988)189
- Adeneye, J.A.
A note on the nutrient and mineral composition of *Leucaena leucocephala* in Western Nigeria 4(1979)221
- Aderibigbe, A.O., see Razzaque, M.A. 14(1986)265
- Aderibigbe, A.O. and Church, D.C.
Evaluation of cardboard and dried poultry waste as feed ingredients for ruminants 18(1987)209
- Aduku, A.O., see Okoh, P.N. 7(1982)359
- Aerts, J.V., see Carlier, L.A. 1(1976)607
- Aerts, J.V., see Boucque, Ch.V. 1(1976)643
- Aerts, J.V., de Brabander, D.L., Cottyn, B.G. and Buysse, F.X.
Comparison of laboratory methods for predicting the organic matter digestibility of forages 2(1977)337
- Aerts, J.V., De Brabander, D.L., Cottyn, B.G., Buysse, F.X., Carlier, L.A. and Moermans, R.J.
Some remarks on the analytical procedure of Van Soest for the prediction of forage digestibility 3(1978)309
- Aerts, J.V., see Cottyn, B.G. 11(1984)137
- Aerts, J.V., De Boever, J.L., Cottyn, B.G., De Brabander, D.L. and Buysse, F.X.
Comparative digestibility of feedstuffs by sheep and cows 12(1984/1985)47
- Aguilera, A., see Alcántara, E. 23(1989)323
- Aguilera, J.F., see Molina, E. 9(1983)1
- Aguilera, J.F., Molina, E. and Prieto, C.
Digestibility and energy value of sweet lupin seed (*Lupinus albus* var. multolupa) in pigs 12(1984/1985)171
- Aguilera, J.F., Prieto, C., Fonolla, J. and Gil, F.
Protein and energy utilisation in rats of diets based on lupin seed (*Lupinus albus* var. multolupa) 15(1986)33
- Aherne, F.X. and Lewis, A.J.
The nutritive value of tower rapeseed meal for swine 3(1978)235
- Aherne, F.X., see Baidoo, S.K. 18(1987)37
- Aherne, F.X., see Baidoo, S.K. 18(1987)45
- Aherne, F.X., see Gibson, D.M. 19(1988)203
- Aii, T. and Stobbs, the late T.H.
Solubility of the protein of tropical pasture species and the rate of its digestion in the rumen 5(1980)183
- Aitken, W.M., see Wright, D.E. 2(1977)93

- Akin, D.E.
Association of rumen fungi with various forage grasses 16(1986/1987)273
- Akin, D.E., Rigsby, L.L., Theodorou, M.K. and Hartley, R.D.
Population changes of fibrolytic rumen bacteria in the presence of phenolic acids and plant extracts 19(1988)261
- Akin, D.E.
Biological structure of lignocellulose and its degradation in the rumen 21(1988)295
- Akin, D.E., Gordon, G.L.R. and Rigsby, L.L.
Comparative fiber degradation by mixed rumen fungi from Australian and U.S.A. cattle 23(1989)305
- Akin, D.E., see Horn, G.W. 24(1989)201
- Al-Athari, A.K. and Guenter, W.
Nutritional value of triticale (Carman) for broiler diets 22(1988)119
- Al-Athari, A.K. and Guenter, W.
The effect of fat level and type on the utilization of triticale (cultivar Carman) by broiler chicks 22(1988)273
- Al-Awadi, A., see Ilian, M.A. 19(1988)327
- Al-Nasser, A., see Razaque, M.A. 14(1986)265
- Alawa, J.P. and Owen, E.
The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149
- Alawa, J.P. and Owen, E.
The effect of supplementing grass silage with sodium hydroxide-treated chopped or milled straw on intake and digestibility by wether sheep 12(1984/1985)215
- Alawa, J.P., Fishwick, G. and Hemingway, R.G.
Fresh and dried brewer's grains as protein supplements to barley straw diets given to pregnant beef cows 19(1988)33
- Alcántara, E., Aguilera, A., Elliott, R. and Shimada, A.
Fermentation and utilization by lambs of sugarcane harvested fresh and ensiled with and without NaOH. 4. Ruminant kinetics 23(1989)323
- Alexander, J.C., see Gregory, K.F. 2(1977)7
- Alexander, J.C., see Santos, J. 8(1983)313
- Alhassan, W.S., Bello, S.A. and Obilana, A.B.
Yield and potential feeding value of straws of grain sorghum cultivars developed in Nigeria 17(1987)285
- Ali, O.M., see Mohammed, T.A. 22(1988)131
- Alibes, X., Muñoz, F. and Rodriguez, J.
Feeding value of apple pomace silage for sheep 11(1984)189
- Alibes, X., Maestre, M.R., Muñoz, F., Combellas, J. and Rodriguez, J.
Nutritive value of almond hulls for sheep 8(1983)63
- Alibes, X., Muñoz, F. and Faci, R.
Anhydrous ammonia-treated cereal straw for animal feeding. Some results from the Mediterranean area 10(1983/1984)239
- Aliu, Y.O., see Kumaresan, A. 11(1984)45
- Allan, S.J., see Partridge, G.G. 13(1985)299
- Allee, G.L., see Pollmann, D.S. 8(1983)99
- Alli, I., Baker, B.E. and Garcia, G.
Studies on the fermentation of chopped sugarcane 7(1982)411
- Alli, I., Fairbairn, R., Baker, B.E. and Garcia, G.
The effects of ammonia on the fermentation of chopped sugarcane 9(1983)291
- Allison, M.J.
Characterization of the flora of the large bowel of pigs: a status report 23(1989)79
- Alumot, E. and Mandel, E.
Long term feeding of concentrate fumigated with carbon tetrachloride to sheep 2(1977)77
- Aluyi, H.S.A., see Aregheore, E.M. 20(1988)233
- Alvira, P., see Rebolé, A. 16(1986/1987)89
- Alvira, P., see Rebolé, A. 22(1988)173
- Alzueta, C. and Gonzalez, G.
Yield, composition and nutritive value of a leaf protein concentrate obtained by fractionating the shelled aerial part of *Pisum sativum* L. 8(1983)155
- Alzueta, C., see González, G. 20(1988)177
- Áman, P., see Graham, H. 10(1983/1984)199
- Áman, P., see Graham, H. 12(1984/1985)195
- Áman, P., see Hesselman, K. 15(1986)83
- Áman, P. and Graham, H.
Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro degradability during maturation 17(1987)15
- Áman, P., see Graham, H. 17(1987)33
- Áman, P., see Pettersson, D. 17(1987)145
- Áman, P., see Nordkvist, E. 17(1987)295
- Áman, P., see Pettersson, D. 20(1988)313
- Áman, P., see Theander, O. 23(1989)205
- Áman, P., see Narasimhalu, P. 24(1989)159
- Amaning-Kwarteng, K., Kellaway, R.C., Spragg, J.C. and Kirby, A.C.
Relative intakes, digestibility and bacterial protein synthesis by sheep and cattle fed high-roughage diets 16(1986/1987)75
- Ammerman, C.B., see Potter, S.G. 12(1984/1985)275
- Anderson, D.H., see Henderson, A.R. 7(1982)303
- Anderson, D.M., see Bowland, J.P. 12(1984/1985)75
- Anderson, P., see Zervas, G. 21(1988)23
- Anderson, R.
Effect of prolonged wilting in poor conditions on the fermentation quality, metabolisability and net energy value of silage given to sheep 12(1984/1985)109
- Andries, J.L., Buysse, F.X., de Brabander, D.L. and Cottyn, B.G.
Isoacids in ruminant nutrition: their role in ruminal and intermediary metabolism and

- possible influences on performances - a review 18(1987)169
- Andries, J.I., see De Boever, J.L. 19(1988)247
- Andrieu, J.
Agronomic factors affecting the growth and composition of the maize plant 1(1976)263
- Andrieu, J.
Factors influencing the composition and nutritive value of ensiled whole-crop maize 1(1976)381
- Angemar, Y., see Holzer, Z. 12(1984/1985)253
- Anglesea, J.D. and Jackson, A.J.
Thiaminase activity in fish silage and moist fish feed 13(1985)39
- Anonymous
Requirements for future research 1(1976)555
- Antongiovanni, M., Giorgetti, A. and Franci, O.
Amino acid composition of two varieties of sorghum grain 5(1980)169
- Anwar, A., see Mohamed, K. 11(1984)199
- Aregheore, E.M., Job, T.A. and Aluyi, H.S.A.
The maize replacement value of cassava flour in rations for growing ewe lambs 20(1988)233
- Argenteria, A., see Caja, G. 2(1977)267
- Arieli, A. and Werner, D.
A comparison between fermentation heat of forages and organic matter digestibility determined by in vitro incubation with rumen fluid 23(1989)333
- Armstrong, D.G., see McMeniman, N.P. 2(1977)255
- Armstrong, D.G., see Rooke, J.A. 13(1985)255
- Armstrong, D.G., see Rooke, J.A. 13(1985)269
- Armstrong, D.G., see Lee, N.-H. 15(1986)301
- Arora, S.K., see Das, B. 4(1979)17
- Arp, S.C., see Schaefer, D.M. 25(1989)23
- Asahida, Y., see Sekine, J. 25(1989)207
- Ash, A.J., see McMeniman, N.P. 19(1988)43
- Ashbell, G., see Levy, D. 16(1986/1987)63
- Ashbell, G., see Weinberg, Z.G. 20(1988)335
- Ashes, J.R. and Peck, N.J.
A simple device for dehulling seeds and grain 3(1978)109
- Ashes, J.R., see Farrell, D.J. 9(1983)99
- Asplund, J.M., see Bowman, J.G.P. 20(1988)19
- Asplund, J.M., see Bowman, J.G.P. 20(1988)33
- Atinkpahoun, H., see Szyliet, O. 3(1978)73
- Atkinson, T., see Livingstone, R.M. 4(1979)295
- Atkinson, T., see Livingstone, R.M. 5(1980)69
- Atuahene, C.C., Donkoh, A. and Swatson, H.
Oil palm slurry (OPS) as a partial replacement for maize in the diets of broiler chickens 17(1987)157
- Atuahene, C.C., see Donkoh, A. 22(1988)139
- Aufriere, J. and Michalet-Doreau, B.
Comparison of methods for predicting digestibility of feeds 20(1988)203
- Auld, D.L., see Thomas, V.M. 11(1984)269
- Aumaitre, A., see Le Dividich, J. 3(1978)15
- Aumaitre, A., see Perez, J.M. 4(1979)109
- Aumaitre, A., Bourdon, D., Peinau, J. and Bengala Freire, J.
Effect of graded levels of raw and processed rapeseed on feed digestibility and nutrient utilization in young pigs 24(1989)275
- Awadel-Kareim, A.M., see El-Tayeb, A.E. 22(1988)333
- Ayyaluswami, P., see Thanu, K. 9(1983)263
- ## B
- Babatunde, G.M., Fetuga, B.L., Odegunmi, J.O. and Oduye, O.O.
The effects of air dried poultry manure on performance, organ weights and haematological measurements of weaner pigs reared in the tropics 4(1979)23
- Babatunde, G.M. and Pond, W.G.
Nutritive value of Nigerian rubber seed (*Hevea brasiliensis*) meal and oil. III. Performance characteristics, relative organ weights, hematocrit and plasma metabolites of growing female rats fed corn diets containing rubber seed meal, soya bean meal or casein 20(1988)125
- Bačvanski, S.
Maize grain or ears in concentrate diets for young fattening bulls 1(1976)393
- Badir, N.A., see El Khidir, O.A. 9(1983)301
- Badve, V.C., see Rangnekar, D.V. 7(1982)61
- Baidoo, S.K., Aherne, F.X., Mitaru, B.N. and Blair, R.
Canola meal as a protein supplement for growing-finishing pigs 18(1987)37
- Baidoo, S.K., Mitaru, B.N., Aherne, F.X. and Blair, R.
The nutritive value of canola meal for early-weaned pigs 18(1987)45
- Baird, B.A., see Livingstone, R.M. 4(1979)295
- Baird, B.A., see Livingstone, R.M. 5(1980)69
- Baker, B.E., see Alli, I. 7(1982)411
- Baker, B.E., see Alli, I. 9(1983)291
- Baker, E.C., see Lynch, G.L. 14(1986)127
- Bakshi, M.P.S., see Langar, P.N. 13(1985)103
- Balogopal, R., see Natanam, R. 25(1989)201
- Baldwin, A., see O'Shea, J. 15(1986)13
- Ballance, G.M., see Campbell, G.L. 16(1986/1987)243
- Balogun, O.O. and Fetuga, B.L.
Lysine requirements of imported European pigs in the humid lowland tropics from 23 to 62 kg liveweight 7(1982)233
- Balogun, O.O. and Fetuga, B.L.A.
Lysine requirement of weanling European pigs fed on yellow maize and groundnut cake diets in the humid lowland tropics 8(1983)197
- Bandyk, C.A., see Pollmann, D.S. 11(1984)261
- Bar-Tsur, A., see Levy, D. 16(1986/1987)63

- Baran, M., Bođa, K. and Siroka, P.
The effect of monensin on rumen fermentation in sheep fed on all-roughage and barley/roughage diets 15(1986)7
- Barber, R.S., Braude, R. and Mitchell, K.G.
The value of 'Pekilo protein' for growing pigs 2(1977)161
- Barber, R.S., Braude, R., Mitchell, K.G. and Pittman, R.J.
The nutritive value of liquid whey, either sour or sweet, when given in restricted amounts to the growing pig 3(1978)163
- Barber, R.S., Braude, R., Hosking, Z.D. and Mitchell, K.G.
Olaquinox as performance-promoting feed additive for growing pigs 4(1979)117
- Barber, R.S., Braude, R., Mitchell, K.G., Partridge, I.G. and Pittman, R.J.
Value of lucerne juice and grass juice as sources of protein for the growing pig 4 (1979)233
- Barber, R.S., Braude, R., Mitchell, K.G., Partridge, I.G. and Pittman, R.J.
Value of freshly produced lucerne juice as a source of supplemental protein for the growing pig 5(1980)215
- Barber, R.S., Braude, R., Mitchell, K.G. and Partridge, I.G.
Lucerne juice as a protein supplement for growing pigs: effects of mineral content of the diet and of the water supply 6(1981)35
- Barber, W.P., see Givens, D.I. 16(1986/1987) 311
- Barge, M.T., see Pace, V. 11(1984)125
- Barker, D.J., see May, P.J. 12(1984/1985)57
- Barnes, B.J. and Ørskov, E.R.
Utilization of alkali-treated grain. 2. Utilization by steers of diets based on hay or straw and mixed with either NaOH-treated or rolled barley 6(1981)347
- Barnes, B.J., see Ørskov, E.R. 6(1981)355
- Barnes, D.A., see Brown, D.L. 18(1987)247
- Barnes, R.J.
Near infra-red spectra of ammonia-treated straw and of isolated cell walls 21(1988)209
- Barnett, C.W. and Batterham, E.S.
Lupinus angustifolius cv. Unicrop as a protein- and energy-source for weaner pigs 6(1981)27
- Barnett, R.D., see Myer, R.O. 22(1988)217
- Barro, C., see González, G. 20(1988)177
- Barry, J.L., see Cherbut, C. 20(1988)327
- Barry, J.L., see Champ, M. 23(1989)195
- Barry, P., see Grenet, E. 19(1988)25
- Bartling, R.R., see Umunna, N.N. 12(1984/1985)151
- Bartling, R.R., see Umunna, N.N. 14(1986)161
- Barton, F.E., see Coelho, M. 20(1988)219
- Barton, G.M., see Hunt, J.R. 3(1978)63
- Barton II, F.E., Wolsink, J.H. and Vedder, H.M.
Near infrared reflectance spectroscopy of untreated and ammoniated barley straw 15 (1986)189
- Barton II, F.E.
Chemistry of lignocellulose: methods of analysis and consequences of structure 21 (1988)279
- Bartsch, B.D., see Valentine, S.C. 16(1986/1987)143
- Bartsch, B.D., see Valentine, S.C. 16(1986/1987)261
- Basmaeli, S., see Abouheif, M.A. 13(1985)215
- Bass, J.M., Parkins, J.J. and Fishwick, G.
The effect of calcium hydroxide treatment on the digestibility of chopped oat straw supplemented with a solution containing urea, calcium, phosphorus, sodium, trace elements and vitamins 7(1982)93
- Batchelder, D.G., see Horn, G.W. 8(1983)35
- Batterham, E.S., see Barnett, C.W. 6(1981)27
- Batterham, E.S., Gorman, T.B.S. and Chvojka, R.
Nutritional value and mercury content of fish silage for growing pigs 9(1983)169
- Batterham, E.S. and Watson, C.
Tryptophan content of feeds, limitations in diets and requirement for growing pigs 13 (1985)171
- Baucells, M.D., see Gasa, J. 25(1989)67
- Bazylak, G., see Masłowska, J. 13(1985)227
- Beames, R.M., see Pringle, D. 9(1983)89
- Beames, R.M., Tait, R.M., Pringle, D.B. and Owen, B.D.
The nutritive value of barley for monogastric animals and for ruminants as affected by artificial drying and alkali and anaerobic storage 15(1986)285
- Beard, J., see Clark, J. 2(1977)153
- Beaumont, P., see Durand, M. 21(1988)197
- Becker, H.N., see Potter, S.G. 12(1984/1985) 275
- Becker, K., see Ohlde, G. 7(1982)191
- Beever, D.E.
The digestion and metabolism of the energy and protein of maize 1(1976)455
- Behnke, K.C., see Skoch, E.R. 6(1981)83
- Behnke, K.C., see Pollmann, D.S. 8(1983)99
- Bell, J.M., see Thacker, P.A. 19(1988)191
- Bell, J.M. and Keith, M.O.
Factors affecting the digestibility by pigs of energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253
- Bell, S.L., see Neill, A.R. 9(1983)71
- Bell, S.L., see Rooke, J.A. 13(1985)269
- Bello, S.A., see Alhassan, W.S. 17(1987)285
- Belyavin, C.G.
Effects of the addition of silica to the diet of the laying hen, with particular reference to egg-shell quality 6(1981)157
- Belyea, R.L., Martz, F.A., Madhisetty, M. and Davis, T.P.
Intake, digestibility and energy utilization of corrugated paper diets 17(1987)57
- Ben-Ghedalia, D. and Miron, J.
The response of wheat straw varieties to mild sulphur dioxide treatment 10(1983/

- 1984)269
- Ben-Ghedalia, D. and Rubinstein, A.
The response of screened manure fibre to ozone and sodium hydroxide treatments 15 (1986)47
- Ben-Ghedalia, D. and Solomon, R.
The effect of dietary barley on carbohydrate digestibility of sulphur dioxide-treated wheat straw by sheep 18(1987)55
- Ben-Ghedalia, D., Miron, J., Est, Y. and Yosef, E.
SO₂ treatment for converting straw into a concentrate-like feed: a growth study with lambs 19(1988)219
- Ben-Ghedalia, D. and Solomon, R.
Amino acids flow to and absorption from the small intestine of sheep fed barley and sulfur dioxide-treated straw at different ratios 22(1988)147
- Ben-Ghedalia, D. and Yosef, E.
SO₂-treated straw as a silage additive: fermentation data on lucerne with particular reference to protein degradation 22(1988)247
- Ben-Ghedalia, D. and Yosef, E.
SO₂-treated straw as a silage additive: the participation of soluble and cell wall monosaccharide residues in lucerne silage fermentation 22(1988)255
- Ben-Ghedalia, D., Yosef, E., Miron, J. and Est, Y.
The effects of starch and pectin-rich diets on quantitative aspects of digestion in sheep 24 (1989)289
- Benevenga, N.J., see Cieslak, D.G. 14(1986)171
- Bengala Freire, J., see Aumaitre, A. 24(1989)275
- Berger, L.L., see Mora, M.I. 9(1983)205
- Berger, L.L., see Lynch, G.L. 14(1986)127
- Berger, L.L., see Lewis, S.M. 17(1987)179
- Berkovitz, S., see Levy, D. 16(1986/1987)63
- Bertrand, D., see Sauvage, D. 13(1985)7
- Bertrand, D., see Robert, P. 16(1986/1987)215
- Berzins, R., see Ha, J.K. 14(1986)117
- Bhargava, P.K., Ørskov, E.R. and Walli, T.K.
Effect of soaking, ensilage and hydrogen peroxide treatment of barley straw on rumen degradability 22(1988)295
- Bickel, H., see Wyss, U. 20(1988)325
- Bienfait, J.M., see Istasse, L. 25(1989)1
- Binder, S.F., see Skoch, E.R. 6(1981)83
- Biston, L., see Istasse, L. 25(1989)1
- Blair, R., see Baidoo, S.K. 18(1987)37
- Blair, R., see Baidoo, S.K. 18(1987)45
- Blanchart, G., see Laurent, F. 13(1985)131
- Blanco, J.L.
Selection objectives for maximum yields of feed with hybrid maize 1(1976)245
- Bland, B., see Panigrahi, S. 22(1988)285
- Boda, K., see Baran, M. 15(1986)7
- Boekholt, H.A., see Van Es, A.J.S. 1(1976)481
- Boer, H., see Tamminga, S. 25(1989)89
- Bokori, J., see Hegedüs, M. 9(1983)79
- Boling, J.A., see Pendulum, L.C. 2(1977)225
- Bolsen, K.K., Tetlow, R.M. and Wilson, R.F.
The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37
- Bolsen, K.K., see Tetlow, R.M. 12(1984/1985)91
- Bolsen, K.K., see Hinds, M.A. 12(1984/1985)205
- Bolt, M.J., see Slabbert, N. 21(1988)31
- Borgida, L.P., see Szylit, O. 3(1978)73
- Borhami, B.E.A. and Sundstøl, F.
Studies on ammonia-treated straw. I. The effects of type and level of ammonia, moisture content and treatment time on the digestibility in vitro and enzyme soluble organic matter of oat straw 7(1982)45
- Borhami, B.E.A., Sundstøl, F. and Garmo, T.H.
Studies on ammonia-treated straw. II. Fixation of ammonia in treated straw by spraying with acids 7(1982)53
- Bothast, R.J., see Montgomery, R.R. 5(1980)337
- Boucque, Ch.V., see Cottyn, B.G. 1(1976)167
- Boucque, Ch.V., Cottyn, B.G. and Buysse, F.X.
Maize silage with or without NPN, dehydrated whole-crop maize pellets or high moisture maize grain for finishing bulls 1 (1976)347
- Boucque, Ch.V., Cottyn, B.G., Aerts, J.V. and Buysse, F.X.
Dried sugar beet pulp as a high energy feed for beef cattle 1(1976)643
- Bouqué, Ch.V., see Fiems, L.O. 8(1983)237
- Bouqué, Ch.V., see Fiems, L.O. 16(1986/1987)287
- Bouqué, Ch.V., Fiems, L.O., Cottyn, B.G., Casteels, M. and Buysse, F.X.
Monensin-sodium as a performance-promoting additive for fattening bulls and its impact on carcass and meat quality characteristics 7(1982)401
- Bourdon, D., see Aumaitre, A. 24(1989)275
- Bowland, J.P., Anderson, D.M. and Thacker, P.A.
Chemical composition and nutritive value of Canadian winter wheats (*Triticum aestivum* L.) given to laboratory rats 12(1984/1985)75
- Bowman, D.K., see Koeln, L.L. 11(1984)181
- Bowman, J.G.P. and Asplund, J.M.
Evaluation of mixed lucerne and caucasian bluestem hay diets fed to sheep 20(1988)19
- Bowman, J.G.P. and Asplund, J.M.
Nitrogen utilization, ruminal fermentation and abomasal nitrogen flow in sheep fed caucasian bluestem hay supplemented with lucerne or urea 20(1988)33
- Boza, J., see Molina, E. 9(1983)1

- Brahmakshatriva, R.D., see Singh, K.S. 6 (1981)63
- Braude, R., see Barber, R.S. 2(1977)161
- Braude, R., see Barber, R.S. 3(1978)163
- Braude, R., see Barber, R.S. 4(1979)117
- Braude, R., see Barber, R.S. 4(1979)233
- Braude, R., see Barber, R.S. 5(1980)215
- Braude, R., see Barber, R.S. 6(1981)35
- Bremel, D.H., see Teh, T.H. 17(1987)257
- Brethour, J., see Hinds, M.A. 12(1984/1985) 205
- Brett, P.A., see Rooke, J.A. 13(1985)255
- Brewer, A., see Williams, P.E.V. 11(1984)103
- Brewer, A., see Williams, P.E.V. 11(1984)115
- Brice, R.E., see Morrison, I.M. 10(1983/1984) 229
- Brice, R.E., see Robinson, P.H. 18(1987)75
- Brigstocke, T.D.A., see Wilson, P.N. 6(1981)1
- Brophy, P.O., see Caffrey, P.J. 19(1988)231
- Broster, W.H., see Smith, T. 10(1983/1984) 189
- Broster, W.H., see Smith, T. 14(1986)29
- Brotz, P.G., see Schaefer, D.M. 25(1989)23
- Brown, D.L. and Chavalim, E.
Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistani napier grass), *Pennisetum* sp. (bana grass), *Impomea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985) 1
- Brown, D.L., Barnes, D.A., Rezende, S.A. and Klasing, K.C.
Yield, composition and feeding value of irrigated *Sesbania sesban* var. *Nubia* leaves harvested at latitude 38°N during a Mediterranean summer 18(1987)247
- Brown, D.L., see Shqueir, A.A. 25(1989)137
- Brunnen, J.M., see Givens, D.I. 18(1987)83
- Brunnen, J.M., see Clark, P. 18(1987)225
- Bruyneel, B., see Vandevoorde, L. 21(1988)99
- Bryant, K.L., see Harper, A.F. 8(1983)69
- Buchanan-Smith, J.G., see Oji, U.I. 4(1979) 187
- Bunting, L.D., McCarthy, D.B., Richardson, C.R. and Tock, R.W.
Pilot plant production of sulphur dioxide-treated mesquite wood. II. Digestibility and feedlot performance by lambs 22(1988)61
- Burke, A.B., see Walldrop, P.W. 3(1978)329
- Bussell, W.D., see Walldrop, P.W. 3(1978)329
- Butler, E.J., see Fenwick, G.R. 6(1981)421
- Butler, E.J., see Pearson, A.W. 8(1983)119
- Butler, K.L., see Minson, D.J. 9(1983)221
- Butterworth, M.H., see Mosi, A.K. 12(1984/1985)241
- Butterworth, M.H. and Mosi, A.
The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16 (1986/1987)99
- Buxton, D.R., see Russell, J.R. 13(1985)69
- Buyse, F.X., see Cottyn, B.G. 1(1976)167
- Buyse, F.X., see Boucque, Ch.V. 1(1976)347
- Buyse, F.X., see Boucque, Ch.V. 1(1976)643
- Buyse, F.X., see Aerts, J.V. 2(1977)337
- Buyse, F.X., see Aerts, J.V. 3(1978)309
- Buyse, F.X., see Bouqué, Ch.V. 7(1982)401
- Buyse, F.X., see Fiems, L.O. 8(1983)237
- Buyse, F.X., see Cottyn, B.G. 11(1984)137
- Buyse, F.X., see Aerts, J.V. 12(1984/1985)47
- Buyse, F.X., see De Boever, J.L. 14(1986)203
- Buyse, F.X., see Fiems, L.O. 16(1986/1987) 287
- Buyse, F.X., see Andries, J.I. 18(1987)169
- Buyse, F.X., see De Boever, J.L. 19(1988)247
- ## C
- Caballero, R., see Treviño, J. 16(1986/1987) 305
- Caffrey, P.J., Miller, C., Brophy, P.O. and Kelleher, D.L.
The effects of method of processing of starters, tallow inclusion and roughage supplementation on the performance of early-weaned calves 19(1988)231
- Caja, G., Galvez, J.F., Argamenteria, A. and Ciria, J.
Inhibition of ruminal deamination in vitro by formaldehyde treatment of sunflower-seed, soya bean and fish meals: response curves to protective treatment 2(1977)267
- Campbell, D.R., see Noland, P.R. 2(1977)219
- Campbell, D.R., see Sharp, R.N. 2(1977)327
- Campbell, D.R., see Noland, P.R. 5(1980)51
- Campbell, G.L., Sosulski, F.W., Classen, H.L. and Ballance, G.M.
Nutritive value of irradiated and β -glucanase-treated wild oat groats (*Avena fatua* L.) for broiler chickens 16(1986/1987)243
- Campbell, G.L., see Thacker, P.A. 19(1988) 191
- Campbell, G.L., see Edney, M.J. 25(1989)193
- Campbell, R.G., see King, R.H. 3(1978)191
- Campher, J.P., see Slabbert, N. 21(1988)31
- Campling, R.C., see Jewell, S.N. 14(1986)81
- Capper, A.L., see Edwards, S.A. 15(1986)129
- Capper, A.L.
Fungal contamination of hydroponic forage 20(1988)163
- Capper, B.S., see Machin, D.H. 11(1984)247
- Capper, B.S.
Genetic variation in the feeding value of cereal straw 21(1988)127
- Carlaw, P.M., see Panigrahi, S. 22(1988)285
- Carlier, L.A., Cottyn, B.G. and Aerts, J.V.
Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607
- Carlier, L.A., see Aerts, J.V. 3(1978)309
- Carlos, G., see Zervas, G. 21(1988)23
- Carlsson, R., Hanczakowski, P. and Kaptur, T.
The quality of the green fraction of leaf pro-

- tein concentrate from *Chenopodium quinoa* Willd. grown at different levels of fertilizer nitrogen 11(1984)239
- Carr, J.R., see Pearson, G. 1(1976)631
- Carr, J.R., see Pearson, G. 2(1977)49
- Carré, B., see Lacassagne, L. 20(1988)59
- Carter, P.M., see Wood, J.F. 9(1983)307
- Carter, P.M., see Machin, D.H. 11(1984)247
- Carter, R.R., see Leibholz, J. 15(1986)141
- Casper, H.H., see Sanderson, M.A. 14(1986)221
- Castaing, J., see Gatel, F. 20(1988)145
- Casteels, M., see Bouqué, Ch.V. 7(1982)401
- Castrillo, C., see Gasa, J. 25(1989)67
- Catalano, A., see Russo, V. 1(1976)25
- Cavani, C., see Manfredini, M. 5(1980)233
- Cayley, G.R., see Lord, K.A. 6(1981)73
- Cayley, G.R., see Lacey, J. 8(1983)323
- Cayley, G.R., see Lord, K.A. 7(1982)287
- Cayley, G.R., see Lacey, J. 8(1983)283
- Centeno, C., see Treviño, J. 16(1986/1987)305
- Cervera, C., Fernandez-Carmona, J. and Marti, J.
Effect of urea on the ensiling process of orange pulp 12(1984/1985)233
- Chahal, D.S., see Moo-Young, M. 4(1979)199
- Chahal, D.S., see Moo-Young, M. 5(1980)175
- Champ, M., Barry, J.L., Hoebler, C. and Delort-Laval, J.
Digestion and fermentation pattern of various dietary fiber sources in the rat 23(1989)195
- Chan, C., see Farrell, D.J. 9(1983)49
- Chandra, S., Prasad, D.A. and Krishna, N.
Effect of sodium hydroxide treatment and/or extrusion cooking on the nutritive value of peanut hulls 12(1984/1985)187
- Chandrasekaran, D., see Subramanian, P.R. 20(1988)343
- Chandrasekaran, D., Kadirvel, R. and Viswanathan, K.
Nutritive value of pungan (*Pongamia glabra* Vent) cake for sheep 22(1988)321
- Chandrasekaran, D., see Viswanathan, K. 22(1988)327
- Charnley, E. and Greenhalgh, J.F.D.
Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19
- Chase Jr., C.C. and Hibberd, C.A.
Effect of level and frequency of maize supplementation on the utilization of low-quality grass hay by beef cows 24(1989)129
- Chavalim, E., see Brown, D.L. 13(1985)1
- Chavez, E.R., Touchburn, S.P. and Moo-Young, M.
Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. I. Composition and nutritive evaluation 22(1988)1
- Chavez, E.R., Touchburn, S.P. and Moo-Young, M.
Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. II. Effect of method of drying and response to amino acid supplementation 22(1988)13
- Chavez, E.R., Touchburn, S.P. and Moo-Young, M.
Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. III. Nutritive value for chicks and piglets 22(1988)23
- Cheng, K.-J., Stewart, C.S., Dinsdale, D. and Costerton, J.W.
Electron microscopy of bacteria involved in the digestion of plant cell walls 10(1983/1984)93
- Cherbut, C., Barry, J.L., Wyers, M. and Delort-Laval, J.
Effect of the nature of dietary fibre on transit time and faecal excretion in the growing pig 20(1988)327
- Cherney, J.H., Johnson, K.D., Tuite, J. and Volenc, J.J.
Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and stored at different moisture contents 17(1987)45
- Chesson, A., see Gordon, A.H. 8(1983)147
- Chesson, A.
Lignin-polysaccharide complexes of the plant cell wall and their effect on microbial degradation in the rumen 21(1988)219
- Chiericato, G.M., see Parigi-Bini, R. 1(1976)301
- Choice, A., see Farrell, D.J. 9(1983)99
- Christensen, D.A., see Horton, G.M.J. 7(1982)1
- Christensen, D.A., see Mann, M.E. 21(1988)57
- Christison, G.I., see Mandell, I.B. 20(1988)111
- Christon, R., see Le Dividich, J. 3(1978)15
- Church, D.C., see Razzaque, M.A. 14(1986)265
- Church, D.C., see Aderibigbe, A.O. 18(1987)209
- Chvojka, R., see Batterham, E.S. 9(1983)169
- Cieslak, D.G., Benevenga, N.J. and Grummer, R.H.
The evaluation of fresh sweet liquid whey as a protein supplement to maize for growing pigs 14(1986)171
- Ciria, J., see Caja, G. 2(1977)267
- Clark, C.F.S., see Ørskov, E.R. 7(1982)135
- Clark, J. and Beard, J.
Prediction of the digestibility of ruminant feeds from their solubility in enzyme solutions 2(1977)153
- Clark, P., Givens, D.I. and Brunnen, J.M.
The chemical composition, digestibility and energy value of fodder-beet roots 18(1987)225
- Classen, H.L., see Campbell, G.L. 16(1986/1987)243
- Classen, H.L., see Thacker, P.A. 19(1988)191
- Classen, H.L., see Edney, M.J. 25(1989)193
- Cobby, J.M., see Mason, V.C. 19(1988)159
- Cobby, J.M., see Givens, D.I. 19(1988)173
- Cobic, T.
Effect of weight at slaughter on the efficiency of fattening young cattle 1(1976)505

- Cockburn, J.E., see Mason, V.C. 24(1989)313
 Cockburn, J.H.S., see Morgan, C.A. 11(1984) 11
- Coelho, M., Hembry, F.G., Barton, F.E. and Saxton, A.M.
 A comparison of microbial, enzymatic, chemical and near-infrared reflectance spectroscopy methods in forage evaluation 20 (1988)219
- Cohen, O., see Silanikove, N. 20(1988)87
 Cohen, R.D.H., see Mann, M.E. 21(1988)57
 Cole, D.J.A., see Green, S. 21(1988)43
 Collins, M.
 Composition and fibre digestion in morphological components of an alfalfa-timothy sward 19(1988)135
- Combellas, J., see Alibés, X. 8(1983)63
 Combs, G.E., see Myer, R.O. 22(1988)217
 Compte, X., see Valdivie, M. 7(1982)185
 Connor, J.K., see Moir, K.W. 2(1977)197
 Conrad, J.H., see Paz, E.A. 11(1984)75
 Cook, D.K., see Schaefer, D.M. 25(1989)23
 Cook, J.E., see Woolford, M.K. 3(1978)89
 Cook, J.E., see Woolford, M.K. 8(1983)107
 Cook, J.E., see Mason, V.C. 24(1989)299
 Cook, J.E., see Mason, V.C. 24(1989)313
 Cooper, E.M., see Mason, V.C. 24(1989)299
 Cooper, E.M., see Mason, V.C. 24(1989)313
 Coppock, C.E., Lanham, J.K. and Horner, J.I.
 A review of the nutritive value and utilization of whole cottonseed, cottonseed meal and associated by-products by dairy cattle 18(1987)89
- Corcuera, P., see Penning, P.D. 5(1980)321
 Cornelissen, J.P., see Grimbergen, A.H.M. 13 (1985)117
 Cornell, J.A., see Myer, R.O. 22(1988)217
 Costerton, J.W., see Cheng, K.-J. 10(1983/1984)93
 Cote, M., see Sibbald, I.R. 8(1983)191
 Cottyn, B.G., Boucque, Ch.V. and Buysse, F.X.
 Survey of the use of maize for livestock feeding in Belgium 1(1976)167
 Cottyn, B.G., see Boucque, Ch.V. 1(1976)347
 Cottyn, B.G., see Carlier, L.A. 1(1976)607
 Cottyn, B.G., see Boucque, Ch.V. 1(1976)643
 Cottyn, B.G., see Aerts, J.V. 2(1977)337
 Cottyn, B.G., see Aerts, J.V. 3(1978)309
 Cottyn, B.G., see Bouqué, Ch.V. 7(1982)401
 Cottyn, B.G., see Fiems, L.O. 8(1983)237
 Cottyn, B.G., Aerts, J.V., Vanacker, J.M., Moermans, R.J. and Buysse, F.X.
 The prediction of the net energy value of compound feedstuffs for dairy cattle 11 (1984)137
 Cottyn, B.G., see Aerts, J.V. 12(1984/1985) 47
 Cottyn, B.G., see De Boever, J.L. 14(1986)203
 Cottyn, B.G., see Fiems, L.O. 16(1986/1987) 287
 Cottyn, B.G., see Andries, J.I. 18(1987)169
 Cottyn, B.G., see De Boever, J.L. 19(1988)247
 Cottyn, B.G. and de Boever, J.L.
 Upgrading of straw by ammoniation 21 (1988)287
- Courtin, M.G. and Spoelstra, S.F.
 Counteracting structure loss in pressed sugar beet pulp silage 24(1989)97
 Cowan, P.J. and Michie, W.
 Turkey feeding regimes with different ages of introduction of a choice of foods 4(1979) 125
- Coxworth, E.C., see Kernan, J.A. 6(1981)257
 Coxworth, E.C., see Kernan, J.A. 11(1984)301
 Coxworth, E.C., see Mandell, I.B. 20(1988)111
 Crickenberger, R.G., see Johnson, T.O. 12 (1984/1985)65
 Crickenberger, R.G., see Ortigues, I. 24(1989) 13
- Crofts, F.C., see Kellaway, R.C. 3(1978)201
 Crofts, R.M.J., see Livingstone, R.M. 4(1979) 295
- Croke, B., see Lahore, R. 3(1978)1
 Crooks, P., see Morgan, C.A. 17(1987)81
 Cross, D.L., see Grotheer, M.D. 14(1986)55
 Crowle, W.L., see Kernan, J.A. 11(1984)301
 Čuperlović, M., see Jovanović, M. 2(1977)351
 Curl, C.L., see Pearson, A.W. 8(1983)119
 Curtis, R.F., see Fenwick, G.R. 5(1980)255
 Cuthbert, N.H., see Wilson, P.N. 6(1981)1
- ## D
- da Costa, J.S.
 Survey of the use of maize for livestock feeding in Portugal 1(1976)137
- Damron, B.L., see Paz, E.A. 11(1984)75
 Daniels, L.B., Smith, M.J., Stallcup, O.T. and Rakes, J.M.
 Nutritive value of ensiled broiler litter for cattle 8(1983)19
- Das, B., Arora, S.K. and Luthra, Y.P.
 Variability in structural carbohydrates and in vitro digestibility of forages. 4. Fenugreek (*Trigonella foenum-graecum*) 4(1979)17
- Datta, S.C.
 Estimation of lysine in compounded poultry diets after enzymic digestion 3(1978)57
- Davidson, J., McIntosh, A.D. and Milne, E.
 Note on the lysinoalanine content of alkali-treated barley 7(1982)217
- Davis, T.P., see Belyea, R.L. 17(1987)57
 de Boer, F.
 Use of maize for livestock feeding in the Netherlands 1(1976)177
 De Boever, J.L., see Aerts, J.V. 12(1984/1985) 47
- De Boever, J.L., Cottyn, B.G., Buysse, F.X., Wainman, F.W. and Vanacker, J.M.
 The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14(1986)203
- De Boever, J.L., Cottyn, B.G., Andries, J.I., Buysse, F.X. and Vanacker, J.M.

- The use of a cellulase technique to predict digestibility, metabolizable and net energy of forages 19(1988)247
- de Boever, J.L., see Cottyn, B.G. 21(1988)287
- De Brabander, D.L., see Aerts, J.V. 3(1978)309
- De Brabander, D.L., see Aerts, J.V. 12(1984/1985)47
- de Brabander, D.L., see Aerts, J.V. 2(1977)337
- de Brabander, D.L., see Andries, J.I. 18(1987)169
- De Groote, G., see Huyghebaert, G. 20(1988)45
- de Groote, G., see Ketels, E. 22(1988)105
- de la Cuesta, D., see Gómez, G. 11(1984)57
- De Munter, G., see Huyghebaert, G. 20(1988)45
- de Paeppe, M., see Eeckhout, W. 24(1989)1
- Decuyper, J.A., see Dierick, N.A. 23(1989)141
- Decuyper, J.A., see Vervaeke, I.J. 23(1989)169
- Deinum, B., Steg, A. and Hof, G.
Measurement and prediction of digestibility of forage maize in The Netherlands 10(1983/1984)301
- Del Monte, P., see Russo, V. 1(1976)25
- Delort-Laval, J., see Szylit, O. 3(1978)73
- Delort-Laval, J., see Cherbut, C. 20(1988)327
- Delort-Laval, J., see Champ, M. 23(1989)195
- Deltoro López, J. and Martínez Pascual, J.L.
Influence of the drying process on the composition of brewer's dried grains 6(1981)163
- Deltoro López, J., Fernandez Carmona, J. and Martínez Pascual, J.L.
Evaluation of brewer's dried grains in the diets of laying hens 6(1981)169
- Deltoro López, J. and Fernandez Carmona, J.
Evaluation of brewer's dried grains in the diets of broiler chickens 6(1981)179
- Demarquilly, C., see Robert, P. 16(1986/1987)215
- Demeyer, D.I., see Kayouli, C. 10(1983/1984)165
- Demeyer, D.I., see Jouany, J.P. 21(1988)229
- Demeyer, D.I., see Dierick, N.A. 23(1989)141
- Demeyer, D.I., see Vervaeke, I.J. 23(1989)169
- Demeyer, D.I., see Henderson, C. 23(1989)227
- Dendooven, R., see Kayouli, C. 10(1983/1984)165
- Deolankar, R.P. and Singh, K.S.
Trypsin inhibitor, mineral availability, and performance of broiler chickens fed on diets based on rice bran 4(1979)133
- Deschard, G., see Tetlow, R.M. 18(1987)271
- Deschard, G., Tetlow, R.M. and Mason, V.C.
Treatment of whole-crop cereals with alkali.
3. Voluntary intake and digestibility studies in sheep given immature wheat ensiled with sodium hydroxide, urea or ammonia 18(1987)283
- Deschard, G., Mason, V.C. and Tetlow, R.M.
Treatment of whole-crop cereals with alkali.
4. Voluntary intake and growth in steers given wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55
- Deyoe, C.W., see Skoch, E.R. 6(1981)83
- Dias-da-Silva, A.A. and Sundstøl, F.
Urea as a source of ammonia for improving the nutritive value of wheat straw 14(1986)67
- Dias-da-Silva, A.A., Mascarenhas Ferreira, A. and Guedes, C.V.M.
Effects of moisture level, treatment time and soya bean addition on the nutritive value of urea-treated maize stover 19(1988)67
- Dias-da-Silva, A.A., see Mascarenhas-Ferreira, A. 25(1989)157
- Dierick, N.A., Vervaeke, I.J., Demeyer, D.I. and Decuyper, J.A.
Approach to the energetic importance of fibre digestion in pigs. I. Importance of fermentation in the overall energy supply 23(1989)141
- Dierick, N.A., see Vervaeke, I.J. 23(1989)169
- Dinsdale, D., see Cheng, K.-J. 10(1983/1984)93
- Dinter, B., see Weinberg, Z.G. 20(1988)335
- Dixon, R.M.
Effects of dietary concentrates on rumen digestion of fibrous feedstuffs 14(1986)193
- Djajanegara, A., Molina, B.T. and Doyle, P.T.
The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12(1984/1985)141
- Djajanegara, A. and Doyle, P.T.
Digestion rates in and outflow rates from the rumen of sheep fed untreated or calcium hydroxide-treated wheat straw 25(1989)179
- D'Mello, J.P.F.
Amino acid supplementation of methanol-grown dried microbial cells in diets for young chicks 4(1979)63
- D'Mello, J.P.F.
Limiting amino acids in dried microbial cells given to young turkeys 4(1979)209
- D'Mello, J.P.F. and Acamovic, T.
Growth performance of, and mimosine excretion by, young chicks fed on *Leucaena leucocephala* 7(1982)247
- D'Mello, J.P.F.
Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen 17(1987)1
- Donkoh, A., see Atuahene, C.C. 17(1987)157
- Donkoh, A., Atuahene, C.C., Kesse, A.G. and Mensah-Asante, B.
The nutritional value of dried coffee pulp (DCP) in broiler chickens's diets 22(1988)139
- Doran, B.E., see Goetsch, A.L. 18(1987)151
- Dove, H., see Freer, M. 11(1984)87
- Doyle, P.T., see Djajanegara, A. 12(1984/1985)141
- Doyle, P.T., see Djajanegara, A. 25(1989)179
- Draper, S.R., see El-Sherbiny, A.E. 5(1980)33

Draper, S.R., see El-Sherbiny, A.E. 5(1980)43
 Dryden, G.McL. and Kempton, T.J.

Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65

Dryden, G.McL. and Leng, R.A.

Treatment of barley straw with ammonia and sulphur dioxide gases under laboratory conditions 14(1986)41

Dryden, G.McL. and Leng, R.A.

Effects of ammonia and sulphur dioxide gases on the composition and digestion of barley straw 19(1988)121

Dumay, C., see Durand, M. 21(1988)197

Durand, M., see Szylit, O. 3(1978)73

Durand, M., Dumay, C., Beaumatin, P. and Morel, M.T.

Use of the rumen simulation technique (RUSITEC) to compare microbial digestion of various by-products 21(1988)197

Dutton, M.F., see Westlake, K. 25(1989)169

E

Eddebbarh, A., see Garrett, J.E. 25(1989)11

Edney, M.J., Campbell, G.L. and Classen, H.L.

The effect of β -glucanase supplementation on nutrient digestibility and growth in broilers given diets containing barley, oat groats or wheat 25(1989)193

Edwards, R.A., see Heron, S.J.E. 19(1988)85

Edwards, S.A., Fairbairn, C.B. and Capper, A.L.

Liquid potato feed for finishing pigs: feeding value, inclusion rate and storage properties 15(1986)129

Eckhout, W.

The energy systems used in the EEC countries: their application to 24 commercial pig-feeds and relationships between them 6(1981)211

Eckhout, W. and Moermans, R.

Prediction of the energy value of commercial pig feeds, according to the various energy systems used in the European Community, that are based on simple analytical parameters of the feed 6(1981)367

Eckhout, W. and de Paepe, M.

Alkali-treated soya flour for piglets 24(1989)1

Ehle, F.R., see Hannah, S.M. 16(1986/1987)51

Ekpenyong, T.E.

Nutrient and amino acid composition of *Leucaena leucocephala* (Lam.) de Wit 15(1986)183

Ekström, N., see Kaspersson, A. 19(1988)299

El-Alaily, H., see Mohamed, K. 11(1984)199

El Boushy, A.R., see Papadopoulos, M.C. 14(1986)279

El-Din, T., see Abel, H.J. 19(1988)111

El Hag, H.M.A. and Greenhalgh, J.F.D.

Intake and digestibility in sheep fed on alkali-treated straw and concentrates, as complete diets or separate feeds 7(1982)271

El Hag, M.G., Kurdi, O.I. and Maghoub, S.O.

Performance and carcass characteristics of Sudan desert sheep and goats on high roughage diets with added fat 13(1985)147

El Khidir, O.A. and Vestergaard Thomsen, K.

The effect of high levels of molasses in combinations with hay on digestibility of organic matter, microbial protein synthesis and volatile fatty acid production in vitro 7(1982)277

El Khidir, O.A. and Vestergaard Thomsen, K.

Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197

El Khidir, O.A., Gumaa, A.Y., Fangali, O.A.I. and Badir, N.A.

The use of Balanites kernel cake in a diet for fattening sheep 9(1983)301

El-Negoumy, A.M., see Moss, B.R. 8(1983)25

El Shazly, K., see Abel, H.J. 19(1988)111

El-Sherbiny, A.E., Draper, S.R. and Topps, J.H.

The determination of available lysine in barley using *Tetrahymena pyriformis* W. 5(1980)33

El-Sherbiny, A.E., Draper, S.R. and Topps, J.H.

A comparison of three methods for the determination of available lysine in barley 5(1980)43

El Tayeb, A.E., see Osman, A.G. 19(1988)351

El-Tayeb, A.E., Abdel-Raheim, F.M. and Awadel-Kareim, A.M.

Comparison of cottonseed cake and dried blood as nitrogen sources for growing Sudan Nubian goats 22(1988)333

Elliot, J.I., see Sibbald, I.R. 8(1983)191

Elliot, J.I., Modler, H.W. and Timbers, G.E.

A continuous process for the production of purified porcine γ -globulin for use in pig-milk replacer supplements 17(1987)213

Elliott, R., see McMeniman, N.P. 19(1988)43

Elliott, R., see Alcántara, E. 23(1989)323

Elliott, R., see Tangtaweewipat, S. 25(1989)123

Ellis, W.C., see Schake, L.M. 7(1982)257

Elwinger, K., see Hesselman, K. 7(1982)351

Ely, D.G., see Pendulum, L.C. 2(1977)225

Englyst, H.

Classification and measurement of plant polysaccharides 23(1989)27

Est, Y., see Ben-Ghedia, D. 19(1988)219

Est, Y., see Ben-Ghedia, D. 24(1989)289

Estergaard Thomsen, K., see Israelsen, M. 3(1978)227

Evans, D.F. and Leibholz, J.

Meat meal in the diet of the early-weaned pig. I. A comparison of meat meal and soya bean meal 4(1979)33

Evans, D.F. and Leibholz, J.

Meat meal in the diet of the early-weaned

- pig. II. Amino acid supplementation 4 (1979)43
- Everington, J.M., see Givens, D.I. 24(1989)27
- F**
- Faci, R., see Alibés, X. 10(1983/1984)239
- Fadel, J.G., see Robinson, P.H. 15(1986)249
- Fahey Jr., G.C., see Guardiola, C.M. 8(1983)129
- Fahey Jr., G.C., see Mora, M.I. 9(1983)205
- Fahey Jr., G.C., see Lynch, G.L. 14(1986)127
- Fahey Jr., G.C., see Lewis, S.M. 17(1987)179
- Fairbairn, C.B., see Edwards, S.A. 15(1986)129
- Fairbairn, R., see Alli, I. 9(1983)291
- Fallon, R.J., Williams, P.E.V. and Innes, G.M. The effects on feed intake, growth and digestibility of nutrients of including calcium soaps of fat in diets for young calves 14 (1986)103
- Fangali, O.A.I., see El Khidir, O.A. 9(1983)301
- Fanta, G.F., see Lewis, S.M. 17(1987)179
- Farrell, D.J. A nutritional evaluation of buckwheat (*Fagopyrum esculentum*) 3(1978)95
- Farrell, D.J., Chan, C., McCrae, F. and McKenzie, R.J. A nutritional evaluation of triticale with pigs 9(1983)49
- Farrell, D.J., Thomson, E., Choice, A., Ashes, J.R., Peck, N.J. and Hogan, J.P. Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983)99
- Farrell, D.J., see Raharjo, Y. 12(1984/1985)29
- Faulkner, D.B., Llamas, G.L., Ward, J.K. and Klopstein, T.J. Improving the intake and nutritive value of wheat straw for beef cows 12(1984/1985)125
- Fay, J.P. and Ovejero, F.M.A. Effect of lactate on the in vitro digestion of *Agropyron elongatum* by rumen microorganisms 16(1986/1987)161
- Fegeros, K., see Papadopoulos, G. 18(1987)303
- Fenwick, G.R. and Curtis, R.F. Rapeseed meal and its use in poultry diets. A review 5(1980)255
- Fenwick, G.R., Pearson, A.W., Greenwood, N.M. and Butler, E.J. Rapeseed meal tannins and egg taint 6 (1981)421
- Fenwick, G.R., see Pearson, A.W. 8(1983)119
- Fernandez Carmona, J., see Deltoro López, J. 6(1981)169
- Fernandez Carmona, J., see Deltoro López, J. 6(1981)179
- Fernandez-Carmona, J., see Cervera, C. 12 (1984/1985)233
- Fernández Carmona, J., see Martínez Pascual, J. 5(1980)1
- Fernández Carmona, J., see Martínez Pascual, J. 5(1980)11
- Fernández Carmona, J., see Martínez Pascual, J. 5(1980)23
- Fetuga, B.L., see Babatunde, G.M. 4(1979)23
- Fetuga, B.L., see Balogun, O.O. 7(1982)233
- Fetuga, B.L.A., see Balogun, O.O. 8(1983)197
- Fetuga, B.L.A., see Ikurior, S.A. 20(1988)251
- Fiallo, J.F., see Wohlt, J.E. 6(1981)115
- Fiems, L.O., see Bouqué, Ch.V. 7(1982)401
- Fiems, L.O., Bouqué, Ch.V., Cottyn, B.G. and Buysse, F.X. Effect of avoparcin on performance, body composition and meat quality of finishing bulls 8(1983)237
- Fiems, L.O., Cottyn, B.G., Bouqué, Ch.V. and Buysse, F.X. Effect of formaldehyde-treated soya bean meal and urea in starters on nitrogen quality, degradability in sacco, sheep digestibility and calf performance 16(1986/1987)287
- Findlay, M., see Partridge, G.G. 13(1985)299
- Findlay, M., see Partridge, G.G. 16(1986/1987)109
- Finley, J.W., see Nelson, M.L. 24(1989)141
- Fischer, J.R., see Goedecken, F.K. 13(1985)47
- Fishwick, G., see Bass, J.M. 7(1982)93
- Fishwick, G., see Alawa, J.P. 19(1988)33
- Flachowsky, G., Wolfgram, D., Wolf, I. and Hennig, A. Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 17(1987)235
- Flám, F., see Vencel, B. 6(1981)235
- Flipot, P., Mason, W. and Lalande, G. Chemical composition and animal performance of grass forage of varying maturity stored as hay or silage 11(1984)35
- Foale, M.A., see Kondos, A.C. 8(1983)85
- Folman, Y., see Holzer, Z. 5(1980)299
- Fonnesbeck, P.V., Garcia de Hernandez, M.M., Kaykay, J.M. and Saiady, M.Y. Estimating yield and nutrient losses due to rainfall on field-drying alfalfa hay 16(1986/1987)7
- Fonolla, J., Prieto, C. and Sanz, R. Influence of age on the nutrient utilization of diets for broilers 6(1981)405
- Fonolla, J., see Aguilera, J.F. 15(1986)33
- Fontenot, J.P., see Rao, N.M. 18(1987)67
- Fontenot, J.P., see Ortigues, I. 24(1989)13
- Fonty, G., see Stewart, C.S. 21(1988)69
- Fonty, G. and Gouet, Ph. Fibre-degrading microorganisms in the mono-gastric digestive tract 23(1989)91
- Fordey, R.A., see Partridge, G.G. 16(1986/1987)109
- Francesch, M., see Lacassagne, L. 20(1988)59
- Franci, O., see Antongiovanni, M. 5(1980)169
- Fraser, J., see Hemingway, R.G. 15(1986)123
- Frederiksen, J.H., see Kataglie, J.A. 4(1979)1
- Freer, M. and Dove, H.

- Rumen degradation of protein in sunflower meal, rapeseed meal and lupin seed placed in nylon bags 11(1984)87
- Froněk, P., see Heger, J. 8(1983)163
- Frydrych, Z. and Heger, J.
Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6 (1981)309
- Frydrych, Z., see Heger, J. 8(1983)163
- Frydrych, Z., see Heger, J. 15(1986)21
- Fundora, O., see Valdivie, M. 7(1982)185
- Funk, M.A., see Goetsch, A.L. 18(1987)151
- G**
- Gabir, S., see Maglad, M.A. 15(1986)69
- Galvez, J.F., see Caja, G. 2(1977)267
- Garcia, G., see Alli, I. 7(1982)411
- Garcia, G., see Alli, I. 9(1983)291
- Garcia de Hernandez, M.M., see Fomesbeck, P.V. 16(1986/1987)7
- Garmo, T.H., see Borhami, B.E.A. 7(1982)53
- Garmo, T.H., see Wanapat, M. 12(1984/1985) 295
- Garrett, J.E., Guessous, F. and Eddebarbar, A.
Utilization of sugar beet molasses and monensin for finishing dairy bullocks 25(1989) 11
- Garrigus, U.S., see Guardiola, C.M. 8(1983) 129
- Gasa, J., Castrillo, C., Baucells, M.D. and Guada, J.A.
By-products from the canning industry as feedstuff for ruminants: digestibility and its prediction from chemical composition and laboratory bioassays 25(1989)67
- Gascoyne, D.J. and Theodorou, M.K.
Consecutive batch culture - a novel technique for the in vitro study of mixed microbial populations from the rumen 21(1988) 183
- Gaspari, F., see Giardini, A. 1(1976)313
- Gaspari, F., see Giardini, A. 1(1976)465
- Gast, C.H., see Mascarenhas Ferreira, A. 9 (1983)19
- Gatel, F., Grosjean, F. and Castaing, J.
Feeding value of ensiled high-moisture maize grain with cob for growing-finishing pigs 20 (1988)145
- Gatel, F., Grosjean, F. and Leuillet, M.
Utilization of white-flowered smooth-seeded spring peas (*Pisum sativum hortense*, CV Amino) by the breeding sow 22(1988)91
- Gauld, S.A., see Mould, F.L. 10(1983/1984)31
- Gauld, S.A., see Williams, P.E.V. 10(1983/ 1984)247
- Geay, Y. and Malterre, C.
Nutritive value of dehydrated whole-crop maize 1(1976)429
- Genizi, A., see Goldman, A. 18(1987)233
- Giardini, A., Gaspari, F., Vecchietini, M. and Schenoni, P.
Effect of maize silage harvest stage on yield, plant composition and fermentation losses 1 (1976)313
- Giardini, A., Vecchietini, M. and Lo Bruno, A.
Energy supplementation of maize silage harvested at different maturity stages 1(1976) 369
- Giardini, A., Lambertini, F., Gaspari, F. and Lo Bruno, A.
Levels, sources and methods of nitrogen supplementation of maize silage for beef production 1(1976)465
- Gibson, D.M., Kennelly, J.J., Aherne, F.X. and Mathison, G.W.
Efficacy of sulfur dioxide as a grain preservative 19(1988)203
- Giger, S., see Sauvant, D. 13(1985)7
- Gil, F., see Aguilera, J.F. 15(1986)33
- Giorgetti, A., see Antongiovanni, M. 5(1980) 169
- Giouzeljannis, A., see Karalazos, A. 20(1988) 13
- Girardeau, J.P., see Gouet, P. 2(1977)297
- Givens, D.I. and Barber, W.P.
Nutritive value of apple pomace for ruminants 16(1986/1987)311
- Givens, D.I. and Brunnen, J.M.
Nutritive value of naked oats for ruminants 18(1987)83
- Givens, D.I., see Clark, P. 18(1987)225
- Givens, D.I., Adamson, A.H. and Cobby, J.M.
The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173
- Givens, D.I., Everington, J.M. and Adamson, A.H.
The digestibility and metabolisable energy content of grass silage and their prediction from laboratory measurements 24(1989)27
- Godeau, J.-M., see Teller, E. 13(1985)25
- Goedeken, F.K., Paterson, J.A., Koeln, L.L., Fischer, J.R. and Williams, J.E.
Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47
- Goetsch, A.L., Owens, F.N., Funk, M.A. and Doran, B.E.
Effects of whole or ground corn with different forms of hay in 85% concentrate diets on digestion and passage rate in beef heifers 18 (1987)151
- Goetsch, A.L., see Hardin, A.C. 25(1989)99
- Goldman, A., Genizi, A., Yulzari, A. and Seligman, N.G.
Improving the reliability of the two-stage in vitro assay for ruminant feed digestibility by calibration against in vivo data from a wide range of sources 18(1987)233
- Gómez, G., see Santos, J. 8(1983)313

- Gómez, G. and Valdivieso, M.
Cassava for animal feeding: effect of variety and plant age on production of leaves and roots 11(1984)49
- Gómez, G., Valdivieso, M., de la Cuesta, D. and Salcedo, T.S.
Effect of variety and plant age on the cyanide content of whole-root cassava chips and its reduction by sun-drying 11(1984)57
- Gomez, M., see Abate, A.N. 10(1983/1984)291
- Gonzalez, G. and Monteagudo, A.
Survey of the use of maize for livestock feeding in Spain 1(1976)129
- Gonzalez, G., see Alzueta, C. 8(1983)155
- González, G., Alzueta, C., Barro, C. and Salvador, A.
Yield and composition of protein concentrate, press cake, green juice and solubles concentrate from wet fractionation of *Sophora japonica* L. foliage 20(1988)177
- González, G., see Rebolé, A. 22(1988)173
- Gonzalez Santillana, R., see Wilkinson, J.M. 3(1978)117
- Gonzalez Santillana, R., see Wilkinson, J.M. 3(1978)133
- Gonzalez Santillana, R. and Wilkinson, J.M.
Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381
- Goode, L., see Johnson, T.O. 12(1984/1985) 65
- Göransson, L., Olsson, O. and Larsson, K.
Effects of different handling systems on pig feed quality 7(1982)11
- Göransson, L. and Ogle, R.B.
Anaerobically stored high moisture cereals for growing pigs 12(1984/1985)159
- Gorbet, D.W., see Myer, R.O. 12(1984/1985) 179
- Gordon, A.H. and Chesson, A.
The effect of prolonged storage on the digestibility and nitrogen content of ammonia-treated barley straw 8(1983)147
- Gordon, G.L.R., see Akin, D.E. 23(1989)305
- Gorman, T.B.S., see Batterham, E.S. 9(1983) 169
- Goto, I. and Minson, D.J.
Prediction of the dry matter digestibility of tropical grasses using a pepsin-cellulase assay 2(1977)247
- Goto, I., see Shimojo, M. 24(1989)173
- Gouet, P., Girardeau, J.P. and Riou, Y.
Inhibition of *Listeria monocytogenes* by defined lactic microflora in gnotobiotic silages of lucerne, fescue, ryegrass and maize - influence of dry matter and temperature 2 (1977)297
- Gouet, Ph., see Stewart, C.S. 21(1988)69
- Gouet, Ph., see Fonty, G. 23(1989)91
- Gould, J.M., see Lewis, S.M. 17(1987)179
- Goulden, K.M., see Whitehead, D.C. 14(1986) 231
- Graham, C.A., see Kellaway, R.C. 3(1978)201
- Graham, H. and Åman, P.
A comparison between degradation in vitro and in sacco of constituents of untreated and ammonia-treated barley straw 10(1983/ 1984)199
- Graham, H., Åman, P., Theander, O., Kolan- kaya, N. and Stewart, C.S.
Influence of heat sterilization and ammonia- tion on straw composition and degradation by pure cultures of cellulolytic rumen bac- teria 12(1984/1985)195
- Graham, H., see Åman, P. 17(1987)15
- Graham, H. and Åman, P.
Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal tract of pigs 17(1987)33
- Graham, H., see Nordkvist, E. 17(1987)295
- Graham, H., see Theander, O. 23(1989)205
- Graham, H., see Narasimhalu, P. 24(1989)159
- Grain, J., see Jouany, J.P. 21(1988)229
- Green, A.G. and Oram, R.N.
Variability for protein and oil quality in *Lupinus albus* 9(1983)271
- Green, S., Wiseman, J. and Cole, D.J.A.
Examination of stability, and its effect on the nutritive value, of fish silage in diets for growing pigs 21(1988)43
- Greenhalgh, J.F.D.
The dilemma of animal feeds and nutrition 1(1976)1
- Greenhalgh, J.F.D., see Manten, A.A. 2(1977) 1
- Greenhalgh, J.F.D., see Pirie, R. 3(1978)143
- Greenhalgh, J.F.D., Pirie, R., Shin, H.T. and Stewart, C.S.
Alkali treatment of straw for ruminants. II. Nutritive value of straw ensiled after alkali treatment. 3(1978)289
- Greenhalgh, J.F.D., see El Hag, H.M.A. 7 (1982)271
- Greenhalgh, J.F.D., see Charmley, E. 18(1987) 19
- Greenwood, N.M., see Fenwick, G.R. 6(1981) 421
- Greenwood, N.M., see Pearson, A.W. 8(1983) 119
- Gregory, K.F., Reade, A.E., Santos-Nunez, J., Alexander, J.C., Smith, R.E. and MacLean, S.J.
Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977) 7
- Greife, H.A., Molnar, S. and Günther, K.-D.
Nitrogen metabolism in broiler chickens consuming the bacterial strain *Alcaligenes eutrophus* 5(1980)241
- Grenet, E. and Barry, P.
Colonization of thick-walled plant tissues by anaerobic fungi 19(1988)25
- Griffith, E.S., see McAllan, A.B. 17(1987)65
- Grigera-Naón, J.J., see Smith, T. 10(1983/ 1984)189
- Grigera-Naón, J.J., see Smith, T. 14(1986)29

- Grimbergen, A.H.M., Cornelissen, J.P. and Stappers, H.P.
The relative availability of phosphorus in inorganic feed phosphates for young turkeys and pigs 13(1985)117
- Grimes, L.W., see Grotheer, M.D. 14(1986)55
- Grosjean, F., see Gatel, F. 20(1988)145
- Grosjean, F., see Gatel, F. 22(1988)91
- Grotheer, M.D., Cross, D.L. and Grimes, L.W.
Effect of ammonia level and time of exposure to ammonia on nutritional and preservative characteristics of dry and high-moisture coastal bermuda grass hay 14(1986)55
- Grubb, D.A., see Ørskov, E.R. 2(1977)307
- Grubb, D.A., see Ørskov, E.R. 6(1981)273
- Grummer, R.H., see Cieslak, D.G. 14(1986)171
- Grummitt, N., see Minson, D.J. 9(1983)221
- Guada, J.A., see Gasa, J. 25(1989)67
- Guardiola, C.M., Fahey Jr., G.C., Spears, J.W., Garrigus, U.S., Izquierdo, O.A. and Pedroza, C.
The effects of sulphur supplementation on cellulose digestion in vitro and on nutrient digestion, nitrogen metabolism and rumen characteristics of lambs fed on good quality fescue and tropical star grass hays 8(1983)129
- Guedes, C.V.M., see Dias-da-Silva, A.A. 19(1986)67
- Guedes, C.V.M., see Mascarenhas-Ferreira, A. 25(1989)157
- Gunter, W., see Al-Athari, A.K. 22(1988)119
- Gunter, W., see Al-Athari, A.K. 22(1988)273
- Gunter, W., see Rotter, B.A. 24(1989)233
- Guessous, F., see Garrett, J.E. 25(1989)11
- Gumaa, A.Y., see El Khidir, O.A. 9(1983)301
- Günther, K.-D., see Greife, H.A. 5(1980)241
- ## H
- Ha, J.K., Kennelly, J.J. and Berzins, R.
Effect of dietary nitrogen source on microbial protein synthesis, dietary protein degradation and nutrient digestion in steers 14(1986)117
- Hadjipanayiotou, M.
The value of urea-treated straw in diets of lactating goats 11(1984)67
- Hadobás, K., see Hegedüs, M. 9(1983)79
- Hahn-Hägerdal, B., Tjerneld, F. and Zacchi, G.
Production of ethanol from lignocellulosic materials 21(1988)175
- Håkansson, J., see Lund, D. 16(1986/1987)119
- Hall, J.M.R., see Wanapat, M. 14(1986)215
- Hall, K.L., see Hardin, A.C. 25(1989)99
- Hamilton, R.M.G. and Trenholm, H.L.
Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin) 11(1984)293
- Hanczakowska, E., see Hanczakowski, P. 6(1981)189
- Hanczakowski, P. and Petzel, M.
Nutritive value of leaf protein coagulum and cereals mixtures 4(1979)91
- Hanczakowski, P., Skraba, B. and Hanczakowska, E.
Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189
- Hanczakowski, P., Skraba, B. and Młodkowski, M.
Nutritive value of leaf-protein concentrate from potato haulm for rats and chicks 6(1981)413
- Hanczakowski, P., see Carlsson, R. 11(1984)239
- Hanczakowski, P. and Skraba, B.
The effect of different precipitating agents on quality of leaf protein concentrate from lucerne 12(1984/1985)11
- Hanczakowski, P. and Skraba, B.
The effect of toxic substances present in some leaf protein concentrates on their nutritive value 24(1989)151
- Hannah, S.M., Stern, M.D. and Ehle, F.R.
Evaluation of a dual flow continuous culture system for estimating bacterial fermentation in vivo of mixed diets containing various soya bean products 16(1986/1987)51
- Hanrahan, T.J.
Effect of pellet size and pellet quality on pig performance 10(1983/1984)277
- Hansen, P., see Opstvedt, J. 18(1987)181
- Hansen, J.F., see Muindi, P.J. 6(1981)197
- Hardin, A.C., Goetsch, A.L., Landis, K.M., Murphy, G.E., Johnson, Z.B. and Hall, K.L.
Intake, digestion and daily gain by cattle consuming bermudagrass (*Cynodon dactylon*) and supplemented with different combinations of ground corn, vegetable oil, urea and corn gluten and blood meals 25(1989)99
- Hargreave, G.T., see Leibholz, J. 5(1980)309
- Harmon, R.J., see Teh, T.H. 17(1987)257
- Harper, A.F., Kornegay, E.T., Bryant, K.L. and Thomas, H.R.
Efficacy of virginiamycin and a commercially-available lactobacillus probiotic in swine diets 8(1983)69
- Harte, F.J.
The current situation regarding the use of maize in Ireland 1(1976)195
- Hartley, R.D., see Theander, O. 10(1983/1984)89
- Hartley, R.D., see Van Soest, P.J. 10(1983/1984)155
- Hartley, R.D., see Whitehead, D.C. 14(1986)231
- Hartley, R.D., see Mason, V.C. 19(1988)159
- Hartley, R.D., see Akin, D.E. 19(1988)261
- Hartley, R.D., see Mason, V.C. 24(1989)299
- Harvey, R.W., see Johnson, T.O. 12(1984/1985)65

- Harvey, R.W., see Ortigues, I. 24(1989)13
- Harvey, R.W. and Spears, J.W.
Effect of blood meal processing method on performance and nitrogen utilization by growing cattle 24(1989)337
- Hasimoglu, S., see Umunna, N.N. 7(1982)375
- Hazlewood, G.P., see Orpin, C.G. 21(1988)161
- Heath, M.E., see Waterworth, D.G. 6(1981)297
- Hector, D.A., see Machin, D.H. 11(1984)247
- Hector, D.A., see Machin, D.H. 15(1986)273
- Hegedüs, M., Bokori, J. and Hadobás, K.
Protein quality of mixtures of animal protein meals 9(1983)79
- Heger, J., see Frydrych, Z. 6(1981)309
- Heger, J., Frydrych, Z. and Froněk, P.
Evaluation of optimum lysine and threonine supplements to a wheat and barley-based diet in rats 8(1983)163
- Heger, J. and Frydrych, Z.
Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21
- Heikonen, M., see Moisis, T. 22(1988)341
- Hembry, F.G., see Coelho, M. 20(1988)219
- Hemingway, R.G., Parkins, J.J. and Fraser, J.
Sugar beet pulp products for dairy cows 15(1986)123
- Hemingway, R.G., see Alawa, J.P. 19(1988)33
- Hemken, R.W., see Teh, T.H. 17(1987)257
- Henderson, A.R., McDonald, P. and Anderson, D.H.
The effect of silage additives containing formaldehyde on the fermentation of ryegrass ensiled at different dry matter levels and on the nutritive value of direct-cut silage 7(1982)303
- Henderson, A.R., see McDonald, P. 8(1983)259
- Henderson, C. and Demeyer, D.I.
The rumen as a model of the microbiology of fibre digestion in the human intestine 23(1989)227
- Hendrosoekarjo, S. and Pearce, G.R.
Utilization by sheep of dried pig faeces containing a high concentration of copper 3(1978)31
- Hennig, A., see Flachowsky, G. 17(1987)235
- Hennig, A., see Schöne, F. 22(1988)45
- Henry, P.R., see Potter, S.G. 12(1984/1985)275
- Hermansen, J.E.
Feed intake and milk yield at increasing supplement of a palmitic and stearic acid-rich type of fat in comparison with animal fat 22(1988)179
- Hermansen, J.E.
Feed intake, milk yield and milk composition by replacing unprotected fat by Ca-soaps for dairy cows 22(1988)193
- Heron, S.J.E., Edwards, R.A. and McDonald, P.
The effects of inoculation, addition of glucose and minging on fermentation and proteolysis in ryegrass ensiled in laboratory silos 19(1988)85
- Hesselman, K., Elwinger, K. and Thomke, S.
Influence of increasing levels of β -glucanase on the productive value of barley diets for broiler chickens 7(1982)351
- Hesselman, K. and Åman, P.
The effect of β -glucanase on the utilization of starch and nitrogen by broiler chickens fed on barley of low- or high-viscosity 15(1986)83
- Hesselman, K., see Pettersson, D. 17(1987)145
- Hibberd, C.A., Wagner, D.G., Schemm, R.L., Mitchell Jr., E.D., Weibel, D.E. and Hintz, R.L.
Response of different sorghum grain and maize varieties to reconstitution 15(1986)231
- Hibberd, C.A., see Chase Jr., C.C. 24(1989)129
- Higgins, T.R. and Spooner, A.E.
Microwave drying of alfalfa compared to field- and oven-drying: effects on forage quality 16(1986/1987)1
- Hinds, M.A., Bolsen, K.K., Brethour, J., Miliken, G. and Hoover, J.
Effects of molasses/urea and bacterial inoculant additives on silage quality, dry matter recovery, and feeding value for cattle 12(1984/1985)205
- Hinks, C.E.
The replacement of skim milk by dried microbial cells and whey in milk diets for calves 2(1977)85
- Hintz, R.L., see Hibberd, C.A. 15(1986)231
- Hlödversson, R. and Kaspersson, A.
Nutrient losses during deterioration of hay in relation to changes in biochemical composition and microbial growth 15(1986)149
- Hlödversson, R.
The nutritive value of white- and dark-flowered cultivars of pea for growing-finishing pigs 17(1987)245
- Hlödversson, R.
The effect of mechanical treatment on the nutritive value of high-moisture barley and the performance of growing-finishing pigs 24(1989)121
- Ho, Y.W., Abdullah, N. and Jalaludin, S.
Colonization of guinea grass by anaerobic rumen fungi in swamp buffalo and cattle 22(1988)161
- Hoadley, C.J., see Mason, V.C. 24(1989)313
- Hoebler, C., see Champ, M. 23(1989)195
- Hof, G., see Deinum, B. 10(1983/1984)301
- Hogan, J.P., see Farrell, D.J. 9(1983)99
- Holland, S.M., see Ørskov, E.R. 8(1983)247
- Holzer, Z., Levy, D. and Folman, Y.
Performance of fattening cattle and lactating beef cows on diets high in chemically treated wheat straw and poultry litter 5(1980)299

- Holzer, Z., Levy, D., Samuel, V., Angemar, Y. and Ilan, D.
Wheat silage for fattening young male cattle 12(1984/1985)253
- Holzer, Z., see Levy, D. 16(1986/1987)63
- Holzgraebe, D.P., see Lewis, S.M. 17(1987)179
- Hoover, J., see Hinds, M.A. 12(1984/1985)205
- Horn, G.W., see Streeter, C.L. 7(1982)325
- Horn, G.W., Batchelder, D.G., Manor, G., Streeter, C.L. and McLaughlin, G.L.
Ammoniation of wheat straw and native grass hay during baling of large round bales 8(1983)35
- Horn, G.W., see Zorrilla-Rios, J. 22(1988)305
- Horn, G.W., Zorrilla-Rios, J. and Akin, D.E.
Influence of stage of forage maturity and ammoniation of wheat straw on ruminal degradation of wheat forage tissues 24(1989)201
- Horner, J.I., see Coppock, C.E. 18(1987)89
- Horton, G.M.J., Nicholson, H.H. and Christensen, D.A.
Ammonia and sodium hydroxide treatment of wheat straw in diets for fattening steers 7(1982)1
- Horvath, D.J., see Reid, R.L. 5(1980)95
- Horvath, P.J., see Jeraci, J.L. 23(1989)121
- Hosking, Z.D., see Barber, R.S. 4(1979)117
- Hove, E.L., see Ruiz Jr., L.P. 2(1977)59
- Huhtanen, P.
The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988)259
- Huhtanen, P. and Robertson, S.
The effect of dietary inclusion of starch, sucrose and xylose on the utilization of dietary energy in sheep 21(1988)11
- Huisman, J., see Van Weerden, E.J. 2(1977)377
- Hulan, H.W., see Zarkadas, C.G. 14(1986)291
- Hulan, H.W., Proudfoot, F.G. and Zarkadas, C.G.
The effect of adding white fish meal containing enzyme digested or untreated stickwater solids to diets for broiler chickens 16(1986/1987)253
- Hunt, C.W., see Koeln, L.L. 11(1984)181
- Hunt, J.R. and Barton, G.M.
Nutritive value of spruce muka (foliage) for the growing chick 3(1978)63
- Husain, R.A.K. and Offer, N.W.
Effect of formaldehyde treatment on the degradation of acid-preserved fish silage protein in vitro 16(1986/1987)297
- Husain, R.A.K., see Offer, N.W. 17(1987)165
- Hussar, N., see Leeson, S. 19(1988)313
- Hutagalung, R.I., see Abdullah, N. 20(1988)79
- Huyghebaert, G., De Munter, G. and De Groote, G.
The metabolisable energy (AME_n) of fats for broilers in relation to their chemical composition 20(1988)45
- Hyslop, J.J., Weir, J., Offer, N.W., Reid, S.J. and Wilcock, J.M.
Effect of formaldehyde treatment on the rumen digestion of barley and wheat grain and its influence on hay intake and digestion by sheep 25(1989)55
- ## I
- Ibbotson, C.F., Mansbridge, R. and Adamson, A.H.
Commercial experience of treating straw with ammonia 10(1983/1984)223
- Ibrahim, M.N.M., Tammings, S. and Zemelink, G.
Effect of urea treatment on rumen degradation characteristics of rice straws 24(1989)83
- Igwe, F.O., see Udedibie, A.B.I. 24(1989)111
- Ihnat, M., see Ivan, M. 9(1983)131
- Ikurior, S.A. and Fetuga, B.L.A.
Chemical and biological evaluation of Nigerian cottonseed meal protein quality 20(1988)251
- Ilan, D., see Holzer, Z. 12(1984/1985)253
- Ilan, D., see Levy, D. 16(1986/1987)63
- Ilian, M.A., Razzaque, M.A., Al-Awadi, A. and Salman, A.J.
Use of fat in diets of sheep in hot environments. I. Effects on performance, carcass characteristics and lipid composition of plasma 19(1988)327
- Ilian, M.A., Razzaque, M.A. and Salman, A.J.
Use of fat in diets of sheep in hot environments. II. Effect on rumen metabolism 19(1988)343
- Ilori, J.O., see Smith, O.B. 11(1984)231
- Ingalls, J.R., see Sharma, H.R. 10(1983/1984)77
- Ingalls, J.R., see Sharma, H.R. 15(1986)101
- Ingalls, J.R., see Mbugi, P.K. 24(1989)267
- Innes, G.M., see Ørskov, E.R. 6(1981)273
- Innes, G.M., see Ørskov, E.R. 6(1981)355
- Innes, G.M., see Williams, P.E.V. 10(1983/1984)247
- Innes, G.M., see Williams, P.E.V. 11(1984)103
- Innes, G.M., see Williams, P.E.V. 11(1984)115
- Innes, G.M., see Williams, P.E.V. 12(1984/1985)1
- Innes, G.M., see Fallon, R.J. 14(1986)103
- Israelson, M., Rexen, B. and Estergaard Thomsen, K.
Cellulose insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3(1978)227
- Istasse, L., Reid, G.W., Tait, C.A.G. and Ørskov, E.R.
Concentrates for dairy cows: effects of feeding method, proportion in diet and type 15(1986)167
- Istasse, L., Smart, R.I. and Ørskov, E.R.

- Comparison between two methods of feeding concentrate to sheep given a diet high or low in concentrate with or without buffering substances 16(1986/1987)37
- Istasse, L., Rollin, F., Kamani Djiele, N., Van Eenae, C., Biston, L. and Bienfait, J.M. Discriminating power of the nylon bag technique and the pepsin-cellulase method to detect less commonly used components included in concentrates for dairy cows 25 (1989)1
- Ivan, M., Ihnat, M. and Veira, D.M. Solubility and flow of calcium, magnesium and phosphorus in the digestive tract of sheep given maize or alfalfa silages 9(1983) 131
- Izquierdo, O.A., see Guardiola, C.M. 8(1983) 129
- ## J
- Jackson, A.J., see Anglesea, J.D. 13(1985)39
- Jackson, M.G.
Review article: the alkali treatment of straws 2(1977)105
- Jackson, M.G.
Who needs feeding standards? 6(1981)101
- Jahreis, G., see Schöne, F. 22(1988)45
- Jain, V.K., Jayal, M.M. and Pathak, N.N.
Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315
- Jakhmola, R.C., see Kamra, D.N. 9(1983)185
- Jakhmola, R.C., see Singh, R. 12(1984/1985) 133
- Jakhmola, R.C., Kundu, S.S., Punj, M.L., Singh, K., Kamra, D.N. and Singh, R. Animal excreta as ruminant feed - scope and limitations under Indian conditions 19 (1988)1
- Jalaludin, S., see Ho, Y.W. 22(1988)161
- James, K.A.C., see Smith, W.C. 18(1987)143
- Janssen, W.M.M.A., see Kan, C.A. 19(1988) 97
- Jayal, M.M., see Jain, V.K. 5(1980)315
- Jayasuriya, M.C.N. and Panditharatne, S. Spent tea leaf as a ruminant feed 3(1978) 219
- Jayasuriya, M.C.N., Perdok, H.B., Ross-Parker, H.M. and van Houtert, M.F.J. Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7 (1982)201
- Jayasuriya, M.C.N., Wijayatunge, C. and Perera, H.G.D. Rumen and post-rumen fermentation of spent tea leaf protein and other protein sources studied by the nylon bag method 7(1982)221
- Jayasuriya, M.C.N. and Pearce, G.R. The effect of urease enzyme on treatment time and the nutritive value of straw treated with ammonia as urea 8(1983)271
- Jelev, K., see Peteva-Vantcheva, Z. 3(1978)23
- Jensen, A.H., see Adams, K.L. 12(1984/1985) 267
- Jensen, A.H., see Adams, K.L. 13(1985)159
- Jensen, A.H., see Adams, K.L. 17(1987)201
- Jeraci, J.L. and Lewis, B.A.
Determination of soluble fiber components: (1-<3; 1-<4)- β -D-glucans and pectins 23 (1989)15
- Jeraci, J.L. and Horvath, P.J.
In vitro fermentation of dietary fiber by human fecal organisms 23(1989)121
- Jewell, S.N. and Campling, R.C.
Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81
- Job, T.A., see Aregheore, E.M. 20(1988)233
- Johnson, K.D., see Cherney, J.H. 17(1987)45
- Johnson, T.O., Harvey, R.W., Goode, L., Linnerud, A.C. and Crickenberger, R.G. Effects of stage of maturity and addition of molasses on nutritive value of maize stover silage 12(1984/1985)65
- Johnson, W.L., see Van Eys, J.E. 18(1987)197
- Johnson, Z.B., see Noland, P.R. 2(1977)219
- Johnson, Z.B., see Waldrup, P.W. 3(1978)329
- Johnson, Z.B., see Noland, P.R. 5(1980)51
- Johnson, Z.B., see Waldrup, P.W. 7(1982)419
- Johnson, Z.B., see Hardin, A.C. 25(1989)99
- Jones, A.S., see Livingstone, R.M. 2(1977)31
- Jones, A.S., see Livingstone, R.M. 2(1977)277
- Joshi, A.L., see Rangnekar, D.V. 7(1982)61
- Jouany, J.P. and Thivend, P.
In vitro effects of avoparcin on protein degradability and rumen fermentation 15 (1986)215
- Jouany, J.P., Demeyer, D.I. and Grain, J. Effect of defaunating the rumen 21(1988) 229
- Jovanović, M. and Čuperlović, M.
Nutritive value of rumen contents for monogastric animals 2(1977)351
- Jovanović, M., see Nikolić, J.A. 15(1986)57
- ## K
- Kabajia, E. and Smith, O.B.
The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171
- Kadirvel, R., see Thanu, K. 9(1983)263
- Kadirvel, R., see Poyyamozhi, V.S. 15(1986) 95
- Kadirvel, R., see Subramanian, P.R. 20(1988) 343
- Kadirvel, R., see Chandrasekaran, D. 22(1988) 321
- Kadirvel, R., see Viswanathan, K. 22(1988) 327

- Kadirvel, R., see Natanam, R. 25(1989)201
- Kalač, P. and Kyzlink, V.
Losses of beta-carotene in red clover in an acid medium during ensiling 4(1979)81
- Kalač, P. and Kyzlink, V.
The enzymic nature of the degradation of beta-carotene in red clover and in other forage crops during silage-making with acid additives 5(1980)59
- Kalač, P.
Losses of beta-carotene in unwilted forage crops during silage-making and feeding 9(1983)63
- Kali, J., see Levy, D. 16(1986/1987)63
- Kameni Djiele, N., see Istasse, L. 25(1989)1
- Kamra, D.N., Singh, R., Jakhmola, R.C. and Srivastava, R.V.N.
Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185
- Kamra, D.N., see Singh, R. 12(1984/1985)133
- Kamra, D.N., see Jakhmola, R.C. 19(1988)1
- Kamra, D.N., Lall, D., Kewalramant, N. and Pathak, N.N.
Ensiling characteristics of cattle waste fermented with wheat straw and green berseem (*Trifolium alexandrinum*) 25(1989)149
- Kan, C.A., Scheele, C.W. and Janssen, W.M.M.A.
The energy content of full-fat soya beans in meal and pelleted feeds for adult cocks and broilers 19(1988)97
- Kaptur, T., see Carlsson, R. 11(1984)239
- Karalazos, A. and Swan, H.
The nutritional value for sheep of molasses and condensed molasses solubles 2(1977)143
- Karalazos, A. and Giouzeljannis, A.
A note on the use of sugar-beet pulp silage and molasses in the diet of lactating dairy cows 20(1988)13
- Karamitros, D.
Sugar beet molasses for growing and fattening pigs 18(1987)131
- Karue, C.N., see Abate, A. 6(1981)15
- Karunajeewa, H. and Tham, S.H.
The influence of oat groats and dietary level of lysine on the laying performance of cross-bred hens 17(1987)271
- Kasperson, A., see Hlödversson, R. 15(1986)149
- Kasperson, A., Lindgren, S. and Ekström, N.
Microbial dynamics in barley grain stored under controlled atmosphere 19(1988)299
- Kategile, J.A. and Frederiksen, J.H.
Effect of level of sodium hydroxide treatment and volume of solution on the nutritive value of maize cobs 4(1979)1
- Kategile, J.A.
Performance of heifers fed on diets based on NaOH-treated maize cobs and the effect of supplementary urea and source of carbohydrates 4(1979)97
- Kategile, J.A., Urio, N.A., Sundstøl, F. and Mzihirwa, Y.G.
Simplified method for alkali treatment of low-quality roughages for use by small-holders in developing countries 6(1981)133
- Kategile, J.A., see Kiangi, E.M.I. 6(1981)377
- Kategile, J.A.
Feed value of Tanzanian sisal wastes: chemical composition and digestibility in vitro 14(1986)255
- Kategile, J.A., Mngulwi, J.G.J. and Abate, A.
Net energy for gain (NE_{gain}) of cassava tops 20(1988)97
- Kategile, J.A., see Sarwatt, S.V. 22(1988)237
- Katz, R.J., see Thomas, V.M. 11(1984)269
- Kaufman, W.
Discussion on session 4 1(1976)485
- Kay, M., see Mbatya, P.B.A. 8(1983)221
- Kay, M., see Mbatya, P.B.A. 8(1983)229
- Kay, M., see Mbatya, P.B.A. 13(1985)281
- Kay, M., see Mbatya, P.B.A. 13(1985)293
- Kaykay, J.M., see Fomesbeck, P.V. 16(1986/1987)7
- Kayongo-Male, H., see Abate, A. 6(1981)15
- Kayongo-Male, H., see Nangole, F.N. 9(1983)121
- Kayouli, C., Demeyer, D.I., Van Nevel, C.J. and Dendooven, R.
Effect of defaunation on straw digestion in sacco and on particle retention in the rumen 10(1983/1984)165
- Keene, A.S., see Mason, V.C. 19(1988)159
- Keene, A.S., see Mason, V.C. 24(1989)299
- Keith, M.O., see Bell, J.M. 24(1989)253
- Kellaway, R.C., Crofts, F.C., Thiago, L.R.L., Redman, R.G., Leibholz, J.M.L. and Graham, C.A.
A new technique for upgrading the nutritive value of roughages under field conditions 3(1978)201
- Kellaway, R.C., see Leibholz, J. 5(1980)309
- Kellaway, R.C., see Thiago, L.R.L. de S. 7(1982)71
- Kellaway, R.C., see Amaning-Kwarteng, K. 16(1986/1987)75
- Kelleher, D.L., see Caffrey, P.J. 19(1988)231
- Kelly, P.
Sugar beet pulp - a review 8(1983)1
- Kempton, T.J., see Abidin, Z. 6(1981)145
- Kempton, T.J., see Dryden, G.M.L. 10(1983/1984)65
- Kennedy, S., see Mbatya, P.B.A. 13(1985)281
- Kennedy, S., see Mbatya, P.B.A. 13(1985)293
- Kennelly, J.J., see Ha, J.K. 14(1986)117
- Kennelly, J.J., see Gibson, D.M. 19(1988)203
- Kernan, J.A., Coxworth, E.C. and Spurr, D.T.
New crop residues and forages for Western Canada: assessment of feeding value in vitro and response to ammonia treatment 6(1981)257
- Kernan, J.A., Crowe, W.L., Spurr, D.T. and Coxworth, E.C.
The nutritional value of crop residue components from several wheat cultivars grown at different fertilizer levels 11(1984)301
- Kernan, J.A., see Mann, M.E. 21(1988)57
- Kerstens, J., see Mascarenhas Ferreira, A. 9

- (1983)19
 Kese, A.G., see Donkoh, A. 22(1988)139
 Ketelaars, E.H., see Papadopoulos, M.C. 14 (1986)279
 Ketels, E. and de Groot, G.
 The nutritional value for broilers of fats characterized by short-chain fatty acids as affected by level of inclusion and age 22 (1988)105
 Kewalramant, N., see Kamra, D.N. 25(1989) 149
 Kharat, S.T., see Rangnekar, D.V. 7(1982)61
 Kiangi, E.M.I. and Kategile, J.A.
 Different sources of ammonia for improving the nutritive value of low quality roughages 6(1981)377
 Kies, A.K., Moughan, P.J. and Smith, W.C.
 The apparent and true ileal digestibility of nitrogen and amino acids in lactic casein for the growing pig 16(1986/1987)169
 Kilkenny, J.B.
 The use of maize for livestock feeding in the United Kingdom 1(1976)199
 King, R.H. and Campbell, R.G.
 Blood meal as a source of protein for grower/finisher pigs 3(1978)191
 King, R.H.
 Lupin-seed meal (*Lupinus albus* cv. Hamburg) as a source of protein for growing pigs 6(1981)285
 Kipnis, T., see Silanikove, N. 20(1988)87
 Kirby, A.C., see Amaning-Kwarteng, K. 16 (1986/1987)75
 Kirchgessner, M., see Roth, F.X. 1(1976)33
 Kirchgessner, M., see Stehr, W. 1(1976)53
 Kirchgessner, M., see Weigand, E. 5(1980)221
 Kirchgessner, M., Schwarz, F.J. and Lindner, H.P.
 Feed consumption and performance of dairy cows with alternate feedings of grass silage and maize silage 6(1981)337
 Kirchgessner, M., see Kreuzer, M. 16(1986/1987)233
 Kirton, A.H., see Wright, D.E. 2(1977)93
 Klasing, K.C., see Brown, D.L. 18(1987)247
 Klasing, K.C., see Shqueir, A.A. 25(1989)137
 Klopfenstein, T.J., see Umunna, N.N. 7(1982) 375
 Klopfenstein, T.J., see Faulkner, D.B. 12 (1984/1985)125
 Klopfenstein, T.J., see Umunna, N.N. 12 (1984/1985)151
 Klopfenstein, T.J., see Umunna, N.N. 14 (1986)161
 Koeln, L.L., Hunt, C.W., Bowman, D.K. and Paterson, J.A.
 Maize whole stillage as a supplemental protein source for growing lambs and finishing steers 11(1984)181
 Koeln, L.L., see Goedeke, F.K. 13(1985)47
 Koes, R.M., see Rule, D.C. 15(1986)113
 Kolankaya, N., see Graham, H. 12(1984/1985) 195
 Kondos, A.C. and Foale, M.A.
 Comparison of the nutritional value of low and medium tannin sorghum grains for pigs 8(1983)85
 Koong, L.J., see Sidahmed, A.E. 8(1983)47
 Kornegay, E.T.
 Protein digestibility of hydrolyzed hog hair meal for pigs 3(1978)323
 Kornegay, E.T., see Harper, A.F. 8(1983)69
 Kornegay, E.T., see Ravindran, V. 17(1987) 227
 Kothandaraman, P., see Narahari, D. 9(1983) 319
 Kothandaraman, P., see Narahari, D. 10 (1983/1984)257
 Koukoura, Z.
 Composition of kermes oak browse as affected by shade and stage of maturity 21 (1988)1
 Kowalczyk, J., Robinson, J.J. and Otwinowska, A.
 The digestion in the small intestine of young bulls of the protein of rapeseed meal treated or untreated with formaldehyde 7(1982)225
 Kreuzer, M., Kirschgessner, M. and Müller, H.L.
 Effect of defaunation on the loss of energy in wethers fed different quantities of cellulose and normal or steamflaked maize starch 16 (1986/1987)233
 Krishna, N., see Chandra, S. 12(1984/1985) 187
 Krishna Reddy, G.V. and Raj Reddy, M.
 Nutritive value of rice straw (*Oryza sativa*) ensiled with animal excreta and rumen digesta 24(1989)69
 Kromann, R.P., see Murdoch, F.R. 2(1977)287
 Kugenheim, Y., see Silanikove, N. 20(1988)87
 Kumaresan, A., Mshelia, T.A. and Aliu, Y.O.
 Biochemical evaluation of bagaruwa seeds (*Acacia nilotica*) for use as livestock feed 11 (1984)45
 Kundu, S.S., see Jakhmola, R.C. 19(1988)1
 Kurdi, O.I., see El Hag, M.G. 13(1985)147
 Kyzlink, V., see Kalac, P. 4(1979)81
 Kyzlink, V., see Kalac, P. 5(1980)59

L

- Laber, B.
 Survey of the use of maize for livestock feeding in Austria 1(1976)153
 Lacassagne, L., Francesch, M., Carré, B. and Melcion, J.P.
 Utilization of tannin-containing and tannin-free faba beans (*Vicia faba*) by young chicks: effects of pelleting feeds on energy, protein and starch digestibility 20(1988)59
 Lacey, J., see Lord, K.A. 6(1981)73
 Lacey, J., Lord, K.A. and Cayley, G.R.
 Chemicals for preventing moulding in damp hay 6(1981)323
 Lacey, J., see Lord, K.A. 7(1982)287
 Lacey, J., Lord, K.A. and Cayley, G.R.

- Problems of testing novel chemicals for the preservation of damp hay 8(1983)283
- Lahore, R. and Croke, B.
Energetics of stockfeed production 3(1978)1
- Laksevela, B.
Adjustments of the protein level for slaughter pigs and supplementation with lysine and methionine 1(1976)619
- Lalande, G., see Flipot, P. 11(1984)35
- Lall, D., Lohan, O.P., Vaid, J., Sharma, O.P. and Negi, S.S.
Can lantana seeds be utilized in lamb rations? 9(1983)29
- Lall, D., see Makkar, H.P.S. 20(1988)1
- Lall, D. and Prasad, T.
Compositional quality of certain unconventional calcium and phosphorus sources in India for use as mineral supplements for livestock 23(1989)343
- Lall, D., see Kamra, D.N. 25(1989)149
- Lambertini, F., see Giardini, A. 1(1976)465
- Lanari, D. and Rioni, M.
Rate of growth, feed utilisation and digestibility observed with diets containing ensiled maize grain or maize ears fed alone or in mixtures 1(1976)401
- Lanari, D., see Rioni, M. 1(1976)531
- Lanari, D.
Discussion on session 5 1(1976)545
- Landis, K.M., see Hardin, A.C. 25(1989)99
- Langar, P.N., Bakshi, M.P.S. and Rana, V.K.
Nutritive evaluation of umrol-wheat straw as a basal ration for buffaloes 13(1985)103
- Lanham, J.K., see Coppock, C.E. 18(1987)89
- Larsson, K., see Göransson, L. 7(1982)11
- Lau, M.M. and van Soest, P.J.
Titratable groups and soluble phenolic compounds as indicators of the digestibility of chemically treated roughages 6(1981)123
- Laurent, F., Blanchart, G. and Vignon, B.
Effects of ethanolamine and ethylenediamine treatments on the composition and degradability of wheat straw 13(1985)131
- Law, D.P., see Minson, D.J. 9(1983)221
- Lawlor, M.J. and O'Shea, J.
The effect of ammoniation on the intake and nutritive value of straw 4(1979)169
- Lawrence, T.L.J.
Effects of micronization on the digestibility of whole soya beans and rapeseeds for the growing pig 3(1978)179
- Lawrence, T.L.J.
The effects of cereal particle size and pelleting on the nutritive value of oat-based diets for the growing pig 8(1983)91
- Le Dividich, J., Christon, R., Peiniau, J. and Aumaitre, A.
Proximate chemical analysis of final cane molasses and effect of feeding 30% molasses on intestinal sucrose and maltase activities in the rat 3(1978)15
- Lechtenberg, V.L., see Moore, K.J. 17(1987)109
- Leclercq, B., see Lessire, M. 7(1982)365
- Leclercq, B., see Mohamed, K. 11(1984)199
- Lee, B.D., see Leeson, S. 11(1984)211
- Lee, N.-H., Rooke, J.A. and Armstrong, D.G.
The digestion by sheep of barley and maize-based diets containing either meat and bone meal or soya bean meal 15(1986)301
- Lee, N.H., see Ørskov, E.R. 8(1983)247
- Leeson, S., see Summers, J.D. 3(1978)243
- Leeson, S., see Slinger, S.J. 3(1978)251
- Leeson, S., see Shen, H. 8(1983)303
- Leeson, S., Summers, J.D. and Lee, B.D.
Nutritive value of single-cell protein produced by *Chaetomium cellulolyticum* grown on maize stover and pulp-mill sludge 11(1984)211
- Leeson, S., see Shen, H. 13(1985)57
- Leeson, S., see Peer, D.J. 13(1985)183
- Leeson, S., see Peer, D.J. 13(1985)191
- Leeson, S., see Peer, D.J. 13(1985)203
- Leeson, S., see Summers, J.D. 16(1986/1987)129
- Leeson, S., Hussar, N. and Summers, J.D.
Feeding and nutritive value of hominy and corn grits for poultry 19(1988)313
- Leibholz, J., see Evans, D.F. 4(1979)33
- Leibholz, J., see Evans, D.F. 4(1979)43
- Leibholz, J.
Meat meal in the diet of the early-weaned pig. III. Meat meal quality and the processing of meat meals 4(1979)53
- Leibholz, J., Kellaway, R.C. and Hargreave, G.T.
Effects of sodium chloride and sodium bicarbonate in the diet on the performance of calves 5(1980)309
- Leibholz, J.
Meat meal in the diet of the early-weaned pig. IV. The supplementation of diets with tryptophane, lysine and methionine 7(1982)27
- Leibholz, J., Love, R.J., Mollah, Y. and Carter, R.R.
The absorption of dietary L-lysine and extruded L-lysine in pigs 15(1986)141
- Leibholz, J.M.L., see Kellaway, R.C. 3(1978)201
- Lelong, C.
Maize and animal production in France 1(1976)141
- Lelong, C., see Malterre, C. 1(1976)417
- Lelong, C.
For beef cattle, all forms of maize can be used 1(1976)521
- Lemerle, C., see Moran, J.B. 20(1988)299
- Leng, R.A., see Dryden, G.McL. 14(1986)41
- Leng, R.A., see Sudana, I.B. 16(1986/1987)25
- Leng, R.A., see Perdok, H.B. 17(1987)121
- Leng, R.A., see Dryden, G.McL. 19(1988)121
- Lessire, M. and Leclercq, B.
Metabolisable energy value of fats in chicks and adult cockerels 7(1982)365
- Letcher, A.J., see Orpin, C.G. 10(1983/1984)145

- Leuillet, M., see Gatel, F. 22(1988)91
- Levanon, D., see Silanikove, N. 20(1988)87
- Levy, D., see Holzer, Z. 5(1980)299
- Levy, D., see Holzer, Z. 12(1984/1985)253
- Levy, D., Bar-Tsur, A., Holzer, Z., Samuel, V., Ashbell, G., Lisker, N., Ilan, D., Berkovitz, S. and Kali, J.
High grain content maize silage in fattening diets of young male cattle 16(1986/1987)63
- Lewis, A.J., see Aherne, F.X. 3(1978)235
- Lewis, B.A., see Jeraci, J.L. 23(1989)15
- Lewis, D., see Oguntuna, T. 12(1984/1985)285
- Lewis, S.M., Holzgraebe, D.P., Berger, L.L., Fahey Jr., G.C., Gould, J.M. and Fanta, G.F.
Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter disappearance in sacco 17(1987)179
- Lindberg, J.E., Malmberg, A. and Theander, O.
The chemical composition and nutritive value for ruminants of four possible energy crops and their residues of anaerobic fermentation 15(1986)197
- Lindgren, S., see Kaspersson, A. 19(1988)299
- Lindner, H.P., see Kirchgessner, M. 6(1981)337
- Linnerud, A.C., see Johnson, T.O. 12(1984/1985)65
- Linnerud, A.C., see Ortigues, I. 24(1989)13
- Lisker, N., see Levy, D. 16(1986/1987)63
- Litsky, J., see Pringle, D. 9(1983)89
- Little, D.A., see Mahyuddin, P. 22(1988)69
- Livingstone, R.M., Jones, A.S. and Mennie, I.
Swedes (*Brassica napus*) for growing pigs: chemical composition and use as a replacement for barley in the diet 2(1977)31
- Livingstone, R.M. and Jones, A.S.
The effect of supplementary protein on the performance of pigs given swedes (*Brassica napus*) as a replacement for barley in the diet 2(1977)277
- Livingstone, R.M., Baird, B.A., Atkinson, T. and Crofts, R.M.J.
The effect of different patterns of thermal processing of potatoes on their digestibility by growing pigs 4(1979)295
- Livingstone, R.M., Baird, B.A. and Atkinson, T.
Cabbage (*Brassica oleracea*) in the diet of growing-finishing pigs 5(1980)69
- Lizama, L.C., Marion, J.E. and McDowell, L.R.
Utilization of aquatic plants *Elodea canadensis* and *Hydrilla verticillata* in broiler chick diets 20(1988)155
- Llamas, G.L., see Faulkner, D.B. 12(1984/1985)125
- Lo Bruno, A., see Giardini, A. 1(1976)369
- Lo Bruno, A., see Giardini, A. 1(1976)465
- Lohan, O.P., see Lall, D. 9(1983)29
- Longland, A.C. and Low, A.G.
Digestion of diets containing molassed or plain sugar-beet pulp by growing pigs 23(1989)67
- Lord, K.A., Cayley, G.R. and Lacey, J.
Laboratory application of preservatives to hay and the effects of irregular distribution on mould development 6(1981)73
- Lord, K.A., see Lacey, J. 8(1983)283
- Lord, K.A., Lacey, J. and Cayley, G.R.
Laboratory tests of 8-quinolinol as an additive to volatile fatty acids for preserving hay and other feed-stuffs 7(1982)287
- Lord, K.A., see Lacey, J. 8(1983)283
- Lough, D.S., see Staples, C.R. 23(1989)277
- Love, R.J., see Leibholz, J. 15(1986)141
- Low, A.G.
Secretory response of the pig gut to non-starch polysaccharides 23(1989)55
- Low, A.G., see Longland, A.C. 23(1989)67
- Lowry, J.B., see Mahyuddin, P. 22(1988)69
- Lüdke, H. and Schöne, F.
Copper and iodine in pig diets with high glucosinolate rapeseed meal. I. Performance and thyroid hormone status of growing pigs fed on a diet with rapeseed meal treated with copper sulphate solution or untreated and supplements of iodine, copper or a quinoxaline derivative 22(1988)33
- Lüdke, H., see Schöne, F. 22(1988)45
- Lund, D. and Håkansson, J.
Nutritional and growth studies with pea-crop meals and peas for growing-finishing pigs 16(1986/1987)119
- Lutfi, A.A.A., see Maglad, M.A. 15(1986)69
- Luthra, Y.P., see Das, B. 4(1979)17
- Lynch, G.L., Berger, L.L., Fahey Jr., G.C. and Baker, E.C.
Effects of isopropyl alcohol treatment of soya bean meal on nitrogen utilization by ruminants 14(1986)127

M

- Macdearmid, A., see Ørskov, E.R. 6(1981)273
- Macdearmid, A., see Ørskov, E.R. 6(1981)355
- Macdearmid, A., see Williams, P.E.V. 10(1983/1984)247
- Macdearmid, A., see Williams, P.E.V. 12(1984/1985)1
- Machin, D.H., Hector, D.A., Capper, B.S. and Carter, P.M.
The utilization by broiler chickens of poultry offal hydrolysed in formic acid 11(1984)247
- Machin, D.H., Silverside, D.E., Hector, D.A. and Parr, W.H.
The utilisation by growing pigs of ruminant offal hydrolysed in formic acid 15(1986)273
- Mackay, B.E., see Walker, H.G. 9(1983)283
- Mackie, I.M.
General review of fish protein hydrolysates 7(1982)113
- Mackie, I.M., see Ritchie, A.H. 7(1982)125
- Mackie, R.I., see Westlake, K. 25(1989)169

- MacLean, S.J., see Gregory, K.F. 2(1977)77
- Macleod, G.K., see Smith, O.B. 5(1980)77
- Macleod, G.K., see Summers, J.D. 5(1980)205
- Madasci, J.P., Parrish, F.W. and McNaughton, J.L.
Treatment of low-tannin sorghum grain for broiler feed 20(1988)69
- Madhisetty, M., see Belyea, R.L. 17(1987)57
- Maestre, M.R., see Alibés, X. 8(1983)63
- Maghoub, S.O., see El Hag, M.G. 13(1985)147
- Maglad, M.A., Lutfi, A.A.A. and Gabir, S.
The effect of grinding groundnut hulls either with or without alkali treatment on digestibility of diet and on ruminal and blood components 15(1986)69
- Mahyuddin, P., Little, D.A. and Lowry, J.B.
Drying treatment drastically affects feed evaluation and feed quality with certain tropical forage species 22(1988)69
- Makkar, H.P.S., Lall, D. and Negi, S.S.
Complexes of urea and formaldehyde as non-protein nitrogen compounds in ruminant rations: a review 20(1988)1
- Makkar, H.P.S., see Negi, S.S. 22(1988)79
- Malmberg, A., see Lindberg, J.E. 15(1986)197
- Malossini, F., see Pace, V. 11(1984)125
- Malterre, C. and Lelong, C.
Utilising the maize crop in various forms for beef production 1(1976)417
- Malterre, C., see Geay, Y. 1(1976)429
- Mandel, E., see Alumot, E. 2(1977)77
- Mandell, I.B., Christison, G.I., Nicholson, H.H. and Coxworth, E.C.
The effect of variation in the water content of wheat straw before ammoniation on its nutritive value for beef cattle 20(1988)111
- Manfredini, M. and Cavani, C.
Distillery effluents as animal feed: the use of condensed beet molasses stillage (CBMS) in broiler feeding 5(1980)233
- Mann, M.E., Cohen, R.D.H., Kernan, J.A., Nicholson, H.H., Christensen, D.A. and Smart, M.E.
The feeding value of ammoniated flax straw, wheat straw and wheat chaff for beef cattle 21(1988)57
- Mann, S.O., see Mould, F.L. 10(1983/1984)15
- Mann, S.P., see Orpin, C.G. 21(1988)161
- Manor, G., see Horn, G.W. 8(1983)35
- Mansbridge, R., see Ibbotson, C.F. 10(1983/1984)223
- Manten, A.A. and Greenhalgh, J.F.D.
Titles of scientific papers 2(1977)1
- Manten, A.A.
Citation decay in animal science 2(1977)195
- Mariani, P., see Russo, V. 1(1976)25
- Marion, J.E., see Lizama, L.C. 20(1988)155
- Marlett, J.A.
Measuring dietary fiber 23(1989)1
- Marquardt, R.R., see Rotter, B.A. 24(1989)233
- Marti, J., see Cervera, C. 12(1984/1985)233
- Martinez Pascual, J. and Fernández Carmona, J.
Composition of citrus pulp 5(1980)1
- Martinez Pascual, J. and Fernández Carmona, J.
Citrus pulp in diets for fattening lambs 5(1980)11
- Martinez Pascual, J. and Fernández Carmona, J.
Citrus pulp in diets for fattening rabbits 5(1980)23
- Martinez Pascual, J.L., see Deltoro López, J. 6(1981)163
- Martinez Pascual, J.L., see Deltoro López, J. 6(1981)169
- Martz, F.A., see Belyea, R.L. 17(1987)57
- Mascarenhas Ferreira, A., Kerstens, J. and Gast, C.H.
The study of several modifications of the neutral detergent fibre procedure 9(1983)19
- Mascarenhas Ferreira, A., see Van Soest, P.J. 10(1983/1984)155
- Mascarenhas Ferreira, A., see Dias-da-Silva, A.A. 19(1988)67
- Mascarenhas-Ferreira, A., Guedes, C.V.M. and Dias-da-Silva, A.A.
Effects of urea treatment on chemical composition and in vitro digestibility on meadow hays of Northern Portugal 25(1989)157
- Maslowska, J. and Bazylak, G.
Thermofractographic determination of monocarbonyl compounds in animal waste fats used as feed fat 13(1985)227
- Mason, V.C., see Tetlow, R.M. 18(1987)257
- Mason, V.C., see Tetlow, R.M. 18(1987)271
- Mason, V.C., see Deschard, G. 18(1987)283
- Mason, V.C., see Deschard, G. 19(1988)55
- Mason, V.C., Hartley, R.D., Keene, A.S. and Cobby, J.M.
The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility in vitro and cell wall degradability 19(1988)159
- Mason, V.C., Keene, A.S., Cook, J.E., Cooper, E.M. and Hartley, R.D.
Oven and stack ammoniation of grass hays. 1. Changes in chemical composition in relation to digestibility in vitro and cell-wall degradability 24(1989)299
- Mason, V.C., Cook, J.E., Cooper, E.M., Hoadley, C.J. and Cockburn, J.E.
Oven and stack ammoniation of grass hays. 2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313
- Mason, W., see Flipot, P. 11(1984)35
- Masoud, S., see Abouheif, M.A. 13(1985)215
- Mathison, G.W., see Gibson, D.M. 19(1988)203
- May, P.J. and Barker, D.J.
Milling barley and lupin grain in diets for cattle 12(1984/1985)57
- Mbatya, P.B.A., see Petchey, A.M. 2(1977)315
- Mbatya, P.B.A., Kay, M. and Smart, R.I.

- Methods of improving the utilization of cereal straw by ruminants. I. Supplements of urea, molasses and dried grass and treatment with sodium hydroxide 8(1983)221
- Mbatya, P.B.A., Kay, M., Smart, R.I. and Milne, J.
Methods of improving the utilization of cereal straw by ruminants. II. Further studies on dried grass as a supplement for lambs 8(1983)229
- Mbatya, P.B.A.
Methods of improving the utilization of cereal straw by ruminants. III. A note on the effect of ensiling straw treated with urea 9(1983)181
- Mbatya, P.B.A., Kay, M., Smart, R.I. and Kennedy, S.
Methods of improving the utilization of cereal straw by ruminants. IV. The effect on intake of level of supplementation with urea and dried grass 13(1985)281
- Mbatya, P.B.A., Kay, M., Smart, R.I. and Kennedy, S.
Methods of improving the utilization of cereal straw by ruminants. V. The effect of molasses and dried grass 13(1985)293
- Mbugi, P.K., Ingalls, J.R. and Sharma, H.R.
Evaluation of pea protein concentrate as a source of protein in milk replacers for Holstein calves 24(1989)267
- McAllan, A.B. and Griffith, E.S.
The effects of different sources of nitrogen supplementation on the digestion of fibre components in the rumen of steers 17(1987)65
- McBurney, M.I. and Thompson, L.U.
Dietary fiber and energy balance: integration of the human ileostomy and in vitro fermentation models 23(1989)261
- McCarthy, D.B., see Bunting, L.D. 22(1988)61
- McCarthy, R.E., see Salyers, A.A. 23(1989)109
- McClure, W.H., see Ortigues, I. 24(1989)13
- McCrae, F., see Farrell, D.J. 9(1983)49
- McDonald, P., see Henderson, A.R. 7(1982)303
- McDonald, P., Proven, M.J. and Henderson, A.R.
The effect of some pre-ensiling treatments on silage composition and nitrogen disappearance in the rumen 8(1983)259
- McDonald, P., see Heron, S.J.E. 19(1988)85
- McDowell, L.R., see Paz, E.A. 11(1984)75
- McDowell, L.R., see Lizama, L.C. 20(1988)155
- McElhiney, R.R. and Tangprasertchai, P.
The effect of dilution levels in premixes on micro-ingredient dispersion in animal feeds 8(1983)139
- McGilliard, A.D., see Adamu, A.M. 22(1988)227
- McGinn, R., see McLellan, A.R. 6(1981)245
- McGinn, R., see McLellan, A.R. 9(1983)107
- McGinn, R. and McLellan, A.R.
Energy metabolism in lambs fed silage diets with sucrose or a growth promoter 16(1986/1987)203
- McIntosh, A.D., see Davidson, J. 7(1982)217
- McKenzie, R.J., see Farrell, D.J. 9(1983)49
- McKirdy, J.A., see Sharma, H.R. 10(1983/1984)77
- McLaughlin, G.L., see Horn, G.W. 8(1983)35
- McLellan, A.R. and McGinn, R.
A calorimetric investigation of the nutritive value of formic acid- and formaldehyde-formic acid-treated grass silages 6(1981)245
- McLellan, A.R. and McGinn, R.
Energy and nitrogen balance, intake and growth rate of lambs fed on unweaned grass silages treated with a formaldehyde-acid mixture or untreated 9(1983)107
- McLellan, A.R., see McGinn, R. 16(1986/1987)203
- McLeod, M.N. and Minson, D.J.
The analytical and biological accuracy of estimating the dry matter digestibility of different legume species 1(1976)61
- McLeod, M.N. and Minson, D.J.
The accuracy of the pepsin-cellulase technique for estimating the dry matter digestibility in vivo of grasses and legumes 3(1978)277
- McLeod, M.N. and Minson, D.J.
A note on Onozuka 3S cellulase as a replacement for Onozuka SS (P1500) cellulase when estimating forage digestibility in vitro 5(1980)347
- McLeod, M.N. and Minson, D.J.
Accuracy of predicting digestibility by the cellulase technique; the effect of pretreatment of forage samples with neutral detergent or acid pepsin 7(1982)83
- McLeod, M.N. and Wilson, J.R.
The effect of method of preservation of forages on their resistance to reduction in particle size with an artificial masticator 24(1989)185
- McMeniman, N.P. and Armstrong, D.G.
Nitrogen levels in low-roughage diets for efficient rumen microbial protein production 2(1977)255
- McMeniman, N.P., Elliott, R. and Ash, A.J.
Supplementation of rice straw with crop by-products. I. Legume straw supplementation 19(1988)43
- McNaughton, J.L., see Madasci, J.P. 20(1988)69
- McNew, R.W., see Zorrilla-Rios, J. 22(1988)305
- McReynolds, W.E., see Rule, D.C. 15(1986)113
- Melcion, J.P., see Lacassagne, L. 20(1988)59
- Mennie, I., see Livingstone, R.M. 2(1977)31
- Mensah-Asante, B., see Donkoh, A. 22(1988)139
- Mergounis, D.
Use of maize for ruminants and pig feeding in Greece 1(1976)125

- Merritt, J.H.
Assessment of the production costs of fish protein hydrolysates 7(1982)147
- Metwally, H., see Abouheif, M.A. 13(1985)215
- Meyer, D.W., see Sanderson, M.A. 14(1986)221
- Michalet-Doreau, B., see Aufrere, J. 20(1988)203
- Michie, W., see Cowan, P.J. 4(1979)125
- Michie, W., see Rose, S.P. 11(1984)221
- Mika, V.
Some remarks on the determination of acid-detergent fibre in feeds with cetyltrimethylammonium bromide versus determination of detergent fibre with alkylarylsulfonic acid 6(1981)91
- Miller, C., see Caffrey, P.J. 19(1988)231
- Miller, M.E., see Wohlt, J.E. 6(1981)115
- Milliken, G., see Hinds, M.A. 12(1984/1985)205
- Milne, E., see Davidson, J. 7(1982)217
- Milne, J., see Mbatya, P.B.A. 8(1983)229
- Minson, D.J., see McLeod, M.N. 1(1976)61
- Minson, D.J., see Goto, I. 2(1977)247
- Minson, D.J., see McLeod, M.N. 3(1978)277
- Minson, D.J., see McLeod, M.N. 5(1980)347
- Minson, D.J.
A flexible, two-component feeding system derived from Blaxter's three-component, metabolisable-energy system 6(1981)223
- Minson, D.J., see McLeod, M.N. 7(1982)83
- Minson, D.J.
Feeding standards are necessary 7(1982)423
- Minson, D.J., Butler, K.L., Grummitt, N. and Law, D.P.
Bias when predicting crude protein, dry matter digestibility and voluntary intake of tropical grasses by near-infrared reflectance 9(1983)221
- Mirgani, T.
Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by goats 17(1987)219
- Miron, J., see Ben-Ghedalia, D. 10(1983/1984)269
- Miron, J., see Ben-Ghedalia, D. 19(1988)219
- Miron, J., see Ben-Ghedalia, D. 24(1989)289
- Mitaru, B.N., see Baidoo, S.K. 18(1987)37
- Mitaru, B.N., see Baidoo, S.K. 18(1987)45
- Mitchell, K.G., see Barber, R.S. 2(1977)161
- Mitchell, K.G., see Barber, R.S. 3(1978)163
- Mitchell, K.G., see Barber, R.S. 4(1979)117
- Mitchell, K.G., see Barber, R.S. 4(1979)233
- Mitchell, K.G., see Barber, R.S. 5(1980)215
- Mitchell, K.G., see Barber, R.S. 6(1981)35
- Mitchell Jr., E.D., see Hibberd, C.A. 15(1986)231
- Młodkowski, M., see Hanczakowski, P. 6(1981)413
- Mngulwi, J.G.J., see Kategile, J.A. 20(1988)97
- Modler, H.W., see Elliot, J.I. 17(1987)213
- Moermans, R., see Eeckhout, W. 6(1981)367
- Moermans, R.J., see Aerts, J.V. 3(1978)309
- Moermans, R.J., see Cottyn, B.G. 11(1984)137
- Mohamed, K., Leclercq, B., Anwar, A., El-Alaily, H. and Soliman, H.
A comparative study of metabolisable energy in ducklings and domestic chicks 11(1984)199
- Mohammed, T.A., see Osman, A.G. 19(1988)351
- Mohammed, T.A. and Ali, O.M.
Effects of wood ash extract treatment on the feeding value and utilization of high-tannin sorghums by broiler chicks 22(1988)131
- Moir, K.W. and Connor, J.K.
A comparison of three fibre methods for predicting the metabolizable energy content of sorghum grain for poultry 2(1977)197
- Moisio, T. and Heikonen, M.
A titration method for silage assessment 22(1988)341
- Molina, B.T., see Djajanegara, A. 12(1984/1985)141
- Molina, E., Boza, J. and Aguilera, J.F.
Nutritive value for ruminants of sugar cane bagasse ensiled after spray treatment with different levels of NaOH 9(1983)1
- Molina, E., see Aguilera, J.F. 12(1984/1985)171
- Mollah, Y., see Leibholz, J. 15(1986)141
- Mølle, K.R. G.
Maize as a forage crop in Denmark 1(1976)183
- Molnar, S., see Greife, H.A. 5(1980)241
- Montague Jr., W.C., see Walker, H.G. 9(1983)283
- Monteagudo, A., see Gonzalez, G. 1(1976)129
- Monteagudo, A.
Cultivation of maize in irrigated areas for silage and grain production and their use by ruminants 1(1976)237
- Montgomery, R.R., Nofsinger, G.W. and Bothast, R.J.
Preservation of high-moisture maize - a comparison of gaseous and liquid anhydrous ammonia with methylene-bis-propionate 5(1980)337
- Moo-Young, M. and Chahal, D.S.
Utilization of cattle manure for single-cell protein production with *Chaetomium cellulolyticum* 4(1979)199
- Moo-Young, M., Chahal, D.S. and Vlach, D.
Utilization of cattle manure for Torula yeast production from straw hydrolysates 5(1980)175
- Moo-Young, M., see Chavez, E.R. 22(1988)1
- Moo-Young, M., see Chavez, E.R. 22(1988)13
- Moo-Young, M., see Chavez, E.R. 22(1988)23
- Moore, K.J. and Lechtenberg, V.L.
Chemical composition and digestion in vitro of orchardgrass hay ammoniated by different techniques 17(1987)109
- Mora, M.I., Fahey Jr., G.C., van der Aar, P.J. and Berger, L.L.
Improving utilization of crop residues. Eval-

- uation of two chemicals used to improve digestibility of crop residues 9(1983)205
- Moran, J.B., see Thompson, C. 16(1986/1987) 225
- Moran, J.B., Lemerle, C. and Trigg, T.E. The intake and digestion of maize silage-based diets by dairy cows and sheep 20 (1988)299
- Morel, M.T., see Durand, M. 21(1988)197
- Morgan, C.A. and Whittemore, C.T. Energy evaluation of feeds and compounded diets for pigs - a review 7(1982)387
- Morgan, C.A., Whittemore, C.T. and Cockburn, J.H.S. The effect of level and source of protein, fibre and fat in the diet on the energy value of compounded pig feeds 11(1984)11
- Morgan, C.A., Whittemore, C.T., Phillips, P. and Crooks, P. The prediction of the energy value of compounded pig foods from chemical analysis 17(1987)81
- Morgan, C.A. and Whittemore, C.T. Dietary fibre and nitrogen excretion and retention by pigs 19(1988)185
- Morgan, C.A., Whittemore, C.T. and Taylor, A. G. Body composition and growth of young pigs as affected by protein source and a growth promoter 24(1989)219
- Morita, Z., see Sekine, J. 25(1989)207
- Morris, J.G., see Sidahmed, A.E. 8(1983)47
- Morrison, I.M. and Brice, R.E. The digestion of untreated and ammonia-treated barley straw in an artificial rumen 10(1983/1984)229
- Morrison, I.M., see Robinson, P.H. 18(1987)75
- Mosi, A., see Butterworth, M.H. 16(1986/1987)99
- Mosi, A.K. and Butterworth, M.H. The voluntary intake and digestibility of combinations of cereal crop residues and legume hay for sheep 12(1984/1985)241
- Moss, B.R., El-Negoumy, A.M. and Newman, C.W. Nutritional value of waxy barleys for broiler chickens 8(1983)25
- Moughan, P.J., see Kies, A.K. 16(1986/1987) 169
- Moughan, P.J., see Smith, W.C. 18(1987)143
- Moughan, P.J., see Smith, W.C. 19(1988)105
- Moughan, P.J., Wilson, M.N., Smits, C.H.M. and Smith, W.C. An evaluation of skim milk powders subjected to different heating conditions during processing as dietary protein sources for the young pig 22(1988)203
- Mould, F.L. and Ørskov, E.R. Manipulation of rumen fluid pH and its influence on cellulolysis in sacco, dry matter degradation and the rumen microflora of sheep offered either hay or concentrate 10 (1983/1984)1
- Mould, F.L., Ørskov, E.R. and Mann, S.O. Associative effects of mixed feeds. I. Effects of type and level of supplementation and the influence of the rumen fluid pH on cellulolysis in vivo and dry matter digestion of various roughages 10(1983/1984)15
- Mould, F.L., Ørskov, E.R. and Gaud, S.A. Associative effects of mixed feeds. II. The effect of dietary addition of bicarbonate salts on the voluntary intake and digestibility of diets containing various proportions of hay and barley 10(1983/1984)31
- Mowat, D.N., see Oji, U.I. 4(1979)177
- Mowat, D.N., see Oji, U.I. 4(1979)187
- Mshelia, T.A., see Kumaresan, A. 11(1984)45
- Mugerwa, J.S., see Odwongo, W.O. 5(1980) 193
- Muindi, P.J., Thomke, S. and Hanssen, J.F. Protein quality studies on rats fed on cassava-root meal enriched with *Cephalosporium eichhorniae* 152 or with conventional plant-protein supplements 6(1981)197
- Müller, H.L., see Kreuzer, M. 16(1986/1987) 233
- Muñoz, F., see Alibés, X. 8(1983)63
- Muñoz, F., see Alibés, X. 10(1983/1984)239
- Muñoz, F., see Alibés, X. 11(1984)189
- Murdock, F.R. and Kromann, R.P. The digestibility and nutritive value for dairy heifers of rations containing cellulose fibre 2(1977)287
- Murphy, G.E., see Hardin, A.C. 25(1989)99
- Mussa, M.A., see Sarwatt, S.V. 22(1988)237
- Myer, R.O. and Gorbet, D.W. Waxy and normal grain sorghums with varying tannin contents in diets for young pigs 12(1984/1985)179
- Myer, R.O., Barnett, R.D., Cornell, J.A. and Combs, G.E. Nutritive value of diets containing tritcale and varying mixtures of tritcale and maize for growing-finishing swine 22(1988)217
- Mzihirwa, Y.G., see Kategile, J.A. 6(1981)133
- ## N
- Nangole, F.N., Kayongo-Male, H. and Said, A.N. Chemical composition, digestibility and feeding value of maize cobs 9(1983)121
- Napoli, G.M. and Santini, F.J. The effect of a protein energy supplement on pasture protein and fibre digestion in the rumen of grazing steers 25(1989)39
- Narahari, D. and Kothandaraman, P. The influence of processing and storage on hydrogen cyanide and tannin contents of para-rubber seed and its products 9(1983) 319
- Narahari, D. and Kothandaraman, P. Chemical composition and nutritional value of para-rubber seed and its products for chickens 10(1983/1984)257
- Narasimhalu, P.

- A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum-pepsin treatment for predicting digestibility and intake of hays in sheep 12(1984/1985)101
- Narasimhalu, P., Graham, H. and Aman, P. Effects of ozonation on the chemical composition and in vitro degradation of triticales internodes and red clover stems 24(1989)159
- Natanam, R., Kadirvel, R. and Balagopal, R. The effect of kernels of karanja (*Pongamia glabra* Vent) on growth and feed efficiency in broiler chicks to 4 weeks of age 25(1989) 201
- Neale, R.J., see Oguntona, T. 12(1984/1985) 285
- Nefzaoui, A. and Vanbelle, M. Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139
- Negi, S.S. Tannins in sal seed (*Shorea robusta*) and sal seed meal limit their utilization as livestock feeds 7(1982)161
- Negi, S.S., see Lall, D. 9(1983)29
- Negi, S.S., see Makkar, H.P.S. 20(1988)1
- Negi, S.S., Singh, B. and Makkar, H.P.S. Rumen degradability of nitrogen in typical cultivated grasses and leguminous fodders 22(1988)79
- Neill, A.R., Bell, S.L. and Williams, K.C. The dye absorption test as an indicator of the nutritive quality of heat-processed navy bean 9(1983)71
- Nelson, M.L. and Finley, J.W. Effect of soft white wheat addition to alfalfa-grass forage on heifer gain, diet digestibility and in vitro digestion kinetics 24 (1989)141
- Nes, I.F., see Skrede, A. 20(1988)287
- Neskar, M., see Rotter, B.A. 24(1989)233
- Newman, C.W., see Moss, B.R. 8(1983)25
- Nicholson, H.H., see Horton, G.M.J. 7(1982)1
- Nicholson, H.H., see Mandell, I.B. 20(1988) 111
- Nicholson, H.H., see Mann, M.E. 21(1988)57
- Nikolić, J.A. and Jovanović, M. Some properties of apple pomace ensiled with and without additives 15(1986)57
- Njoku, P.C., see Okoh, P.N. 7(1982)359
- Njoku, P.C. Effect of dietary ascorbic acid (vitamin C) supplementation on the performance of broiler chickens in a tropical environment 16 (1986/1987)17
- Nofsinger, G.W., see Montgomery, R.R. 5 (1980)337
- Noland, P.R., Campbell, D.R., Sharp, R.N. and Johnson, Z.B. Influence of pericarp and endosperm colour and type on digestibility of grain sorghum by pigs 2(1977)219
- Noland, P.R., see Sharp, R.N. 2(1977)327
- Noland, P.R., Campbell, D.R. and Johnson, Z.B. Use of unextracted sunflower seeds as a protein source 5(1980)51
- Nordkvist, E., Graham, H. and Aman, P. Degradation in vitro and in sacco of red clover leaves and stems 17(1987)295
- Nwoko, E. and Oji, U.I. Variation in metabolizable energy content of raw or autoclaved white and brown varieties of three tropical grain legumes 13 (1985)141
- Nwoko, E. A comparison of metabolizable energy content of eight common feed ingredients determined with young guinea fowls (keets) and pullet chicks 15(1986)1
- ## O
- Obilana, A.B., see Alhassan, W.S. 17(1987) 285
- Obilana, A.T., see Okoh, P.N. 7(1982)359
- Ocio, E., Viñaras, R. and Rey, J.Ma. House fly larvae meal grown on municipal organic waste as a source of protein in poultry diets 4(1979)227
- Odebumi, J.O., see Babatunde, G.M. 4 (1979)23
- Oduye, O.O., see Babatunde, G.M. 4(1979)23
- Odwongo, W.O. and Mugerwa, J.S. Performance of calves on diets containing *Amaranthus* leaf meal 5(1980)193
- Offer, N.W., see Husain, R.A.K. 16(1986/ 1987)297
- Offer, N.W. and Husain, R.A.K. Fish silage as a protein supplement for early-weaned calves 17(1987)165
- Offer, N.W., see Hyslop, J.J. 25(1989)55
- Ogle, R.B., see Göransson, L. 12(1984/1985) 159
- Oguntona, T., Neale, R.J. and Lewis, D. The effect of pelleting on the utilisation of yeast grown on n-alkanes by broiler chicks 12(1984/1985)285
- Ohajuruka, O.A. and Palmquist, D.L. Response of high-producing dairy cows to high levels of dried corn gluten feed 24 (1989)191
- Ohlde, G. and Becker, K. Suitability of cell-wall constituents as predictors of organic matter digestibility in some tropical and subtropical by-products 7(1982)191
- Oji, U.I. and Mowat, D.N. Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979) 177
- Oji, U.I., Mowat, D.N. and Buchanan-Smith, J.G. Nutritive value of thermoammoniated and

- steam-treated maize stover II. Rumen metabolites and rate of passage 4(1979)187
- Oji, U.I., see Nwoko, E. 13(1985)141
- Oke, O.L.
Problems in the use of cassava as animal feed 3(1978)345
- Okoh, P.N., Obilana, A.T., Njoku, P.C. and Aduku, A.O.
Proximate analysis, amino acid composition and tannin content of improved Nigerian sorghum varieties and their potential in poultry feeds 7(1982)359
- Oldham, J.D., see Phipps, R.H. 4(1979)163
- Olsson, O., see Göransson, L. 7(1982)11
- Omole, T.A. and Onwudike, O.C.
Investigations of the treatment of sawdust for rabbit feeding. 1. Effect of sodium hydroxide treatment 6(1981)43
- Omole, T.A. and Onwudike, O.C.
Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51
- Omole, T.A., see Sonaiya, E.B. 8(1983)211
- Onesiroso, P., see Smith, O.B. 11(1984)231
- Onwudike, O.C., see Omole, T.A. 6(1981)43
- Onwudike, O.C., see Omole, T.A. 6(1981)51
- Onwudike, O.C.
Palm kernel meal as a feed for poultry. 1. Composition of palm kernel meal and availability of its amino acids to chicks 16(1986/1987)179
- Onwudike, O.C.
Palm kernel meal as a feed for poultry. 2. Diets containing palm kernel meal for starter and grower pullets 16(1986/1987)187
- Onwudike, O.C.
Palm kernel meal as a feed for poultry. 3. Replacement of groundnut cake by palm kernel meal in broiler diets 16(1986/1987)195
- Onwudike, O.C.
Palm kernel meal as a feed for poultry. 4. Use of palm kernel meal by laying birds 20(1988)279
- Opstvedt, J., Søbstad, G. and Hansen, P.
Fish protein concentrates for pre-ruminant calves: effects of processing temperatures and source of fish on protein digestibility and biological value 18(1987)181
- Oram, R.N., see Green, A.G. 9(1983)271
- Orpin, C.G.
The role of ciliate protozoa and fungi in the rumen digestion of plant cell walls 10(1983/1984)121
- Orpin, C.G. and Letcher, A.J.
Effect of absence of ciliate protozoa on rumen fluid volume, flow rate and bacterial populations in sheep 10(1983/1984)145
- Orpin, C.G., Hazlewood, G.P. and Mann, S.P.
Possibilities of the use of recombinant-DNA techniques with rumen micro-organisms 21(1988)161
- Ørskov, E.R. and Grubb, D.A.
The use of whole barley diets fortified with solutions of urea, minerals and vitamins for lambs 2(1977)307
- Ørskov, E.R., Macdearmid, A., Grubb, D.A. and Innes, G.M.
Utilization of alkali-treated grain. 1. Alkali-treated grain in complete diets for steers and lambs 6(1981)273
- Ørskov, E.R., see Barnes, B.J. 6(1981)347
- Ørskov, E.R., Barnes, B.J., Macdearmid, A., Williams, P.E.V. and Innes, G.M.
Utilization of alkali-treated grain. 3. Utilization by steers of NaOH-treated and rolled barley in silage-based diets 6(1981)355
- Ørskov, E.R., Soliman, H.S. and Clark, C.F.S.
Use of fish protein hydrolysate in milk replacers 7(1982)135
- Ørskov, E.R., Reid, G.W., Holland, S.M., Tait, C.A.G. and Lee, N.H.
The feeding value for ruminants of straw and whole-crop barley and oats treated with anhydrous or aqueous ammonia or urea 8(1983)247
- Ørskov, E.R., see Mould, F.L. 10(1983/1984)1
- Ørskov, E.R., see Mould, F.L. 10(1983/1984)15
- Ørskov, E.R., see Mould, F.L. 10(1983/1984)31
- Ørskov, E.R., see Istasse, L. 15(1986)167
- Ørskov, E.R., see Istasse, L. 16(1986/1987)37
- Ørskov, E.R., see Silva, A.T. 19(1988)277
- Ørskov, E.R., see Silva, A.T. 19(1988)289
- Ørskov, E.R., see Bhargava, P.K. 22(1988)295
- Ortigues, I., Crickenberger, R.G., Fontenot, J.P., Harvey, R.W., McClure, W.H. and Linnerud, A.C.
Distribution of liquid supplement injected into large bales of roughage and comparison of nitrogen supplements for stocker cattle fed low-quality roughages 24(1989)13
- Osafo, E.L.K., see Smith, O.B. 20(1988)189
- Osborn, D.F.
Principles governing the use of chemical methods for assessing the nutritive value of forages: a review 3(1978)265
- Osborn, D.F., Terry, R.A., Spooner, M.C. and Tetlow, R.M.
Use of processing to explore the factors affecting the digestion of forage cell walls 6(1981)387
- Osei, S.A. and Twumasi, I.K.
Effects of oven-dried cassava peel meal on the performance and carcass characteristics of broiler chickens 24(1989)247
- O'Shea, J., see Lawlor, M.J. 4(1979)169
- O'Shea, J. and Baldwin, A.
Evaluation of the increase in feeding value of barley straw following mild treatment with sulphur dioxide 15(1986)13
- Oslage, H.J., see Schulz, E. 1(1976)9
- Osman, A.G., El Tayeb, A.E., Sulieman, A.H. and Mohammed, T.A.
Effect of sorghum straw alone or in combination with molasses and nitrogen sources

- on performance of Sudan desert lambs 19 (1988)351
- Otwinowska, A., see Kowalczyk, J. 7(1982)225
- Oura, R., see Sekine, J. 25(1989)207
- Ovejero, F.M.A., see Fay, J.P. 16(1986/1987) 161
- Overend, M.A., see Rooke, J.A. 13(1985)255
- Owen, B.D., see Beames, R.M. 15(1986)285
- Owen, E., see Alawa, J.P. 11(1984)149
- Owen, E., see Alawa, J.P. 12(1984/1985)215
- Owen, E., see Wrathall, J.H.M. 24(1989)57
- Owens, F.N., see Goetsch, A.L. 18(1987)151
- ## P
- Pace, V., Barge, M.T., Settineri, D. and Malossini, F.
Comparison of forage digestibility in vitro with enzymic solubility 11(1984)125
- Pahlow, G., see Weinberg, Z.G. 20(1988)335
- Palmer, A.Z., see Potter, S.G. 12(1984/1985) 275
- Palmquist, D.L., see Ohajuruka, O.A. 24 (1989)191
- Pandit, N.N., see Singh, A.P. 3(1978)299
- Pandita, N.N., see Singh, A.P. 9(1983)143
- Panditharatne, S., see Jayasuriya, M.C.N. 3 (1978)219
- Panigrahi, S., Bland, B. and Carlaw, P.M.
The nutritive value of tamarind seeds for broiler chicks 22(1988)285
- Papadopoulos, G. and Ziras, E.
Nutrient composition of Greek cottonseed meal 18(1987)295
- Papadopoulos, G., Fegeros, K. and Ziras, E.
Evaluation of Greek cottonseed meal. 2. Use in rations for fattening pigs 18(1987)303
- Papadopoulos, M.C., El Boushy, A.R., Roodbeen, A.E. and Ketelaars, E.H.
Effects of processing time and moisture content on amino acid composition and nitrogen characteristics of feather meal 14(1986) 279
- Papadopoulos, M.C.
The effect of enzymatic treatment on amino acid content and nitrogen characteristics of feather meal 16(1986/1987)151
- Parigi-Bini, R. and Chiericato, G.M.
Influence of ensiling maize ears and grain on chemical composition, conservation losses and digestibility 1(1976)301
- Parkins, J.J., see Bass, J.M. 7(1982)93
- Parkins, J.J., see Hemingway, R.G. 15(1986) 123
- Parr, W.H., see Machin, D.H. 15(1986)273
- Parrish, F.W., see Madasci, J.P. 20(1988)69
- Parthasarathy, M. and Pradhan, K.
Fermentation characteristics and feeding value of ensiled poultry litter containing wheat straw, bagasse or sawdust 7(1982)341
- Partridge, G.G., Allan, S.J. and Findlay, M.
Studies on the nutritive value of roots, cabbage and grass silage for growing commercial rabbits 13(1985)299
- Partridge, G.G., Findlay, M. and Fordyce, R.A.
Fat supplementation of diets for growing rabbits 16(1986/1987)109
- Partridge, I.G., see Barber, R.S. 4(1979)233
- Partridge, I.G., see Barber, R.S. 5(1980)215
- Partridge, I.G., see Barber, R.S. 6(1981)35
- Partridge, I.G.
The digestion of diets containing manioc by young growing pigs 12(1984/1985)119
- Patchell, M.R., see Smith, W.C. 19(1988)105
- Paterson, J.A., see Koeln, L.L. 11(1984)181
- Paterson, J.A., see Goedecken, F.K. 13(1985) 47
- Pathak, N.N., see Jain, V.K. 5(1980)315
- Pathak, N.N., see Kamra, D.N. 25(1989)149
- Patterson, D.C. and Walker, N.
The use of effluent from grass silage in the diet of finishing pigs. I. Variation in composition of effluents 4(1979)263
- Patterson, D.C. and Walker, N.
The use of effluent from grass silage in the diet of finishing pigs. II. Assessment of nutritive value of fresh and stored effluent 4 (1979)275
- Payne, E., see Wright, D.E. 2(1977)93
- Paz, E.A., McDowell, L.R., Conrad, J.H. and Damron, B.L.
Biological availability of phosphorus from basic steel slag 11(1984)75
- Pearce, G.R., see Hendrosoekarjo, S. 3(1978) 31
- Pearce, G.R., see Stanogins, G. 3(1978)155
- Pearce, G.R., see Jayasuriya, M.C.N. 8(1983) 271
- Pearson, A.W., see Fenwick, G.R. 6(1981)421
- Pearson, A.W., Greenwood, N.M., Butler, E.J., Curl, C.L. and Fenwick, G.R.
The involvement of trimethylamine oxide in fish meal in the production of egg taint 8 (1983)119
- Pearson, G. and Carr, J.R.
Lupin-seed meal (*Lupinus angustifolius* cv. Uniwhite) as a protein supplement to barley-based diets for growing pigs 1(1976) 631
- Pearson, G. and Carr, J.R.
A comparison between meals prepared from the seeds of different varieties of lupin as protein supplements to barley-based diets for growing pigs 2(1977)49
- Pearson, G., see Smith, W.C. 18(1987)143
- Pearson, G., see Smith, W.C. 19(1988)105
- Peck, N.J., see Ashes, J.R. 3(1978)109
- Peck, N.J., see Farrell, D.J. 9(1983)99
- Pedroza, C., see Guardiola, C.M. 8(1983)129
- Peer, D.J. and Leeson, S.
Feeding value of hydroponically sprouted barley for poultry and pigs 13(1985)183
- Peer, D.J. and Leeson, S.
Nutrient content of hydroponically sprouted barley 13(1985)191
- Peer, D.J. and Leeson, S.

- Nutrient and trypsin inhibitor content of hydroponically sprouted soya beans 13 (1985)203
- Peers, D.G., Taylor, A.G. and Whittemore, C.T.
The influence of feeding level and level of dietary inclusion on the digestibility of barley meal in the pig 2(1977)41
- Peiniau, J., see Le Dividich, J. 3(1978)15
- Peiniau, J., see Aumaitre, A. 24(1989)275
- Pendulum, L.C., Boling, J.A. and Ely, D.G.
Abomasal and plasma nitrogen constituents of steers fed on different physical forms of shelled maize 2(1977)225
- Penning, P.D., Corcuera, P. and Treacher, T.T.
Effect of dry-matter concentration of milk substitute and method of feeding on intake and performance by lambs 5(1980)321
- Perdok, H.B., see Jayasuriya, M.C.N. 7(1982) 201
- Perdok, H.B. and Leng, R.A.
Hyperexcitability in cattle fed ammoniated roughages 17(1987)121
- Perera, H.G.D., see Jayasuriya, M.C.N. 7 (1982)221
- Perez, J.M. and Aumaitre, A.
Waxy versus regular maize: energy value for growing pigs and utilization in piglet diets 4(1979)109
- Petchev, A.M. and Mbatya, P.B.A.
The intake of steers and heifers fed on mixtures of alkali-treated or untreated straw with grass silage 2(1977)315
- Petchev, A.M.
Performance of calves fed on milk replacers containing fish protein hydrolysate 7(1982) 141
- Peterson, C.L., see Thomas, V.M. 11(1984)269
- Peteva-Vantcheva, Z. and Platikanov, N.
Forage maize in the rations of dairy cows and fattening cattle in Bulgaria 1(1976)45
- Peteva-Vantcheva, Z. and Jeleu, K.
Maize silage and whole maize plant pellets as the basic forages in complete feeds for milk cows 3(1978)23
- Pettersson, D., Hesselman, K. and Åman, P.
Nutritional value for chickens of dried distillers-spent-grain from barley and de-hulled barley 17(1987)145
- Pettersson, D. and Åman, P.
Effects of enzyme supplementation of diets based on wheat, rye or triticale on their productive value for broiler chickens 20(1988) 313
- Petzel, M., see Hanczakowski, P. 4(1979)91
- Pfeffer, E., see Abate, A. 14(1986)1
- Pfugfelder, R.L., Rooney, L.W. and Schake, L.M.
The role of germination in sorghum reconstitution 14(1986)243
- Phillips, P., see Morgan, C.A. 17(1987)81
- Phipps, R.H. and Weller, R.F.
Effect of agronomic factors on the yield of maize and on its composition in relation to its use by ruminants 1(1976)251
- Phipps, R.H. and Oldham, J.D.
Amino acid composition of maize silage produced in the United Kingdom 4(1979)163
- Pike, D.J., see Wrathall, J.H.M. 24(1989)57
- Pirie, R. and Greenhalgh, J.F.D.
Alkali treatment of straw for ruminants. I. Utilization of complete diets containing straw by beef cattle 3(1978)143
- Pirie, R., see Greenhalgh, J.F.D. 3(1978)289
- Pittman, R.J., see Barber, R.S. 3(1978)163
- Pittman, R.J., see Barber, R.S. 4(1979)233
- Pittman, R.J., see Barber, R.S. 5(1980)215
- Platikanov, N., see Peteva-Vantcheva, Z. 1 (1976)45
- Playne, M.J.
Differences between cattle and sheep in their digestion and relative intake of a mature tropical grass hay 3(1978)41
- Playne, M.J.
Estimation of the digestibility of low-quality hays by cattle from measurements made with sheep 3(1978)51
- Pollmann, D.S., Behnke, K.C. and Allee, G.L.
The effect of diet complexity and physical form on the performance of starter pigs and feed processing efficiency 8(1983)99
- Pollmann, D.S. and Bandyk, C.A.
Stability of viable lactobacillus products 11 (1984)261
- Pond, W.G., see Babatunde, G.M. 20(1988) 125
- Potter, S.G., Ammerman, C.B., Henry, P.R., Becker, H.N. and Palmer, A.Z.
Effect of sugarcane condensed molasses solubles, sugarcane molasses and monensin on performance and volatile fatty acid production in finishing steers 12(1984/1985)275
- Poutiainen, E., see Pulli, S. 1(1976)187
- Poyyamozi, V.S. and Kadirvel, R.
The value of banana stalk as a feed for goats 15(1986)95
- Pradhan, K., see Parthasarathy, M. 7(1982) 341
- Prasad, C.M., see Singh, K.S. 4(1979)143
- Prasad, C.M., see Singh, K.S. 6(1981)63
- Prasad, D.A., see Chandra, S. 12(1984/1985) 187
- Prasad, J., see Singh, A.P. 9(1983)325
- Prasad, T., see Lall, D. 23(1989)343
- Preston, R.L., see Rule, D.C. 15(1986)113
- Prieto, C., see Fonolla, J. 6(1981)405
- Prieto, C., see Aguilera, J.F. 12(1984/1985) 171
- Prieto, C., see Aguilera, J.F. 15(1986)33
- Prieto, F., see Szylit, O. 3(1978)73
- Pringle, D., Beames, R.M., Tait, R.M. and Litsky, J.
Effect of storage and processing of barley on its utilization by pigs and rats 9(1983)89
- Pringle, D.B., see Beames, R.M. 15(1986)285
- Proudfoot, F.G., see Zarkadas, C.G. 14(1986)

- 291
Proudfoot, F.G., see Hulan, H.W. 16(1986/1987)253
Proven, M.J., see McDonald, P. 8(1983)259
Pulli, S., Poutiainen, E. and Syrjala, L.
Prospects of growing maize in Finland 1 (1976)187
Pulungan, H., see Van Eys, J.E. 18(1987)197
Punj, M.L., see Seghal, J.P. 9(1983)155
Punj, M.L., see Jakhmola, R.C. 19(1988)1
Pusztai, A.J., see Williams, P.E.V. 12(1984/1985)1
Pyle, C., see Wright, D.E. 2(1977)93
- R**
Radosevich, S.R., see Sidahmed, A.E. 8(1983)47
Raharjo, Y. and Farrell, D.J.
A new biological method for determining amino acid digestibility in poultry feedstuffs using a simple cannula, and the influence of dietary fibre on endogenous amino acid output 12(1984/1985)29
Raj Reddy, M., see Krishna Reddy, G.V. 24 (1989)69
Rajaguru, A.S.B., see Ravindran, V. 17(1987)227
Rakes, J.M., see Daniels, L.B. 8(1983)19
Rana, V.K., see Langar, P.N. 13(1985)103
Rangkuti, M., see Van Eys, J.E. 18(1987)197
Rangnekar, D.V., Badve, V.C., Kharat, S.T., Sobale, B.N. and Joshi, A.L.
Effect of high-pressure steam treatment on chemical composition and digestibility in vitro of roughages 7(1982)61
Rao, N.M. and Fontenot, J.P.
Ensiling of rumen contents and blood with wheat straw 18(1987)67
Ravindran, V., Kornegay, E.T. and Rajaguru, A.S.B.
Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227
Razzaque, M.A., Al-Nasser, A., Salman, A.J., Aderibigbe, A.O. and Church, D.C.
Corrugated cardboard and dried poultry manure as dietary ingredients for sheep 14 (1986)265
Razzaque, M.A., see Ilian, M.A. 19(1988)327
Razzaque, M.A., see Ilian, M.A. 19(1988)343
Reade, A.E., see Gregory, K.F. 2(1977)7
Rebolé, A. and Alvira, P.
Composition of vine-branches with leaves of *Vitis vinifera* L. fresh and ensiled using different additives (previous results) 16(1986/1987)89
Rebolé, A., Alvira, P. and González, G.
Digestibility in vivo of ensiled grapevines (branches and leaves): influence of the system of analysis in the detergent fibre scheme on the prediction of digestibility 22(1988)173
Rechkemmer, G., see Von Engelhardt, W. 23 (1989)43
Reddy, G.V.N. and Reddy, M.R.
Effect of ammonia treatment and processing of whole cotton plants as sole source of roughage in complete feeds for growing cross-bred calves 13(1985)93
Reddy, G.V.N. and Reddy, M.R.
Effects of ammonia treatment and processing of cotton straw as the sole source of roughage in complete feeds on rumen characteristics with Murrah buffaloes 14(1986)95
Reddy, M.R., see Reddy, G.V.N. 13(1985)93
Reddy, M.R., see Reddy, G.V.N. 14(1986)95
Redman, R.G., see Kellaway, R.C. 3(1978)201
Reid, G.W., see Ørskov, E.R. 8(1983)247
Reid, G.W., see Istasse, L. 15(1986)167
Reid, R.L. and Horvath, D.J.
Soil chemistry and mineral problems in farm livestock. A review 5(1980)95
Reid, S.J., see Hyslop, J.J. 25(1989)55
Rekib, A., see Singh, A.P. 9(1983)325
Rexen, B.
Enzyme solubility - a method for evaluating the digestibility of alkali-treated straw 2 (1977)205
Rexen, B., see Israelsen, M. 3(1978)227
Rexen, B.
Use of enzymes for improvement of feed 6 (1981)105
Rexen, F. and Vestergaard Thomsen, K.
The effect on digestibility of a new technique for alkali treatment of straw 1(1976)73
Rey, J.Ma., see Ocio, E. 4(1979)227
Rezende, S.A., see Brown, D.L. 18(1987)247
Richardson, C.R., see Bunting, L.D. 22(1988)61
Riggs, J.K., see Schake, L.M. 7(1982)257
Rigsby, L.L., see Akin, D.E. 19(1988)261
Rigsby, L.L., see Akin, D.E. 23(1989)305
Rioni, M., see Lanari, D. 1(1976)401
Rioni, M. and Lanari, D.
Bull-beef production from the maize crop in Italy in large and small units 1(1976)531
Riopérez, J., see Tortuero, F. 25(1989)79
Riou, Y., see Gouet, P. 2(1977)297
Ritchie, A.H. and Mackie, I.M.
Preparation of fish protein hydrolysates 7 (1982)125
Robert, P., Bertrand, D. and Demarquilly, C.
Prediction of forage digestibility by principal component analysis of near infrared reflectance spectra 16(1986/1987)215
Roberts, D.J.
The substitution of grass silage by a straw and concentrate mixture for dairy cows 20 (1988)135
Robertson, S., see Huhtanen, P. 21(1988)11
Robinson, J.J., see Kowalczyk, J. 7(1982)225
Robinson, P.H., Fadel, J.G. and Tamminga, S.
Evaluation of mathematical models to de-

- scribe neutral detergent residue in terms of its susceptibility to degradation in the rumen 15(1986)249
- Robinson, P.H., Brice, R.E. and Morrison, I.M.
Effect of straw treatment and laboratory of incubation on degradation of neutral detergent residue in the rumen 18(1987)75
- Robinson, P.H., see Tamminga, S. 25(1989)89
- Rodriguez, J., see Alibes, X. 8(1983)63
- Rodriguez, J., see Alibes, X. 11(1984)189
- Rodriguez, M.L., see Tortuero, F. 25(1989)79
- Rollin, F., see Istasse, L. 25(1989)1
- Romita, A.
Survey of the maize situation in Italy 1 (1976)109
- Rönnau, K., see Von Engelhardt, W. 23(1989)43
- Roodbeen, A.E., see Papadopoulos, M.C. 14 (1986)279
- Rooke, J.A., Brett, P.A., Overend, M.A. and Armstrong, D.G.
The energetic efficiency of rumen microbial protein synthesis in cattle given silage-based diets 13(1985)255
- Rooke, J.A., Bell, S.L. and Armstrong, D.G.
The chemical composition of grass silages prepared with and without pre-treatment with inoculants containing *Lactobacillus plantarum* 13(1985)269
- Rooke, J.A., see Lee, N.-H. 15(1986)301
- Rooney, L.W., see Pflugfelder, R.L. 14(1986)243
- Rose, S.P. and Michie, W.
Meat and bone and fish meals in balancer feeds for choice-fed broilers 11(1984)221
- Ross-Parker, H.M., see Jayasuriya, M.C.N. 7 (1982)201
- Rossmagel, B.G., see Thacker, P.A. 19(1988)191
- Roth, F.X. and Kirchgessner, M.
Methanol-fermentation protein in veal calf nutrition 1(1976)33
- Rotter, B.A., Nesar, M., Guenter, W. and Marquardt, R.R.
Effect of enzyme supplementation on the nutritive value of hullless barley in chicken diets 24(1989)233
- Rubinstein, A., see Ben-Ghedalia, D. 15(1986)47
- Ruiz Jr., L.P., White, S.F. and Hove, E.L.
The alkaloid content of sweet lupin seed used in feeding trials on pigs and rats 2 (1977)59
- Rule, D.C., Preston, R.L., Koes, R.M. and McReynolds, W.E.
Feeding value of sprouted wheat (*Triticum aestivum*) for beef cattle finishing diets 15 (1986)113
- Rundgren, M.
Low-glucosinolate rapeseed products for pigs - a review 9(1983)239
- Rundgren, M.
Evaluation of triticale given to pigs, poultry and rats 19(1988)359
- Russell, J.R. and Buxton, D.R.
Storage of large round bales of hay harvested at different moisture concentrations and treated with sodium diacetate and/or covered with plastic 13(1985)69
- Russell, J.R.
Influence of harvest date on the nutritive value and ensiling characteristics of maize stover 14(1986)11
- Russell, J.R., see Adamu, A.M. 20(1988)241
- Russell, J.R., see Adamu, A.M. 22(1988)227
- Russo, V., Catalano, A., Mariani, P. and Del Monte, P.
Utilization of yeast grown on *n*-paraffins in the feeding of heavy pigs, from weaning to slaughter 1(1976)25

S

- Sadiq, M., see Slinger, S.J. 3(1978)251
- Saiady, M.Y., see Fannesbeck, P.V. 16(1986/1987)7
- Said, A.N., see Nangole, F.N. 9(1983)121
- Sakata, T., see Von Engelhardt, W. 23(1989)43
- Salcedo, T.S., see Gómez, G. 11(1984)57
- Salman, A.J., see Razzaque, M.A. 14(1986)265
- Salman, A.J., see Ilian, M.A. 19(1988)327
- Salman, A.J., see Ilian, M.A. 19(1988)343
- Salmon, R.E.
Effects of pelleting, added sodium bentonite and fat in a wheat-based diet on performance and carcass characteristics of small white turkeys 12(1984/1985)223
- Salvador, A., see González, G. 20(1988)177
- Salyers, A.A. and McCarthy, R.E.
Assessing the importance of host-derived polysaccharides as carbon sources for bacteria growing in the human colon 23(1989)109
- Samuel, V., see Holzer, Z. 12(1984/1985)253
- Samuel, V., see Levy, D. 16(1986/1987)63
- Sanderson, M.A., Meyer, D.W. and Casper, H.H.
Dicoumarol concentrations in sweetclover hay treated with preservatives and in spoiled hay of high- and low-coumarin cultivars of sweetclover 14(1986)221
- Santini, F.J., see Napoli, G.M. 25(1989)39
- Santos, J., Gómez, G. and Alexander, J.C.
Production of fungal protein from rasped fresh cassava roots using 200- and 3000-litre fermentors 8(1983)313
- Santos-Nunez, J., see Gregory, K.F. 2(1977)7
- Sanz, R., see Fonolla, J. 6(1981)405
- Sarwatt, S.V., Mussa, M.A. and Kategile, J.A.
The nutritive value of ensiled forages cut at three stages of growth 22(1988)237
- Sauer, W.C. and Thacker, P.A.
Apparent ileal and faecal digestibility of

- amino acids in barley-based diets supplemented with soya bean meal or canola meal for growing pigs 14(1986)183
- Sauvant, D., Bertrand, D. and Giger, S.
Variations and prevision of the in sacco dry matter digestion of concentrates and by-products 13(1985)7
- Sauveur, B. and Thapon, J.L.
Effects of dietary maize solubles and minerals on albumen quality of fresh and stored brown-shelled eggs 8(1983)177
- Savić, R.
The use of techniques of maize cultivation to maximise feed for beef production 1(1976)229
- Savory, R., see Wood, J.F. 9(1983)307
- Saxton, A.M., see Coelho, M. 20(1988)219
- Schaefer, D.M., Brotz, P.G., Arp, S.C. and Cook, D.K.
Inoculation of corn silage and high-moisture corn with lactic acid bacteria and its effects on the subsequent fermentations and on feedlot performance of beef steers 25(1989)23
- Schake, L.M., Ellis, W.C., Suarez, W.A. and Riggs, J.K.
Preservation of sorghum plant portions harvested, processed and ensiled at ten stages of maturity 7(1982)257
- Schake, L.M., see Pflugfelder, R.L. 14(1986)243
- Scheele, C.W., see Kan, C.A. 19(1988)97
- Schemm, R.L., see Hibberd, C.A. 15(1986)231
- Schenoni, P., see Giardini, A. 1(1976)313
- Schneeberger, H.
Survey of the use of maize for livestock feeding in Switzerland 1(1976)149
- Schöne, F., see Lüdke, H. 22(1988)33
- Schöne, F., Lüdke, H., Hennig, A. and Jahreis, G.
Copper and iodine in pig diets with high glucosinolate rapeseed meal. II. Influence of iodine supplements for rations with rapeseed meal untreated or treated with copper ions on performance and thyroid hormone status of growing pigs 22(1988)45
- Schultz, J.A. and Tolan, J.W.
Particle size distribution of a feed premix containing amprolium 3(1978)211
- Schulz, E. and Oslage, H.J.
Composition and nutritive value of single-cell protein (SCP) 1(1976)9
- Schwarz, F.J., see Kirchgessner, M. 6(1981)337
- Seghal, J.P. and Punj, M.L.
Utilization of alkali-treated and neutralized wheat straw-based rations for growing goat kids 9(1983)155
- Sekine, J., Morita, Z., Oura, R. and Asahida, Y.
Water-drinking frequency and quantity of water drunk by steers fed hay or fresh forage 25(1989)207
- Seligman, N.G., see Goldman, A. 18(1987)233
- Setälä, J. and Syrjälä-Qvist, L.
Degradation of crude protein and quality of undegradable protein in untreated or formaldehyde-treated rapeseed meal 12(1984/1985)19
- Settineri, D., see Pace, V. 11(1984)125
- Sharma, H.R., Ingalls, J.R. and McKirdy, J.A.
Feeding value of alkali-treated whole rye (*Secale cereale* L.) grain for lactating cows and its digestibility for sheep 10(1983/1984)77
- Sharma, H.R., White, B. and Ingalls, J.R.
Utilization of whole rape (Canola) seed and sunflower seeds as sources of energy and protein in calf starter diets 15(1986)101
- Sharma, H.R., see Mbugi, P.K. 24(1989)267
- Sharma, O.P., see Lall, D. 9(1983)29
- Sharp, R.N., see Noland, P.R. 2(1977)219
- Sharp, R.N., Noland, P.R. and Campbell, D.R.
Relationships between tristimulus colour values and digestibility of grain sorghum by swine 2(1977)327
- Shelby, T., see Slabbert, N. 21(1988)31
- Shen, H., Summers, J.D. and Leeson, S.
The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303
- Shen, H., Summers, J.D. and Leeson, S.
Influence of supplemental fat and steam pelleting on the response of broilers to diets varying in energy density 13(1985)57
- Shimada, A., see Alcántara, E. 23(1989)323
- Shimojo, M. and Goto, I.
Effect of sodium silicate on forage digestion with rumen fluid of goats or cellulase using culture solutions adjusted for pH 24(1989)173
- Shin, H.T., see Greenhalgh, J.F.D. 3(1978)289
- Shqueir, A.A., Brown, D.L. and Klasing, K.C.
Canavanine content and toxicity of sesbania leaf meal for growing chicks 25(1989)137
- Sibbald, I.R., Elliot, J.I. and Cote, M.
Comparison of bioavailable energy values of diets measured with poultry and swine 8(1983)191
- Sidahmed, A.E., Morris, J.G., Radosevich, S.R. and Koong, L.J.
Seasonal changes in composition and intake of chaparral by Spanish goats 8(1983)47
- Silanikove, N., Cohen, O., Levanon, D., Kipnis, T. and Kugenheim, Y.
Preservation and storage of green panic (*Panicum maximum*) as moist hay with urea 20(1988)87
- Silva, A.T. and Ørskov, E.R.
Fibre degradation in the rumens of animals receiving hay, untreated or ammonia-treated straw 19(1988)277
- Silva, A.T. and Ørskov, E.R.
The effect of five different supplements on the degradation of straw in sheep given untreated barley straw 19(1988)289
- Silverside, D.E., see Machin, D.H. 15(1986)273

- Singh, A.P. and Pandit, N.N.
Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299
- Singh, A.P. and Pandita, N.N.
Studies on fermentation of sorghum silage during storage, and its effect on milch animals 9(1983)143
- Singh, A.P., Prasad, J. and Rekib, A.
Note on high-nitrate para grass (*Brachiaria mutica*) silage given to rabbits 9(1983)325
- Singh, B., see Negi, S.S. 22(1988)79
- Singh, K., see Jakhmola, R.C. 19(1988)1
- Singh, K.S., see Deolankar, R.P. 4(1979)133
- Singh, K.S. and Prasad, C.M.
Feeding value of sunflower and groundnut cakes for broilers 4(1979)143
- Singh, K.S., Prasad, C.M. and Brahmakshatriya, R.D.
Feeding value of sunflower- and groundnut-cakes for laying hens 6(1981)63
- Singh, R., see Kamra, D.N. 9(1983)185
- Singh, R., Kamra, D.N. and Jakhmola, R.C.
Ensilaging of leguminous green forages in combination with different dry roughages and molasses 12(1984/1985)133
- Singh, R., see Jakhmola, R.C. 19(1988)1
- Siroka, P., see Baran, M. 15(1986)7
- Siviter, J.W., see Smith, T. 10(1983/1984)189
- Siviter, J.W., see Smith, T. 14(1986)29
- Skoch, E.R., Behnke, K.C., Deyoe, C.W. and Binder, S.F.
The effect of steam-conditioning rate on the pelleting process 6(1981)83
- Skraba, B., see Hanczakowski, P. 6(1981)189
- Skraba, B., see Hanczakowski, P. 6(1981)413
- Skraba, B., see Hanczakowski, P. 12(1984/1985)11
- Skraba, B., see Hanczakowski, P. 24(1989)151
- Skrede, A. and Nes, I.F.
Slaughterhouse by-products preserved by *Lactobacillus plantarum* fermentation as feed for mink and foxes 20(1988)287
- Slabbert, N., Bolt, M.J., Shelby, T. and Campher, J.P.
The utilization by beef steers of a soft high lysine and a hard normal size maize cultivar in whole grain feedlot diets 21(1988)31
- Slinger, S.J., Leeson, S., Summers, J.D. and Sadiq, M.
Influence of steam pelleting on the feeding value of tower and candle rapeseed products for broiler chickens 3(1978)251
- Smart, M.E., see Mann, M.E. 21(1988)57
- Smart, R.L., see Mbatya, P.B.A. 8(1983)221
- Smart, R.L., see Mbatya, P.B.A. 8(1983)229
- Smart, R.L., see Mbatya, P.B.A. 13(1985)281
- Smart, R.L., see Mbatya, P.B.A. 13(1985)293
- Smart, R.L., see Istasse, L. 16(1986/1987)37
- Smith, M.A., Watson Jr., C.E., Watson, V.H. and Trevathan, L.E.
Effects of *Drechslera sorokiniana* infection on the yield and quality of tall fescue forage 15(1986)41
- Smith, M.J., see Daniels, L.B. 8(1983)19
- Smith, O.B. and Macleod, G.K.
Digestibility of wet cage layer excreta and an excreta-supplemented diet by cattle as determined by two methods 5(1980)77
- Smith, O.B., Ilori, J.O. and Onesirosoan, P.
The proximate composition and nutritive value of the winged bean *Psophocarpus tetragonolobus* (L.) DC for broilers 11(1984)231
- Smith, O.B. and Adegbola, A.A.
Studies on the feeding value of agro-industrial by-products. III. Digestibility of cocoa-pod and cocoa-pod-based diets by ruminants 13(1985)249
- Smith, O.B., see Kabajja, E. 20(1988)171
- Smith, O.B., Osafo, E.L.K. and Adegbola, A.A.
Studies on the feeding value of agro-industrial by-products: strategies for improving the utilisation of cocoa-pod-based diets by ruminants 20(1988)189
- Smith, R.E., see Gregory, K.F. 2(1977)7
- Smith, T., Grigera-Naón, J.J., Broster, W.H. and Siviter, J.W.
Ammonia versus sodium hydroxide treatment of straw for growing cattle 10(1983/1984)189
- Smith, T., Grigera-Naón, J.J., Broster, W.H. and Siviter, J.W.
The use of barley straw as a feed 14(1986)29
- Smith, W.C., see Kies, A.K. 16(1986/1987)169
- Smith, W.C., Moughan, P.J., Pearson, G. and James, K.A.C.
Comparative bioavailable energy values of five ground cereal grains measured with growing rats and pigs 18(1987)143
- Smith, W.C., Moughan, P.J., Pearson, G. and Patchell, M.R.
A comparison of bioavailable energy values of ground cereal grains measured with adult cockerels and growing pigs 19(1988)105
- Smith, W.C., see Moughan, P.J. 22(1988)203
- Smits, C.H.M., see Moughan, P.J. 22(1988)203
- Sobale, B.N., see Rangnekar, D.V. 7(1982)61
- Søbstad, G., see Opstvedt, J. 18(1987)181
- Soliman, H., see Mohamed, K. 11(1984)199
- Soliman, H.S., see Ørskov, E.R. 7(1982)135
- Solomon, R., see Ben-Ghedalia, D. 18(1987)55
- Solomon, R., see Ben-Ghedalia, D. 22(1988)147
- Sonaiya, E.B. and Omole, T.A.
Cassava meal and cassava peel meal in diets for growing pigs 8(1983)211
- Sosulski, F.W., see Campbell, G.L. 16(1986/1987)243
- Spears, J.W., see Guardiola, C.M. 8(1983)129
- Spears, J.W., see Harvey, R.W. 24(1989)337
- Spolstra, S.F., see Courtin, M.G. 24(1989)97
- Spooner, A.E., see Higgins, T.R. 16(1986/1987)1
- Spooner, M.C., see Osbourn, D.F. 6(1981)387

- Spragg, J.C., see Amaning-Kwarteng, K. 16 (1986/1987)75
- Spurr, D.T., see Kernan, J.A. 6(1981)257
- Spurr, D.T., see Kernan, J.A. 11(1984)301
- Srećković, A.
Review of production and utilisation of maize for animal nutrition in Yugoslavia 1 (1976)115
- Srivastava, R.V.N., see Kamra, D.N. 9(1983)185
- Stallcup, O.T., see Daniels, L.B. 8(1983)19
- Stanogias, G. and Pearce, G.R.
Digestibility by cattle of diets containing dried pig faeces 3(1978)155
- Staples, C.R. and Lough, D.S.
Efficacy of supplemental dietary neutralizing agents for lactating dairy cows. A review 23(1989)277
- Stappers, H.P., see Grimbergen, A.H.M. 13 (1985)117
- Steg, A., see Deinum, B. 10(1983/1984)301
- Steg, A. and van der Meer, J.M.
Differences in chemical composition and digestibility of beet and cane molasses 13 (1985)83
- Stehr, W. and Kirchgessner, M.
The relationship between the intake of herbage grazed by dairy cows and its digestibility 1(1976)53
- Stephen, A.M.
The physiological effects of cellulose in the human large intestine 23(1989)241
- Stern, M.D., see Hannah, S.M. 16(1986/1987)51
- Stern, M.D., see Waltz, D.M. 25(1989)111
- Stewart, C.S., see Greenhalgh, J.F.D. 3(1978)289
- Stewart, C.S., see Theander, O. 10(1983/1984)89
- Stewart, C.S., see Cheng, K.-J. 10(1983/1984)93
- Stewart, C.S., see Graham, H. 12(1984/1985)195
- Stewart, C.S., Fonty, G. and Gouet, Ph.
The establishment of rumen microbial communities 21(1988)69
- Stobbs, the late T.H., see Aii, T. 5(1980)183
- Streeter, C.L. and Horn, G.W.
Effect of treatment of wheat straw with ammonia and peracetic acid on digestibility in vitro and cell wall composition 7(1982)325
- Streeter, C.L., see Horn, G.W. 8(1983)35
- Suarez, W.A., see Schake, L.M. 7(1982)257
- Subramanian, P.R., Kadirvel, R., Viswanathan, K. and Chandrasekaran, D.
In vitro studies and short-term feeding trial in lambs to evaluate plantain sheath (*Musa sapientum*) as a feed for ruminants 20(1988)343
- Sudana, I.B. and Leng, R.A.
Effects of supplementing a wheat straw diet with urea or a urea-molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25
- Suliman, A.H., see Osman, A.G. 19(1988)351
- Summers, J.D. and Leeson, S.
Utilization of processed pig hair in poultry diets 3(1978)243
- Summers, J.D., see Slinger, S.J. 3(1978)251
- Summers, J.D., Macleod, G.K. and Warner, W.C.
Chemical composition of culinary wastes and their potential as a feed for ruminants 5(1980)205
- Summers, J.D., see Shen, H. 8(1983)303
- Summers, J.D., see Leeson, S. 11(1984)211
- Summers, J.D., see Shen, H. 13(1985)57
- Summers, J.D. and Leeson, S.
Influence of nutrient density on feed consumption, weight gain and gut capacity of broilers, Leghorns and turkeys reared to 26 days of age 16(1986/1987)129
- Summers, J.D., see Leeson, S. 19(1988)313
- Sundstøl, F., see Kategile, J.A. 6(1981)133
- Sundstøl, F., see Borhami, B.E.A. 7(1982)45
- Sundstøl, F., see Borhami, B.E.A. 7(1982)53
- Sundstøl, F.
Ammonia treatment of straw: methods for treatment and feeding experience in Norway 10(1983/1984)173
- Sundstøl, F., see Wanapat, M. 12(1984/1985)295
- Sundstøl, F., see Dias-da-Silva, A.A. 14(1986)67
- Sundstøl, F., see Wanapat, M. 14(1986)215
- Suttle, N.F.
Reducing the potential copper toxicity of concentrates to sheep by the use of molybdenum and sulphur supplements 2(1977)235
- Swan, H., see Karalazos, A. 2(1977)143
- Swatson, H., see Atuahene, C.C. 17(1987)157
- Syrjala, L., see Pulli, S. 1(1976)187
- Syrjala-Qvist, L., see Setälä, J. 12(1984/1985)19
- Szylił, O., Durand, M., Borgida, L.P., Atinkpahoun, H., Prieto, F. and Delort-Laval, J.
Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73

T

- Tait, C.A.G., see Ørskov, E.R. 8(1983)247
- Tait, C.A.G., see Istasse, L. 15(1986)167
- Tait, R.M., see Pringle, D. 9(1983)89
- Tait, R.M., see Beames, R.M. 15(1986)285
- Tamminga, S., see Robinson, P.H. 15(1986)249
- Tamminga, S. and van Vuuren, A.M.
Formation and utilization of end products of ligno cellulose degradation in ruminants 21(1988)141
- Tamminga, S., see Ibrahim, M.N.M. 24(1989)83
- Tamminga, S., Robinson, P.H., Vogt, M. and

- Boer, H.
Rumen ingesta kinetics of cell wall components in dairy cows 25(1989)89
- Tangprasertchai, P., see McElhiney, R.R. 8 (1983)139
- Tangtaweewipat, S. and Elliott, R.
Nutritional value of pigeonpea (*Cajanus cajan*) meal in poultry diets 25(1989)123
- Tatterson, I.N.
Fish silage - preparation, properties and uses 7(1982)153
- Taylor, A. G., see Morgan, C.A. 24(1989)219
- Taylor, A.G., see Peers, D.G. 2(1977)41
- Teh, T.H., Hemken, R.W., Bremel, D.H. and Harmon, R.J.
Comparison of buffers on rumen functions, turnover rate and gastric secretions in Holstein steers 17(1987)257
- Telfer, S.B., see Zervas, G. 21(1988)23
- Teller, E. and Godeau, J.-M.
The effect of nitrogen supplements to maize silage on the amount and composition of the nitrogen fraction entering the intestine of cattle 13(1985)25
- Terry, R.A., see Osbourn, D.F. 6(1981)387
- Tetlow, R.M., see Osbourn, D.F. 6(1981)387
- Tetlow, R.M., see Bolsen, K.K. 9(1983)37
- Tetlow, R.M.
The effect of urea on the preservation and digestibility in vitro of perennial ryegrass 10 (1983/1984)49
- Tetlow, R.M., see Woolford, M.K. 11(1984)159
- Tetlow, R.M., Wilson, R.F. and Bolsen, K.K.
Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91
- Tetlow, R.M. and Mason, V.C.
Treatment of whole-crop cereals with alkali.
1. The influence of sodium hydroxide and ensiling on the chemical composition and in vitro digestibility of rye, barley and wheat crops harvested at increasing maturity and dry matter content 18(1987)257
- Tetlow, R.M., Mason, V.C. and Deschard, G.
Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271
- Tetlow, R.M., see Deschard, G. 18(1987)283
- Tetlow, R.M., see Deschard, G. 19(1988)55
- Tewe, O.O.
Effect of cassava-based diets varying in cyanide content on the performance and physiopathology of the African giant rat (*Cricetomys gambianus* Waterhouse) 11 (1984)1
- Thacker, P.A., see Bowland, J.P. 12(1984/1985)75
- Thacker, P.A., see Sauer, W.C. 14(1986)183
- Thacker, P.A., Bell, J.M., Classen, H.L., Campbell, G.L. and Rossmagel, B.G.
The nutritive value of hullless barley for swine 19(1988)191
- Tham, S.H., see Karunajeewa, H. 17(1987)271
- Thanu, K., Kadirvel, R. and Ayyaluswami, P.
The effect of nutrient supplementation on the feeding value of kapok seed for poultry 9(1983)263
- Thapon, J.L., see Sauveur, B. 8(1983)177
- Theander, O., Hartley, R.D. and Stewart, C.S.
A note on the OECD collaborative experiment on ammonia-treated straw 10(1983/1984)89
- Theander, O., see Graham, H. 12(1984/1985)195
- Theander, O., see Lindberg, J.E. 15(1986)197
- Theander, O.
Possibilities for multipurpose fractionation and alternative uses of plant biomass 21 (1988)267
- Theander, O., Westerlund, E., Åman, P. and Graham, H.
Plant cell walls and monogastric diets 23 (1989)205
- Theodorou, M.K., see Akin, D.E. 19(1988)261
- Theodorou, M.K., see Gascoyne, D.J. 21 (1988)183
- Thiago, L.R.L., see Kellaway, R.C. 3(1978)201
- Thiago, L.R.L. de S. and Kellaway, R.C.
Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71
- Thiesen, J.
Detoxification of aflatoxins in groundnut meal 2(1977)67
- Thivend, P., see Jouany, J.P. 15(1986)215
- Thomas, H.R., see Harper, A.F. 8(1983)69
- Thomas, V.M., Katz, R.J., Auld, D.L. and Peterson, C.L.
Value of mechanically extracted rape and safflower oilseed meals as protein supplements for growing lambs 11(1984)269
- Thomke, S., see Muindi, P.J. 6(1981)197
- Thomke, S., see Hesselman, K. 7(1982)351
- Thompson, C. and Moran, J.B.
The rate of dry matter disappearance from nylon bags of maize silage/grain diets as influenced by basal ration and processing of test samples 16(1986/1987)225
- Thompson, L.U., see McBurney, M.I. 23(1989)261
- Thomson, E., see Farrell, D.J. 9(1983)99
- Timbers, G.E., see Elliot, J.I. 17(1987)213
- Tjerneld, F., see Hahn-Hägerdal, B. 21(1988)175
- Tock, R.W., see Bunting, L.D. 22(1988)61
- Tolan, J.W., see Schultz, J.A. 3(1978)211
- Topps, J.H., see El-Sherbiny, A.E. 5(1980)33
- Topps, J.H., see El-Sherbiny, A.E. 5(1980)43
- Tortuero, F., Riopérez, J. and Rodríguez, M.L.
Nutritional value for rabbits of olive pulp and the effects on their visceral organs 25 (1989)79
- Touchburn, S.P., see Chavez, E.R. 22(1988)1
- Touchburn, S.P., see Chavez, E.R. 22(1988)13

- Touchburn, S.P., see Chavez, E.R. 22(1988)23
 Treacher, T.T., see Penning, P.D. 5(1980)321
 Trenholm, H.L., see Hamilton, R.M.G. 11 (1984)293
 Trenkle, A., see Adamu, A.M. 20(1988)241
 Trenkle, A., see Adamu, A.M. 22(1988)227
 Trevathan, L.E., see Smith, M.A. 15(1986)41
 Treviño, J., Centeno, C. and Caballero, R.
 The chemical composition of pea plant parts as related to harvesting time 16(1986/1987) 305
 Trigg, T.E., see Moran, J.B. 20(1988)299
 Tuite, J., see Cherney, J.H. 17(1987)45
 Tullis, J.B. and Whittemore, C.T.
 Digestibility of a fully hydrogenated tallow for growing pigs 5(1980)87
 Twumasi, I.K., see Osei, S.A. 24(1989)247
- U**
- Udedibie, A.B.I. and Igwe, F.O.
 Dry matter yield and chemical composition of pigeon pea (*C. cajan*) leaf meal and the nutritive value of pigeon pea leaf meal and grain meal for laying hens 24(1989)111
 Uden, P. and Van Soest, P.J.
 The determination of digesta particle size in some herbivores 7(1982)35
 Uden, P.
 The effect of intake and hay: concentrate ratio upon digestibility and digesta passage 11(1984)167
 Uden, P.
 Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279
 Uden, P.
 The effect of grinding and pelleting hay on digestibility, fermentation rate, digesta passage and rumen and faecal particle size in cows 19(1988)145
 Umunna, N.N., Klopfenstein, T.J., Hasimoglu, S. and Woods, W.R.
 Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375
 Umunna, N.N., Bartling, R.R. and Klopfenstein, T.J.
 High temperature and pressure processing of maize cobs. I. Digestibility in vitro of processed cobs 12(1984/1985)151
 Umunna, N.N., Bartling, R.R. and Klopfenstein, T.J.
 High temperature and pressure processing of maize cobs. II. Animal response and effect of sodium metabisulphite 14(1986)161
 Urio, N.A., see Kategile, J.A. 6(1981)133
- Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989) 45
 Vaid, J., see Lall, D. 9(1983)29
 Valdivie, M., Compte, X. and Fundora, O.
 The utilization of torula yeast in diets for White Leghorn birds during growth and laying periods 7(1982)185
 Valdivieso, M., see Gómez, G. 11(1984)49
 Valdivieso, M., see Gómez, G. 11(1984)57
 Valentine, S.C. and Bartsch, B.D.
 Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammermilled lupin grain in oaten hay or oaten pasture based diets 16(1986/1987)143
 Valentine, S.C. and Bartsch, B.D.
 Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261
 van der Aar, P.J., see Mora, M.I. 9(1983)205
 van der Honing, Y., see Van Es, A.J.S. 1(1976) 481
 van der Meer, J.M., see Steg, A. 13(1985)83
 Van Eenaeme, C., see Istasse, L. 25(1989)1
 van Es, A.J.S., van der Honing, Y. and Boekholt, H.A.
 The nutritive value (energy) of maize silage 1(1976)481
 Van Eys, J.E., Pulungan, H., Rangkuti, M. and Johnson, W.L.
 Cassava meal as supplement to napier grass diets for growing sheep and goats 18(1987) 197
 van Houtert, M.F.J., see Jayasuriya, M.C.N. 7(1982)201
 Van Nevel, C.J., see Kayouli, C. 10(1983/1984)165
 Van Soest, P.J., see Uden, P. 7(1982)35
 Van Soest, P.J., Mascarenhas Ferreira, A. and Hartley, R.D.
 Chemical properties of fibre in relation to nutritive quality of ammonia-treated forages 10(1983/1984)155
 van Soest, P.J., see Lau, M.M. 6(1981)123
 van Vuuren, A.M., see Tamminga, S. 21(1988) 141
 Van Weerden, E.J. and Huisman, J.
 Digestibility of protein and amino acids of a fermentation single-cell protein for veal calves 2(1977)377
 Vanacker, J.M., see Cottyn, B.G. 11(1984)137
 Vanacker, J.M., see De Boever, J.L. 14(1986) 203
 Vanacker, J.M., see De Boever, J.L. 19(1988) 247
 Vanbelle, M., see Nefzaoui, A. 14(1986)139
 Vandevoorde, L., Bruyneel, B. and Verstraete, W.
 In-reactor cellulose conversion 21(1988)99
 Vecchiattini, M., see Giardini, A. 1(1976)369
 Vecchiattini, M., see Giardini, A. 1(1976)313
 Vedder, H.M., see Barton II, F.E. 15(1986)189
 Veira, D.M., see Ivan, M. 9(1983)131
 Vencl, B. and Flám, F.

V

- Vadiveloo, J.
 The intake and digestibility in goats of

- Predicting the energetic value of forage from digestibility of organic matter in vitro or solubility in potassium hydroxide 6(1981) 235
- Verstraete, W., see Vandevoorde, L. 21(1988) 99
- Vervaeke, I.J., see Dierick, N.A. 23(1989)141
- Vervaeke, I.J., Dierick, N.A., Demeyer, D.I. and Decuyper, J.A.
Approach to the energetic importance of fibre digestion in pigs. II. An experimental approach to hindgut digestion 23(1989)169
- Vestergaard Thomsen, K., see Rexen, F. 1 (1976)73
- Vestergaard Thomsen, K., see Waagepetersen, J. 2(1977)131
- Vestergaard Thomsen, K., see El Khidir, O.A. 7(1982)277
- Vestergaard Thomsen, K., see El Khidir, O.A. 9(1983)197
- Vignon, B., see Laurent, F. 13(1985)131
- Viñaras, R., see Ocio, E. 4(1979)227
- Viswanathan, K., see Subramanian, P.R. 20 (1988)343
- Viswanathan, K., see Chandrasekaran, D. 22 (1988)321
- Viswanathan, K., Kadirvel, R. and Chandrasekaran, D.
Nutritive value of banana stalk (*Musa cavendishii*) as a feed for sheep 22(1988)327
- Vlach, D., see Moo-Young, M. 5(1980)175
- Vogt, M., see Tamminga, S. 25(1989)89
- Volenc, J.J., see Cherney, J.H. 17(1987)45
- Von Engelhardt, W., Rönau, K., Reckemmer, G. and Sakata, T.
Absorption of short-chain fatty acids and their role in the hindgut of monogastric animals 23(1989)43
- ## W
- Waagepetersen, J. and Vestergaard Thomsen, K.
Effect on digestibility and nitrogen content of barley straw of different ammonia treatments 2(1977)131
- Wagner, D.G., see Hibberd, C.A. 15(1986)231
- Wainman, F.W., see De Boever, J.L. 14(1986) 203
- Waldroup, P.W., Bussell, W.D., Burke, A.B. and Johnson, Z.B.
Comparison of the relative biological availability and stability of commercial sources of vitamin K activity 3(1978)329
- Waldroup, P.W., Whelchel, D.L. and Johnson, Z.B.
Variation in nutrient content of samples of dried bakery product 7(1982)419
- Walker, H.G., Mackay, B.E. and Montague Jr., W.C.
Composition and enzymatic digestibility of oregon grass straws 9(1983)283
- Walker, N.
The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21
- Walker, N.
Treatment of the haem fraction of abattoir blood with formic acid to produce a solid state, low dry matter protein supplement for use in finishing-pig diets 2(1977)367
- Walker, N., see Patterson, D.C. 4(1979)263
- Walker, N., see Patterson, D.C. 4(1979)275
- Walker, N.
A comparison of wheat- or barley-based diets given ad libitum as meal or pellets to finishing pigs 22(1988)263
- Walker, N.
Observations on the feeding of non-lactating sows on graded levels of liquid whey 24 (1989)327
- Walli, T.K., see Bhargava, P.K. 22(1988)295
- Walter, K.
Economic aspects of high-plane feeding of bulls with maize silage 1(1976)513
- Waltz, D.M. and Stern, M.D.
Evaluation of various methods for protecting soya bean protein from degradation by rumen bacteria 25(1989)111
- Wanapat, M., Sundstøl, F. and Garmo, T.H.
A comparison of alkali treatment methods to improve the nutritive value of straw. I. Digestibility and metabolizability 12(1984/1985)295
- Wanapat, M., Sundstøl, F. and Hall, J.M.R.
A comparison of alkali treatment methods used to improve the nutritive value of straw. II. In sacco and in vitro degradation relative to in vivo digestibility 14(1986)215
- Ward, J.K., see Faulkner, D.B. 12(1984/1985) 125
- Warner, W.C., see Summers, J.D. 5(1980)205
- Waterworth, D.G. and Heath, M.E.
Pruteen in the diet of breeding pigs: reproductive performance 6(1981)297
- Watson, C., see Batterham, E.S. 13(1985)171
- Watson, V.H., see Smith, M.A. 15(1986)41
- Watson Jr., C.E., see Smith, M.A. 15(1986)41
- Weibel, D.E., see Hibberd, C.A. 15(1986)231
- Weigand, E. and Kirchgessner, M.
Protein and energy value of vinnase for pigs 5(1980)221
- Weinberg, Z.G., Pahlow, G., Dinter, B. and Ashbell, G.
The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20(1988)335
- Weir, J., see Hyslop, J.J. 25(1989)55
- Weller, R.F., see Phipps, R.H. 1(1976)251
- Werner, D., see Arieli, A. 23(1989)333
- Westerlund, E., see Theander, O. 23(1989)205
- Westlake, K., Mackie, R.I. and Dutton, M.F.
In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations 25(1989)169
- Whelchel, D.L., see Waldroup, P.W. 7(1982) 419
- White, B., see Sharma, H.R. 15(1986)101

- White, S.F., see Ruiz Jr., L.P. 2(1977)59
- Whitehead, D.C., Goulden, K.M. and Hartley, R.D.
Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231
- Whittemore, C.T., see Peers, D.G. 2(1977)41
- Whittemore, C.T.
The potato (*Solanum tuberosum*) as a source of nutrients for pigs, calves and fowl - a review 2(1977)171
- Whittemore, C.T., see Tullis, J.B. 5(1980)87
- Whittemore, C.T., see Morgan, C.A. 7(1982)387
- Whittemore, C.T., see Morgan, C.A. 11(1984)11
- Whittemore, C.T., see Morgan, C.A. 17(1987)81
- Whittemore, C.T., see Morgan, C.A. 19(1988)185
- Whittemore, C.T., see Morgan, C.A. 24(1989)219
- Wignjoesastro, N. and Young, A.W.
Digestibility of diets containing increasing levels of NaOH-treated or untreated wheat straw 7(1982)331
- Wijayatunge, C., see Jayasuriya, M.C.N. 7(1982)221
- Wilcock, J.M., see Hyslop, J.J. 25(1989)55
- Wilkins, R.J.
Discussion on session 2 1(1976)273
- Wilkinson, J.M.
Discussion on session I 1(1976)215
- Wilkinson, J.M.
Voluntary intake and efficiency of utilisation of whole-crop maize silage 1(1976)441
- Wilkinson, J.M. and Gonzalez Santillana, R.
Ensiled alkali-treated straw. I. Effect of level and type of alkali on the composition and digestibility in vitro of ensiled barley straw 3(1978)117
- Wilkinson, J.M. and Gonzalez Santillana, R.
Ensiled alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensiled or frozen alkali-treated straw and ryegrass silage 3(1978)133
- Wilkinson, J.M.
Laboratory evaluation of ensiled mixtures of alkali-treated cattle excreta and whole-crop maize 3(1978)335
- Wilkinson, J.M., see Gonzalez Santillana, R. 3(1978)381
- Wilkinson, J.M., see Woolford, M.K. 8(1983)107
- Williams, A.G.
Factors affecting the formation of polysaccharide-degrading enzymes by rumen micro-organisms 21(1988)191
- Williams, J.E., see Goedeken, F.K. 13(1985)47
- Williams, K.C., see Neill, A.R. 9(1983)71
- Williams, P.E.V., see Ørskov, E.R. 6(1981)355
- Williams, P.E.V.
Digestibility studies on ammonia-treated straw 10(1983/1984)213
- Williams, P.E.V., Macdearmid, A., Innes, G.M. and Gauld, S.A.
Ammonia-treated barley straw and rolled barley offered either together, in a mixed ration, or successively to beef steers 10(1983/1984)247
- Williams, P.E.V., Innes, G.M. and Brewer, A.
Ammonia treatment of straw via the hydrolysis of urea. I. Effects of dry matter and urea concentrations on the rate of hydrolysis of urea 11(1984)103
- Williams, P.E.V., Innes, G.M. and Brewer, A.
Ammonia treatment of straw via the hydrolysis of urea. II. Additions of soya bean (urease), sodium hydroxide and molasses; effects on the digestibility of urea-treated straw 11(1984)115
- Williams, P.E.V., Pusztai, A.J., Macdearmid, A. and Innes, G.M.
The use of kidney beans (*Phaseolus vulgaris*) as protein supplements in diets for young rapidly growing beef steers 12(1984/1985)1
- Williams, P.E.V., see Fallon, R.J. 14(1986)103
- Wilson, J.R., see McLeod, M.N. 24(1989)185
- Wilson, M.N., see Moughan, P.J. 22(1988)203
- Wilson, P.N., Brigstocke, T.D.A. and Cuthbert, N.H.
Some factors affecting the future composition of UK compound animal feeds 6(1981)1
- Wilson, R.F., see Bolsen, K.K. 9(1983)37
- Wilson, R.F., see Tetlow, R.M. 12(1984/1985)91
- Wiseman, J.
A note on the nutritive value of dried instant coffee residue for broiler chickens and turkey poults 10(1983/1984)285
- Wiseman, J., see Green, S. 21(1988)43
- Wohlt, J.E., Fiallo, J.F. and Miller, M.E.
Composition of by-products of the essential-oil industry and their potential as feeds for ruminants 6(1981)115
- Wolf, I., see Flachowsky, G. 17(1987)235
- Wolfram, D., see Flachowsky, G. 17(1987)235
- Wolsink, J.H., see Barton II, F.E. 15(1986)189
- Wood, J.F., Carter, P.M. and Savory, R.
Investigations into the effects of processing on the retention of the carotenoid fractions of *Leucaena leucocephala* during storage, and the effects of processing on mimosine concentration 9(1983)307
- Wood, J.F.
The functional properties of feed raw materials and their effect on the production and quality of feed pellets 18(1987)1
- Woods, W.R., see Umunna, N.N. 7(1982)375
- Woolford, M.K. and Cook, J.E.
A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3(1978)89

- Woolford, M.K., Wilkinson, J.M. and Cook, J.E.
Investigations on the effect of sodium bentonite and other moisture absorbents on the production of effluent from grass silage 8 (1983)107
- Woolford, M.K. and Tetlow, R.M.
The effect of anhydrous ammonia and moisture content on the preservation and chemical composition of perennial ryegrass hay 11(1984)159
- Wrathall, J.H.M., Owen, E. and Pike, D.J.
Upgrading barley straw for goats: the effectiveness of a sodium hydroxide and urea dip method 24(1989)57
- Wright, D.E., Payne, E., Pyle, C., Aitken, W.M. and Kirton, A.H.
Poly-unsaturated fat in young ruminants 2 (1977)93
- Wyers, M., see Cherbut, C. 20(1988)327
- Wyss, U. and Bickel, H.
Ripe beans of *Canavalia ensiformis* (jack-bean) as feed ingredient for monogastric animals 20(1988)325
- Yusef, E., see Ben-Ghedalia, D. 19(1988)219
- Yusef, E., see Ben-Ghedalia, D. 22(1988)247
- Yusef, E., see Ben-Ghedalia, D. 22(1988)255
- Yusef, E., see Ben-Ghedalia, D. 24(1989)289
- Young, A.W., see Wignjosoesastro, N. 7(1982)331
- Yulzari, A., see Goldman, A. 18(1987)233
- F.G.
The amino acid and mineral composition of white fish meal containing enzyme-digested or untreated stickwater solids 14(1986)291
- Zarkadas, C.G., see Hulan, H.W. 16(1986/1987)253
- Zemmelink, G., see Ibrahim, M.N.M. 24 (1989)83
- Zervas, G.
Treatment of dairy sheep with soluble glass boluses containing copper, cobalt and selenium 19(1988)79
- Zervas, G., Telfer, S.B., Carlos, G. and Anderson, P.
The effect of soluble-glass boluses containing copper, cobalt and selenium on the blood composition of ewes 21(1988)23
- Zimmer, E. and Zscheischler, J.
Use of maize for livestock feeding in the Federal Republic of Germany 1(1976)157
- Zimmer, E.
Efficiency of harvesting and conservation 1 (1976)289
- Zimmer, E.
Discussion on session 3 1(1976)327
- Ziras, E., see Papadopoulos, G. 18(1987)295
- Ziras, E., see Papadopoulos, G. 18(1987)303
- Zorrilla-Rios, J., see Horn, G.W. 24(1989)201
- Zorrilla-Rios, J., Horn, G.W. and McNew, R.W.
Effect of ammoniation and energy supplementation on the utilization of wheat straw by sheep 22(1988)305
- Zscheischler, J., see Zimmer, E. 1(1976)157

Y

Z

- Zacchi, G., see Hahn-Hägerdal, B. 21(1988)175
- Zarkadas, C.G., Hulan, H.W. and Proudfoot,

SUBJECT INDEX

Animal Feed Science and Technology, volumes 1-25

A

Acid treatment

Studies on ammonia-treated straw. II. Fixation of ammonia in treated straw by spraying with acids 7(1982)53

Accuracy of predicting digestibility by the cellulase technique; the effect of pretreatment of forage samples with neutral detergent or acid pepsin 7(1982)83

Aflatoxins

Detoxification of aflatoxins in groundnut meal 2(1977)67

In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations 25(1989)169

African giant rat

Effect of cassava-based diets varying in cyanide content on the performance and physiopathology of the African giant rat (*Cricetomys gambianus* Waterhouse) 11(1984)1

African yam

Variation in metabolizable energy content of raw or autoclaved white and brown varieties of three tropical grain legumes 13(1985)141

Agronomic factors

Cultivation of maize in irrigated areas for silage and grain production and their use by ruminants 1(1976)237

Effect of agronomic factors on the yield of maize and on its composition in relation to its use by ruminants 1(1976)251

Agronomic factors affecting the growth and composition of the maize plant 1(1976)263

Soil chemistry and mineral problems in farm livestock. A review 5(1980)95

The quality of the green fraction of leaf protein concentrate from *Chenopodium quinoa* Willd. grown at different levels of fertilizer nitrogen 11(1984)239

The nutritional value of crop residue components from several wheat cultivars grown at different fertilizer levels 11(1984)301

Estimating yield and nutrient losses due to rainfall on field-drying alfalfa hay 16(1986/1987)7

Agropyron elongatum

Effect of lactate on the in vitro digestion of *Agropyron elongatum* by rumen microorganisms 16(1986/1987)161

Albumen quality

Effects of dietary maize solubles and minerals on albumen quality of fresh and stored brown-shelled eggs 8(1983)177

Alcaligenes eutrophus

Nitrogen metabolism in broiler chickens consuming the bacterial strain *Alcaligenes eutrophus* 5(1980)241

Alfalfa

Microwave drying of alfalfa compared to field- and oven-drying: effects on forage quality 16(1986/1987)1

Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and stored at different moisture contents 17(1987)45

Composition and fibre digestion in morphological components of an alfalfa-timothy sward 19(1988)135

Effect of soft white wheat addition to alfalfa-grass forage on heifer gain, diet digestibility and in vitro digestion kinetics 24(1989)141

Alkali treatment

The effect on digestibility of a new technique for alkali treatment of straw 1(1976)73

Review article: the alkali treatment of straws 2(1977)105

Enzyme solubility - a method for evaluating the digestibility of alkali-treated straw 2(1977)205

The intake of steers and heifers fed on mixtures of alkali-treated or untreated straw with grass silage 2(1977)315

Ensiled alkali-treated straw. I. Effect of level and type of alkali on the composition and digestibility in vitro of ensiled barley straw 3(1978)117

Ensiled alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensiled or frozen alkali-treated straw and ryegrass silage 3(1978)133

Alkali treatment of straw for ruminants. I. Utilization of complete diets containing straw by beef cattle 3(1978)143

Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3(1978)227

Alkali treatment of straw for ruminants. II. Nutritive value of straw ensiled after alkali treatment. 3(1978)289

Laboratory evaluation of ensiled mixtures of alkali-treated cattle excreta and whole-crop maize 3(1978)335

- Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381
- Effect of level of sodium hydroxide treatment and volume of solution on the nutritive value of maize cobs 4(1979)1
- Performance of heifers fed on diets based on NaOH-treated maize cobs and the effect of supplementary urea and source of carbohydrates 4(1979)97
- Investigations of the treatment of sawdust for rabbit feeding. 1. Effect of sodium hydroxide treatment 6(1981)43
- Titratable groups and soluble phenolic compounds as indicators of the digestibility of chemically treated roughages 6(1981)123
- Simplified method for alkali treatment of low-quality roughages for use by small-holders in developing countries 6(1981)133
- Utilization of alkali-treated grain. 1. Alkali-treated grain in complete diets for steers and lambs 6(1981)273
- Utilization of alkali-treated grain. 2. Utilization by steers of diets based on hay or straw and mixed with either NaOH-treated or rolled barley 6(1981)347
- Utilization of alkali-treated grain. 3. Utilization by steers of NaOH-treated and rolled barley in silage-based diets 6(1981)355
- Ammonia and sodium hydroxide treatment of wheat straw in diets for fattening steers 7(1982)1
- Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7(1982)201
- Note on the lysinoalanine content of alkali-treated barley 7(1982)217
- Intake and digestibility in sheep fed on alkali-treated straw and concentrates, as complete diets or separate feeds 7(1982)271
- Digestibility of diets containing increasing levels of NaOH-treated or untreated wheat straw 7(1982)331
- Methods of improving the utilization of cereal straw by ruminants. I. Supplements of urea, molasses and dried grass and treatment with sodium hydroxide 8(1983)221
- Nutritive value for ruminants of sugar cane bagasse ensiled after spray treatment with different levels of NaOH 9(1983)1
- Utilization of alkali-treated and neutralized wheat straw-based rations for growing goat kids 9(1983)155
- Feeding value of alkali-treated whole rye (*Secale cereale* L.) grain for lactating cows and its digestibility for sheep 10(1983/1984)77
- Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91
- Effect of sodium hydroxide treatment and/or extrusion cooking on the nutritive value of peanut hulls 12(1984/1985)187
- A comparison of alkali treatment methods to improve the nutritive value of straw. I. Digestibility and metabolizability 12(1984/1985)295
- Organic waste ash as possible source of alkali for animal feed treatment 13(1985)237
- Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139
- A comparison of alkali treatment methods used to improve the nutritive value of straw. II. In sacco and in vitro degradation relative to in vivo digestibility 14(1986)215
- Corrugated cardboard and dried poultry manure as dietary ingredients for sheep 14(1986)265
- The response of screened manure fibre to ozone and sodium hydroxide treatments 15(1986)47
- The effect of grinding groundnut hulls either with or without alkali treatment on digestibility of diet and on ruminal and blood components 15(1986)69
- The nutritive value of barley for monogastric animals and for ruminants as affected by artificial drying and alkali and anaerobic storage 15(1986)285
- Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by goats 17(1987)219
- Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 17(1987)235
- Treatment of whole-crop cereals with alkali. 1. The influence of sodium hydroxide and ensiling on the chemical composition and in vitro digestibility of rye, barley and wheat crops harvested at increasing maturity and dry matter content 18(1987)257
- Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271
- Treatment of whole-crop cereals with alkali. 3. Voluntary intake and digestibility studies in sheep given immature wheat ensiled with sodium hydroxide, urea or ammonia 18(1987)283
- Treatment of whole-crop cereals with alkali. 4. Voluntary intake and growth in steers given

- wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55
- Fermentation and utilization by lambs of sugarcane harvested fresh and ensiled with and without NaOH. 4. Ruminant kinetics 23 (1989)323
- Alkali-treated soya flour for piglets 24(1989)1
- Upgrading barley straw for goats: the effectiveness of a sodium hydroxide and urea dip method 24(1989)57
- Alkaloid content**
- The alkaloid content of sweet lupin seed used in feeding trials on pigs and rats 2(1977)59
- Almond hulls**
- Nutritive value of almond hulls for sheep 8 (1983)63
- Amaranthus hybridus***
- Performance of calves on diets containing *Amaranthus* leaf meal 5(1980)193
- Amino acids, essential**
- Utilization of processed pig hair in poultry diets 3(1978)243
- Nitrogen metabolism in broiler chickens consuming the bacterial strain *Alcaligenes eutrophus* 5(1980)241
- Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21
- Palm kernel meal as a feed for poultry. 1. Composition of palm kernel meal and availability of its amino acids to chicks 16(1986/1987)179
- Amino acids, general**
- Amino acid supplementation of methanol-grown dried microbial cells in diets for young chicks 4(1979)63
- Ammonia treatment**
- Detoxification of aflatoxins in groundnut meal 2(1977)67
- Effect on digestibility and nitrogen content of barley straw of different ammonia treatments 2(1977)131
- The effect of ammoniation on the intake and nutritive value of straw 4(1979)169
- Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979)177
- Nutritive value of thermoammoniated and steam-treated maize stover II. Rumen metabolites and rate of passage 4(1979)187
- Preservation of high-moisture maize - a comparison of gaseous and liquid anhydrous ammonia with methylene-bis-propionate 5 (1980)337
- Effects of treatment of barley straw with anhydrous ammonia and supplementation with heat-treated protein meals on feed intake and liveweight performance of growing lambs 6(1981)145
- New crop residues and forages for Western Canada: assessment of feeding value in vitro and response to ammonia treatment 6(1981)257
- Different sources of ammonia for improving the nutritive value of low quality roughages 6(1981)377
- Ammonia and sodium hydroxide treatment of wheat straw in diets for fattening steers 7 (1982)1
- Studies on ammonia-treated straw. I. The effects of type and level of ammonia, moisture content and treatment time on the digestibility in vitro and enzyme soluble organic matter of oat straw 7(1982)45
- Studies on ammonia-treated straw. II. Fixation of ammonia in treated straw by spraying with acids 7(1982)53
- Effect of treatment of wheat straw with ammonia and peracetic acid on digestibility in vitro and cell wall composition 7(1982)325
- Ammoniation of wheat straw and native grass hay during baling of large round bales 8 (1983)35
- The effect of prolonged storage on the digestibility and nitrogen content of ammonia-treated barley straw 8(1983)147
- The feeding value for ruminants of straw and whole-crop barley and oats treated with anhydrous or aqueous ammonia or urea 8 (1983)247
- The effect of urease enzyme on treatment time and the nutritive value of straw treated with ammonia as urea 8(1983)271
- Methods of improving the utilization of cereal straw by ruminants. III. A note on the effect of ensiling straw treated with urea 9(1983)181
- Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185
- Composition and enzymatic digestibility of oregon grass straws 9(1983)283
- The effects of ammonia on the fermentation of chopped sugarcane 9(1983)291
- The effect of urea on the preservation and digestibility in vitro of perennial ryegrass 10(1983/1984)49

- Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65
- A note on the OECD collaborative experiment on ammonia-treated straw 10(1983/1984)89
- Chemical properties of fibre in relation to nutritive quality of ammonia-treated forages 10(1983/1984)155
- Ammonia treatment of straw: methods for treatment and feeding experience in Norway 10(1983/1984)173
- Ammonia versus sodium hydroxide treatment of straw for growing cattle 10(1983/1984)189
- A comparison between degradation *in vitro* and *in sacco* of constituents of untreated and ammonia-treated barley straw 10(1983/1984)199
- Digestibility studies on ammonia-treated straw 10(1983/1984)213
- Commercial experience of treating straw with ammonia 10(1983/1984)223
- The digestion of untreated and ammonia-treated barley straw in an artificial rumen 10(1983/1984)229
- Anhydrous ammonia-treated cereal straw for animal feeding. Some results from the Mediterranean area 10(1983/1984)239
- Ammonia-treated barley straw and rolled barley offered either together, in a mixed ration, or successively to beef steers 10(1983/1984)247
- Ammonia treatment of straw via the hydrolysis of urea. I. Effects of dry matter and urea concentrations on the rate of hydrolysis of urea 11(1984)103
- Ammonia treatment of straw via the hydrolysis of urea. II. Additions of soya bean (urease), sodium hydroxide and molasses; effects on the digestibility of urea-treated straw 11(1984)115
- The effect of anhydrous ammonia and moisture content on the preservation and chemical composition of perennial ryegrass hay 11(1984)159
- Influence of heat sterilization and ammoniation on straw composition and degradation by pure cultures of cellulolytic rumen bacteria 12(1984/1985)195
- Effect of ammonia treatment and processing of whole cotton plants as sole source of roughage in complete feeds for growing cross-bred calves 13(1985)93
- Effects of ethanolamine and ethylenediamine treatments on the composition and degradability of wheat straw 13(1985)131
- Methods of improving the utilization of cereal straw by ruminants. IV. The effect on intake of level of supplementation with urea and dried grass 13(1985)281
- Methods of improving the utilization of cereal straw by ruminants. V. The effect of molasses and dried grass 13(1985)293
- The use of barley straw as a feed 14(1986)29
- Treatment of barley straw with ammonia and sulphur dioxide gases under laboratory conditions 14(1986)41
- Effect of ammonia level and time of exposure to ammonia on nutritional and preservatory characteristics of dry and high-moisture coastal bermuda grass hay 14(1986)55
- Urea as a source of ammonia for improving the nutritive value of wheat straw 14(1986)67
- Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81
- Effects of ammonia treatment and processing of cotton straw as the sole source of roughage in complete feeds on rumen characteristics with Murrah buffaloes 14(1986)95
- Near infrared reflectance spectroscopy of untreated and ammoniated barley straw 15(1986)189
- Chemical composition and digestion *in vitro* of orchardgrass hay ammoniated by different techniques 17(1987)109
- Hyperexcitability in cattle fed ammoniated roughages 17(1987)121
- Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter disappearance *in sacco* 17(1987)179
- Treatment of whole-crop cereals with alkali. 3. Voluntary intake and digestibility studies in sheep given immature wheat ensiled with sodium hydroxide, urea or ammonia 18(1987)283
- Treatment of whole-crop cereals with alkali. 4. Voluntary intake and growth in steers given wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55
- Effects of ammonia and sulphur dioxide gases on the composition and digestion of barley straw 19(1988)121
- The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility *in vitro* and cell wall degradability 19(1988)159
- The effect of ammoniation on the nutritive value of wheat, barley and oat straws.

- II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements** 19(1988)173
- Fibre degradation in the rumens of animals receiving hay, untreated or ammonia-treated straw** 19(1988)277
- The effect of variation in the water content of wheat straw before ammoniation on its nutritive value for beef cattle** 20(1988)111
- The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage** 20(1988)335
- The feeding value of ammoniated flax straw, wheat straw and wheat chaff for beef cattle** 21(1988)57
- Near infra-red spectra of ammonia-treated straw and of isolated cell walls** 21(1988)209
- Upgrading of straw by ammoniation** 21(1988)287
- Effect of ammoniation and energy supplementation on the utilization of wheat straw by sheep** 22(1988)305
- Effect of urea treatment on rumen degradation characteristics of rice straws** 24(1989)83
- Influence of stage of forage maturity and ammoniation of wheat straw on ruminal degradation of wheat forage tissues** 24(1989)201
- Oven and stack ammoniation of grass hays.**
1. Changes in chemical composition in relation to digestibility in vitro and cell-wall degradability 24(1989)299
- Oven and stack ammoniation of grass hays.**
2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313
- Effects of urea treatment on chemical composition and in vitro digestibility on meadow hays of Northern Portugal** 25(1989)157
- Amprolium**
Particle size distribution of a feed premix containing amprolium 3(1978)211
- Amylase**
Prediction of the digestibility of ruminant feeds from their solubility in enzyme solutions 2(1977)153
- Antibiotics**
A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3(1978)89
- Efficacy of virginiamycin and a commercially-available lactobacillus probiotic in swine diets** 8(1983)69
- Apple pomace**
Feeding value of apple pomace silage for sheep 11(1984)189
- Some properties of apple pomace ensiled with and without additives** 15(1986)57
- Nutritive value of apple pomace for ruminants** 16(1986/1987)311
- Arginine**
Limiting amino acids in dried microbial cells given to young turkeys 4(1979)209
- Arhar**
Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315
- Artichoke**
The chemical composition and nutritive value for ruminants of four possible energy crops and their residues of anaerobic fermentation 15(1986)197
- Aspergillus fumigatus**
Production of fungal protein from rasped fresh cassava roots using 200- and 3000-litre fermentors 8(1983)313
- Avoparcin**
Effect of avoparcin on performance, body composition and meat quality of finishing bulls 8(1983)237
- In vitro effects of avoparcin on protein degradability and rumen fermentation** 15(1986)215
- ## B
- Bacteria**
A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3(1978)89
- Nitrogen metabolism in broiler chickens consuming the bacterial strain *Alcaligenes eutrophus*** 5(1980)241
- Protein quality studies on rats fed on cassava-root meal enriched with *Cephalosporium eichhorniae* 152 or with conventional plant-protein supplements** 6(1981)197
- The effect of some pre-ensiling treatments on silage composition and nitrogen disappearance in the rumen** 8(1983)259
- Electron microscopy of bacteria involved in the digestion of plant cell walls** 10(1983/1984)93
- Effect of absence of ciliate protozoa on rumen fluid volume, flow rate and bacterial populations in sheep** 10(1983/1984)145

- Stability of viable lactobacillus products 11 (1984)261
- Influence of heat sterilization and ammoniation on straw composition and degradation by pure cultures of cellulolytic rumen bacteria 12(1984/1985)195
- Effects of molasses/urea and bacterial inoculant additives on silage quality, dry matter recovery, and feeding value for cattle 12 (1984/1985)205
- The chemical composition of grass silages prepared with and without pre-treatment with inoculants containing *Lactobacillus plantarum* 13(1985)269
- Evaluation of a dual flow continuous culture system for estimating bacterial fermentation in vivo of mixed diets containing various soya bean products 16(1986/1987)51
- Population changes of fibrolytic rumen bacteria in the presence of phenolic acids and plant extracts 19(1988)261
- The establishment of rumen microbial communities 21(1988)69
- Possibilities of the use of recombinant-DNA techniques with rumen micro-organisms 21 (1988)161
- Characterization of the flora of the large bowel of pigs: a status report 23(1989)79
- Fibre-degrading microorganisms in the monogastric digestive tract 23(1989)91
- Assessing the importance of host-derived polysaccharides as carbon sources for bacteria growing in the human colon 23(1989) 109
- In vitro fermentation of dietary fiber by human fecal organisms 23(1989)121
- Inoculation of corn silage and high-moisture corn with lactic acid bacteria and its effects on the subsequent fermentations and on feedlot performance of beef steers 25(1989) 23
- Evaluation of various methods for protecting soya bean protein from degradation by rumen bacteria 25(1989)111
- In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations 25(1989)169
- Bagaruwa seeds**
Biochemical evaluation of bagaruwa seeds (*Acacia nilotica*) for use as livestock feed 11(1984)45
- Balanites kernel cake**
The use of Balanites kernel cake in a diet for fattening sheep 9(1983)301
- Banana stalk**
The value of banana stalk as a feed for goats 15(1986)95
Nutritive value of banana stalk (*Musa cavendishi*) as a feed for sheep 22(1988)327
- Barley**
Lupin-seed meal (*Lupinus angustifolius* cv. Uniwhite) as a protein supplement to barley-based diets for growing pigs 1(1976) 631
Swedes (*Brassica napus*) for growing pigs: chemical composition and use as a replacement for barley in the diet 2(1977)31
A comparison between meals prepared from the seeds of different varieties of lupin as protein supplements to barley-based diets for growing pigs 2(1977)49
The effect of supplementary protein on the performance of pigs given swedes (*Brassica napus*) as a replacement for barley in the diet 2(1977)277
The use of whole barley diets fortified with solutions of urea, minerals and vitamins for lambs 2(1977)307
Nutritive value of leaf protein coagulum and cereals mixtures 4(1979)91
The determination of available lysine in barley using *Tetrahymena pyriformis* W. 5 (1980)33
A comparison of three methods for the determination of available lysine in barley 5 (1980)43
Utilization of alkali-treated grain. 1. Alkali-treated grain in complete diets for steers and lambs 6(1981)273
Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6(1981)309
Utilization of alkali-treated grain. 2. Utilization by steers of diets based on hay or straw and mixed with either NaOH-treated or rolled barley 6(1981)347
Utilization of alkali-treated grain. 3. Utilization by steers of NaOH-treated and rolled barley in silage-based diets 6(1981)355
Influence of increasing levels of β -glucanase on the productive value of barley diets for broiler chickens 7(1982)351
Nutritional value of waxy barleys for broiler chickens 8(1983)25
Evaluation of optimum lysine and threonine supplements to a wheat and barley-based diet in rats 8(1983)163
The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation

- and digestibility *in vitro* of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37
- Effect of storage and processing of barley on its utilization by pigs and rats 9(1983)89
- Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983)99
- Associative effects of mixed feeds. II. The effect of dietary addition of bicarbonate salts on the voluntary intake and digestibility of diets containing various proportions of hay and barley 10(1983/1984)31
- Anaerobically stored high moisture cereals for growing pigs 12(1984/1985)159
- Feeding value of hydroponically sprouted barley for poultry and pigs 13(1985)183
- Nutrient content of hydroponically sprouted barley 13(1985)191
- Apparent ileal and faecal digestibility of amino acids in barley-based diets supplemented with soya bean meal or canola meal for growing pigs 14(1986)183
- The effect of monensin on rumen fermentation in sheep fed on all-roughage and barley/roughage diets 15(1986)7
- Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21
- The effect of β -glucanase on the utilization of starch and nitrogen by broiler chickens fed on barley of low- or high-viscosity 15(1986)83
- The nutritive value of barley for monogastric animals and for ruminants as affected by artificial drying and alkali and anaerobic storage 15(1986)285
- Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261
- Nutritional value for chickens of dried distillers-spent-grain from barley and dehulled barley 17(1987)145
- The effect of dietary barley on carbohydrate digestibility of sulphur dioxide-treated wheat straw by sheep 18(1987)55
- Treatment of whole-crop cereals with alkali. 1. The influence of sodium hydroxide and ensiling on the chemical composition and *in vitro* digestibility of rye, barley and wheat crops harvested at increasing maturity and dry matter content 18(1987)257
- Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271
- The nutritive value of hulless barley for swine 19(1988)191
- The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988)259
- Amino acids flow to and absorption from the small intestine of sheep fed barley and sulfur dioxide-treated straw at different ratios 22(1988)147
- A comparison of wheat- or barley-based diets given *ad libitum* as meal or pellets to finishing pigs 22(1988)263
- The effect of mechanical treatment on the nutritive value of high-moisture barley and the performance of growing-finishing pigs 24(1989)121
- Effect of enzyme supplementation on the nutritive value of hulless barley in chicken diets 24(1989)233
- Factors affecting the digestibility by pigs of energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253
- Effect of formaldehyde treatment on the rumen digestion of barley and wheat grain and its influence on hay intake and digestion by sheep 25(1989)55
- The effect of β -glucanase supplementation on nutrient digestibility and growth in broilers given diets containing barley, oat groats or wheat 25(1989)193
- Barley grain**
- Note on the lysinoalanine content of alkali-treated barley 7(1982)217
- Milling barley and lupin grain in diets for cattle 12(1984/1985)57
- Microbial dynamics in barley grain stored under controlled atmosphere 19(1988)299
- Barley meal**
- The influence of feeding level and level of dietary inclusion on the digestibility of barley meal in the pig 2(1977)41
- Barley straw**
- Effect on digestibility and nitrogen content of barley straw of different ammonia treatments 2(1977)131
- Ensiled alkali-treated straw. I. Effect of level and type of alkali on the composition and digestibility *in vitro* of ensiled barley straw 3(1978)117

- Alkali treatment of straw for ruminants. I. Utilization of complete diets containing straw by beef cattle 3(1978)143
- Alkali treatment of straw for ruminants. II. Nutritive value of straw ensiled after alkali treatment. 3(1978)289
- Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381
- Effects of treatment of barley straw with anhydrous ammonia and supplementation with heat-treated protein meals on feed intake and liveweight performance of growing lambs 6(1981)145
- Studies on ammonia-treated straw. II. Fixation of ammonia in treated straw by spraying with acids 7(1982)53
- The feeding value for ruminants of straw and whole-crop barley and oats treated with anhydrous or aqueous ammonia or urea 8(1983)247
- The effect of urease enzyme on treatment time and the nutritive value of straw treated with ammonia as urea 8(1983)271
- Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65
- A comparison between degradation in vitro and in sacco of constituents of untreated and ammonia-treated barley straw 10(1983/1984)199
- Digestibility studies on ammonia-treated straw 10(1983/1984)213
- The digestion of untreated and ammonia-treated barley straw in an artificial rumen 10(1983/1984)229
- Ammonia-treated barley straw and rolled barley offered either together, in a mixed ration, or successively to beef steers 10(1983/1984)247
- A comparison of alkali treatment methods to improve the nutritive value of straw. I. Digestibility and metabolizability 12(1984/1985)295
- Treatment of barley straw with ammonia and sulphur dioxide gases under laboratory conditions 14(1986)41
- A comparison of alkali treatment methods used to improve the nutritive value of straw. II. In sacco and in vitro degradation relative to in vivo digestibility 14(1986)215
- Evaluation of the increase in feeding value of barley straw following mild treatment with sulphur dioxide 15(1986)13
- Near infrared reflectance spectroscopy of untreated and ammoniated barley straw 15(1986)189
- Effect of straw treatment and laboratory of incubation on degradation of neutral detergent residue in the rumen 18(1987)75
- Fresh and dried brewer's grains as protein supplements to barley straw diets given to pregnant beef cows 19(1988)33
- Effects of ammonia and sulphur dioxide gases on the composition and digestion of barley straw 19(1988)121
- The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility in vitro and cell wall degradability 19(1988)159
- The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173
- Fibre degradation in the rumsens of animals receiving hay, untreated or ammonia-treated straw 19(1988)277
- The effect of five different supplements on the degradation of straw in sheep given untreated barley straw 19(1988)289
- Upgrading of straw by ammoniation 21(1988)287
- Effect of soaking, ensilage and hydrogen peroxide treatment of barley straw on rumen degradability 22(1988)295
- Upgrading barley straw for goats: the effectiveness of a sodium hydroxide and urea dip method 24(1989)57
- Basic steel slag**
Biological availability of phosphorus from basic steel slag 11(1984)75
- Bean, winged**
The proximate composition and nutritive value of the winged bean *Psophocarpus tetragonolobus* (L.) DC for broilers 11(1984)231
- Bermuda grass**
Effect of ammonia level and time of exposure to ammonia on nutritional and preservative characteristics of dry and high-moisture coastal bermuda grass hay 14(1986)55
- Bermudagrass**
Intake, digestion and daily gain by cattle consuming bermudagrass (*Cynodon dactylon*) and supplemented with different combinations of ground corn, vegetable oil, urea and corn gluten and blood meals 25(1989)99

β -glucanase

Influence of increasing levels of β -glucanase on the productive value of barley diets for broiler chickens 7(1982)351

The effect of β -glucanase on the utilization of starch and nitrogen by broiler chickens fed on barley of low- or high-viscosity 15(1986) 83

The effect of β -glucanase supplementation on nutrient digestibility and growth in broilers given diets containing barley, oat groats or wheat 25(1989)193

Bicarbonate salts

Associative effects of mixed feeds. II. The effect of dietary addition of bicarbonate salts on the voluntary intake and digestibility of diets containing various proportions of hay and barley 10(1983/1984)31

Blood

The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21

Treatment of the haem fraction of abattoir blood with formic acid to produce a solid state, low dry matter protein supplement for use in finishing-pig diets 2(1977)367

Blood meal as a source of protein for grower/finisher pigs 3(1978)191

Ensiling of rumen contents and blood with wheat straw 18(1987)87

Comparison of cottonseed cake and dried blood as nitrogen sources for growing Sudan Nubian goats 22(1988)333

Blood meal

Protein quality of mixtures of animal protein meals 9(1983)79

Effect of blood meal processing method on performance and nitrogen utilization by growing cattle 24(1989)337

Bone meal

Protein quality of mixtures of animal protein meals 9(1983)79

Meat and bone and fish meals in balancer feeds for choice-fed broilers 11(1984)221

Botanical composition

Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71

Brewer's dried grains

Influence of the drying process on the composition of brewer's dried grains 6(1981)163

Evaluation of brewer's dried grains in the diets of laying hens 6(1981)169

Evaluation of brewer's dried grains in the diets of broiler chickens 6(1981)179

Fresh and dried brewer's grains as protein supplements to barley straw diets given to pregnant beef cows 19(1988)33

Buckwheat

A nutritional evaluation of buckwheat (*Fagopyrum esculentum*) 3(1978)95

Buffalo

Nutritive evaluation of uromol-wheat straw as a basal ration for buffaloes 13(1985)103

Effects of ammonia treatment and processing of cotton straw as the sole source of roughage in complete feeds on rumen characteristics with Murrah buffaloes 14(1986) 95

Colonization of guinea grass by anaerobic rumen fungi in swamp buffalo and cattle 22 (1988)161

Buffers

Comparison between two methods of feeding concentrate to sheep given a diet high or low in concentrate with or without buffering substances 16(1986/1987)37

Comparison of buffers on rumen functions, turnover rate and gastric secretions in Holstein steers 17(1987)257

Efficacy of supplemental dietary neutralizing agents for lactating dairy cows. A review 23(1989)277

By-products-animal waste

The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21

Nutritive value of rumen contents for monogastric animals 2(1977)351

Utilization by sheep of dried pig faeces containing a high concentration of copper 3 (1978)31

Digestibility by cattle of diets containing dried pig faeces 3(1978)155

Utilization of processed pig hair in poultry diets 3(1978)243

Protein digestibility of hydrolyzed hog hair meal for pigs 3(1978)323

The effects of air dried poultry manure on performance, organ weights and haematological measurements of weaner pigs reared in the tropics 4(1979)23

Utilization of cattle manure for single-cell protein production with *Chaetomium cellulolyticum* 4(1979)199

Digestibility of wet cage layer excreta and an excreta-supplemented diet by cattle as determined by two methods 5(1980)77

Utilization of cattle manure for *Torula yeast* production from straw hydrolysates 5(1980) 175

Performance of fattening cattle and lactating beef cows on diets high in chemically

- treated wheat straw and poultry litter 5 (1980)299
- Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51
- Fermentation characteristics and feeding value of ensiled poultry litter containing wheat straw, bagasse or sawdust 7(1982)341
- Nutritive value of ensiled broiler litter for cattle 8(1983)19
- The utilization by broiler chickens of poultry offal hydrolysed in formic acid 11(1984)247
- Chemical preparation of NaOH-keratin hydrolysate for improving the nutritive value of wheat straw 13(1985)215
- Thermofractographic determination of monocarbonyl compounds in animal waste fats used as feed fat 13(1985)227
- Organic waste ash as possible source of alkali for animal feed treatment 13(1985)237
- Effects of processing time and moisture content on amino acid composition and nitrogen characteristics of feather meal 14(1986)279
- The utilisation by growing pigs of ruminant offal hydrolysed in formic acid 15(1986)273
- The effect of enzymatic treatment on amino acid content and nitrogen characteristics of feather meal 16(1986/1987)151
- Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 17 (1987)235
- Ensiling of rumen contents and blood with wheat straw 18(1987)67
- Evaluation of cardboard and dried poultry waste as feed ingredients for ruminants 18 (1987)209
- Animal excreta as ruminant feed - scope and limitations under Indian conditions 19 (1988)1
- Slaughterhouse by-products preserved by *Lactobacillus plantarum* fermentation as feed for mink and foxes 20(1988)287
- Nutritive value of rice straw (*Oryza sativa*) ensiled with animal excreta and rumen digesta 24(1989)69
- Ensiling characteristics of cattle waste fermented with wheat straw and green berseem (*Trifolium alexandrinum*) 25(1989)149
- By-products-apple pomace**
Some properties of apple pomace ensiled with and without additives 15(1986)57
- By-products-banana stalk**
The value of banana stalk as a feed for goats 15(1986)95
- By-products-brewer's dried grains**
Influence of the drying process on the composition of brewer's dried grains 6(1981)163
Evaluation of brewer's dried grains in the diets of laying hens 6(1981)169
Evaluation of brewer's dried grains in the diets of broiler chickens 6(1981)179
- By-products-canning industry residues**
By-products from the canning industry as feedstuff for ruminants: digestibility and its prediction from chemical composition and laboratory bioassays 25(1989)67
- By-products-cardboard**
Evaluation of cardboard and dried poultry waste as feed ingredients for ruminants 18 (1987)209
- By-products-cocoa**
Studies on the feeding value of agro-industrial by-products. III. Digestibility of cocoa-pod and cocoa-pod-based diets by ruminants 13 (1985)249
Studies on the feeding value of agro-industrial by-products: strategies for improving the utilisation of cocoa-pod-based diets by ruminants 20(1988)189
- By-products-coffee residue**
A note on the nutritive value of dried instant coffee residue for broiler chickens and turkey poult 10(1983/1984)285
The nutritional value of dried coffee pulp (DCP) in broiler chickens's diets 22(1988)139
- By-products-cottonseed**
A review of the nutritive value and utilization of whole cottonseed, cottonseed meal and associated by-products by dairy cattle 18 (1987)89
- By-products-crop residues**
New crop residues and forages for Western Canada: assessment of feeding value in vitro and response to ammonia treatment 6(1981)257
Improving utilization of crop residues. Evaluation of two chemicals used to improve digestibility of crop residues 9(1983)205
The nutritional value of crop residue components from several wheat cultivars grown at different fertilizer levels 11(1984)301
Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter disappearance in sacco 17(1987)179
Supplementation of rice straw with crop by-products. I. Legume straw supplementation 19(1988)43
Nutritive value of banana stalk (*Musa cavendishii*) as a feed for sheep 22(1988)327

By-products-culinary waste

Chemical composition of culinary wastes and their potential as a feed for ruminants 5 (1980)205

By-products-distillery waste

Distillery effluents as animal feed: the use of condensed beet molasses stillage (CBMS) in broiler feeding 5(1980)233

Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47

Nutritional value for chickens of dried distillers-spent-grain from barley and dehulled barley 17(1987)145

By-products-dried bakery product

Variation in nutrient content of samples of dried bakery product 7(1982)419

By-products-fish silage

Fish silage as a protein supplement for early-weaned calves 17(1987)165

By-products-general

Suitability of cell-wall constituents as predictors of organic matter digestibility in some tropical and subtropical by-products 7(1982)191

Variations and prevision of the in sacco dry matter digestion of concentrates and by-products 13(1985)7

Use of the rumen simulation technique (RUSITEC) to compare microbial digestion of various by-products 21(1988)197

By-products-grapevine

Digestibility in vivo of ensiled grapevines (branches and leaves): influence of the system of analysis in the detergent fibre scheme on the prediction of digestibility 22 (1988)173

By-products-maize whole stillage

Maize whole stillage as a supplemental protein source for growing lambs and finishing steers 11(1984)181

By-products-oil industry

Composition of by-products of the essential-oil industry and their potential as feeds for ruminants 6(1981)115

By-products-olive cake

Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139

By-products-orange peel

The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20 (1988)335

By-products-palm oil mill effluent

The intake and digestibility in goats of *Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989)45

By-products-sal seed meal

Tannins in sal seed (*Shorea robusta*) and sal seed meal limit their utilization as livestock feeds 7(1982)161

By-products-sawdust

Investigations of the treatment of sawdust for rabbit feeding. 1. Effect of sodium hydroxide treatment 6(1981)43

Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51

By-products-silage effluent

The use of effluent from grass silage in the diet of finishing pigs. I. Variation in composition of effluents 4(1979)263

The use of effluent from grass silage in the diet of finishing pigs. II. Assessment of nutritive value of fresh and stored effluent 4(1979) 275

Investigations on the effect of sodium bentonite and other moisture absorbents on the production of effluent from grass silage 8(1983)107

By-products-sisal waste

Feed value of Tanzanian sisal wastes: chemical composition and digestibility in vitro 14 (1986)255

By-products-vine

Composition of vine-branches with leaves of *Vitis vinifera* L. fresh and ensiled using different additives (previous results) 16(1986/1987)89

By-products-whey

The nutritive value of liquid whey, either sour or sweet, when given in restricted amounts to the growing pig 3(1978)163

The evaluation of fresh sweet liquid whey as a protein supplement to maize for growing pigs 14(1986)171

By-products-wood residue

Composition of kermes oak browse as affected by shade and stage of maturity 21(1988)1

Pilot plant production of sulphur dioxide-treated mesquite wood. II. Digestibility and feedlot performance by lambs 22(1988) 61

C

Cabbage

Cabbage (*Brassica oleracea*) in the diet of growing-finishing pigs 5(1980)69

Studies on the nutritive value of roots, cabbage and grass silage for growing commercial rabbits 13(1985)299

Calcium

Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Compositional quality of certain unconventional calcium and phosphorus sources in India for use as mineral supplements for livestock 23(1989)343

Calcium, magnesium

Solubility and flow of calcium, magnesium and phosphorus in the digestive tract of sheep given maize or alfalfa silages 9(1983)131

Calcium hydroxide treatment

The effect of calcium hydroxide treatment on the digestibility of chopped oat straw supplemented with a solution containing urea, calcium, phosphorus, sodium, trace elements and vitamins 7(1982)93

The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91

The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12(1984/1985)141

Digestion rates in and outflow rates from the rumen of sheep fed untreated or calcium hydroxide-treated wheat straw 25(1989)179

Calcium soap

The effects on feed intake, growth and digestibility of nutrients of including calcium soaps of fat in diets for young calves 14(1986)103

Calcium-soaps

Feed intake, milk yield and milk composition by replacing unprotected fat by Ca-soaps for dairy cows 22(1988)193

Calorimetry

A calorimetric investigation of the nutritive value of formic acid- and formaldehyde-formic acid-treated grass silages 6(1981)245

Canavanine

Canavanine content and toxicity of sesbania leaf meal for growing chicks 25(1989)137

Cane

Proximate chemical analysis of final cane molasses and effect of feeding 30% molasses on intestinal sucrase and maltase activities in the rat 3(1978)15

Canola meal

Apparent ileal and faecal digestibility of amino acids in barley-based diets supplemented with soya bean meal or canola meal for growing pigs 14(1986)183

Canola meal as a protein supplement for growing-finishing pigs 18(1987)37

The nutritive value of canola meal for early-weaned pigs 18(1987)45

Factors affecting the digestibility by pigs of energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253

Canola rapeseed

The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303

Carbon tetrachloride

Long term feeding of concentrate fumigated with carbon tetrachloride to sheep 2(1977)77

Cardboard

Corrugated cardboard and dried poultry manure as dietary ingredients for sheep 14(1986)265

Evaluation of cardboard and dried poultry waste as feed ingredients for ruminants 18(1987)209

Carotenes

Losses of beta-carotene in red clover in an acid medium during ensiling 4(1979)81

The enzymic nature of the degradation of beta-carotene in red clover and in other forage crops during silage-making with acid additives 5(1980)59

Losses of beta-carotene in unwilted forage crops during silage-making and feeding 9(1983)63

Investigations into the effects of processing on the retention of the carotenoid fractions of *Leucaena leucocephala* during storage, and the effects of processing on mimosine concentration 9(1983)307

Casein

The apparent and true ileal digestibility of nitrogen and amino acids in lactic casein for the growing pig 16(1986/1987)169

Nutritive value of Nigerian rubber seed (*Hevea brasiliensis*) meal and oil. III. Performance characteristics, relative organ weights, hematocrit and plasma metabolites of growing female rats fed corn diets containing rubber seed meal, soya bean meal or casein 20(1988)125

Cassava

Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977)7

Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73

Problems in the use of cassava as animal feed 3(1978)345

Cassava meal and cassava peel meal in diets for growing pigs 8(1983)211

Effect of cassava-based diets varying in cyanide content on the performance and physiopathology of the African giant rat (*Cricetomys gambianus* Waterhouse) 11(1984)1

Cassava for animal feeding: effect of variety and plant age on production of leaves and roots 11(1984)49

Cassava chips

Effect of variety and plant age on the cyanide content of whole-root cassava chips and its reduction by sun-drying 11(1984)57

Cassava flour

The maize replacement value of cassava flour in rations for growing ewe lambs 20(1988)233

Cassava meal

Protein quality studies on rats fed on cassava-root meal enriched with *Cephalosporium eichhorniae* 152 or with conventional plant-protein supplements 6(1981)197

Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227

Cassava meal as supplement to napier grass diets for growing sheep and goats 18(1987)197

Effects of oven-dried cassava peel meal on the performance and carcass characteristics of broiler chickens 24(1989)247

Cassava roots

Production of fungal protein from rasped fresh cassava roots using 200- and 3000-litre fermentors 8(1983)313

Cassava tops

Net energy for gain (NE_{gain}) of cassava tops 20(1988)97

Cattle-beef

The use of techniques of maize cultivation to maximise feed for beef production 1(1976)229

Maize silage with or without NPN, dehydrated whole-crop maize pellets or high moisture maize grain for finishing bulls 1(1976)347

Maize grain or ears in concentrate diets for young fattening bulls 1(1976)393

Rate of growth, feed utilisation and digestibility observed with diets containing ensiled maize grain or maize ears fed alone or in mixtures 1(1976)401

Utilising the maize crop in various forms for beef production 1(1976)417

Levels, sources and methods of nitrogen supplementation of maize silage for beef production 1(1976)465

Economic aspects of high-plane feeding of bulls with maize silage 1(1976)513

For beef cattle, all forms of maize can be used 1(1976)521

Bull-beef production from the maize crop in Italy in large and small units 1(1976)531

Dried sugar beet pulp as a high energy feed for beef cattle 1(1976)643

Abomasal and plasma nitrogen constituents of steers fed on different physical forms of shelled maize 2(1977)225

Alkali treatment of straw for ruminants. I. Utilization of complete diets containing straw by beef cattle 3(1978)143

Performance of fattening cattle and lactating beef cows on diets high in chemically treated wheat straw and poultry litter 5(1980)299

Utilization of alkali-treated grain. 1. Alkali-treated grain in complete diets for steers and lambs 6(1981)273

Utilization of alkali-treated grain. 2. Utilization by steers of diets based on hay or straw and mixed with either NaOH-treated or rolled barley 6(1981)347

Utilization of alkali-treated grain. 3. Utilization by steers of NaOH-treated and rolled barley in silage-based diets 6(1981)355

Ammonia and sodium hydroxide treatment of wheat straw in diets for fattening steers 7(1982)1

The digestion in the small intestine of young bulls of the protein of rapeseed meal

- treated or untreated with formaldehyde 7 (1982)225
- Monensin-sodium as a performance-promoting additive for fattening bulls and its impact on carcass and meat quality characteristics 7(1982)401
- Effect of avoparcin on performance, body composition and meat quality of finishing bulls 8(1983)237
- Ammonia versus sodium hydroxide treatment of straw for growing cattle 10(1983/1984) 189
- Ammonia-treated barley straw and rolled barley offered either together, in a mixed ration, or successively to beef steers 10 (1983/1984)247
- Maize whole stillage as a supplemental protein source for growing lambs and finishing steers 11(1984)181
- The use of kidney beans (*Phaseolus vulgaris*) as protein supplements in diets for young rapidly growing beef steers 12(1984/1985) 1
- Milling barley and lupin grain in diets for cattle 12(1984/1985)57
- Improving the intake and nutritive value of wheat straw for beef cows 12(1984/1985) 125
- Wheat silage for fattening young male cattle 12(1984/1985)253
- Effect of sugarcane condensed molasses solubles, sugarcane molasses and monensin on performance and volatile fatty acid production in finishing steers 12(1984/1985)275
- Methods of improving the utilization of cereal straw by ruminants. IV. The effect on intake of level of supplementation with urea and dried grass 13(1985)281
- Methods of improving the utilization of cereal straw by ruminants. V. The effect of molasses and dried grass 13(1985)293
- Effect of dietary nitrogen source on microbial protein synthesis, dietary protein degradation and nutrient digestion in steers 14 (1986)117
- Effects of dietary concentrates on rumen digestion of fibrous feedstuffs 14(1986)193
- Feeding value of sprouted wheat (*Triticum aestivum*) for beef cattle finishing diets 15 (1986)113
- High grain content maize silage in fattening diets of young male cattle 16(1986/1987)63
- Relative intakes, digestibility and bacterial protein synthesis by sheep and cattle fed high-roughage diets 16(1986/1987)75
- The effects of different sources of nitrogen supplementation on the digestion of fibre components in the rumen of steers 17(1987) 65
- Comparison of buffers on rumen functions, turnover rate and gastric secretions in Holstein steers 17(1987)257
- Effects of whole or ground corn with different forms of hay in 85% concentrate diets on digestion and passage rate in beef heifers 18(1987)151
- Fresh and dried brewer's grains as protein supplements to barley straw diets given to pregnant beef cows 19(1988)33
- Treatment of whole-crop cereals with alkali. 4. Voluntary intake and growth in steers given wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55
- The effect of variation in the water content of wheat straw before ammoniation on its nutritive value for beef cattle 20(1988)111
- Effects of ruminal degradability of the protein supplement on the utilization of maize-stover silage by growing beef cattle 20 (1988)241
- The utilization by beef steers of a soft high lysine and a hard normal size maize cultivar in whole grain feedlot diets 21(1988)31
- The feeding value of ammoniated flax straw, wheat straw and wheat chaff for beef cattle 21(1988)57
- Rumen degradability of nitrogen in typical cultivated grasses and leguminous fodders 22(1988)79
- Effects of added dietary urea on the utilization of maize stover silage by growing beef cattle 22(1988)227
- Effect of level and frequency of maize supplementation on the utilization of low-quality grass hay by beef cows 24(1989)129
- Inoculation of corn silage and high-moisture corn with lactic acid bacteria and its effects on the subsequent fermentations and on feedlot performance of beef steers 25(1989) 23
- The effect of a protein energy supplement on pasture protein and fibre digestion in the rumen of grazing steers 25(1989)39
- Water-drinking frequency and quantity of water drunk by steers fed hay or fresh forage 25(1989)207
- Cattle-calf**
- Methanol-fermentation protein in veal calf nutrition 1(1976)33
- Effect of weight at slaughter on the efficiency of fattening young cattle 1(1976)505
- The replacement of skim milk by dried microbial cells and whey in milk diets for calves 2(1977)85

- Digestibility of protein and amino acids of a fermentation single-cell protein for veal calves 2(1977)377
- Ensilaged alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensilaged or frozen alkali-treated straw and ryegrass silage 3(1978)133
- Spent tea leaf as a ruminant feed 3(1978)219
- Performance of calves on diets containing *Amaranthus* leaf meal 5(1980)193
- Effects of sodium chloride and sodium bicarbonate in the diet on the performance of calves 5(1980)309
- Performance of calves fed on milk replacers containing fish protein hydrolysate 7(1982)141
- Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375
- Effect of ammonia treatment and processing of whole cotton plants as sole source of roughage in complete feeds for growing cross-bred calves 13(1985)93
- The effects on feed intake, growth and digestibility of nutrients of including calcium soaps of fat in diets for young calves 14(1986)103
- Utilization of whole rape (*Canola*) seed and sunflower seeds as sources of energy and protein in calf starter diets 15(1986)101
- Effect of formaldehyde-treated soya bean meal and urea in starters on nitrogen quality, degradability in sacco, sheep digestibility and calf performance 16(1986/1987)287
- Fish silage as a protein supplement for early-weaned calves 17(1987)165
- Fish protein concentrates for pre-ruminant calves: effects of processing temperatures and source of fish on protein digestibility and biological value 18(1987)181
- The effects of method of processing of starters, tallow inclusion and roughage supplementation on the performance of early-weaned calves 19(1988)231
- Evaluation of pea protein concentrate as a source of protein in milk replacers for Holstein calves 24(1989)267
- Effect of blood meal processing method on performance and nitrogen utilization by growing cattle 24(1989)337
- Cattle-dairy**
- Forage maize in the rations of dairy cows and fattening cattle in Bulgaria 1(1976)45
- The relationship between the intake of herbage grazed by dairy cows and its digestibility 1(1976)53
- The digestibility and nutritive value for dairy heifers of rations containing cellulose fibre 2(1977)287
- Maize silage and whole maize plant pellets as the basic forages in complete feeds for milk cows 3(1978)23
- Performance of heifers fed on diets based on NaOH-treated maize cobs and the effect of supplementary urea and source of carbohydrates 4(1979)97
- Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315
- Dry matter, protein, energy and fibre intake by dairy heifers grazing a Rhodes grass (*Chloris gayana*) pasture 6(1981)15
- Feed consumption and performance of dairy cows with alternate feedings of grass silage and maize silage 6(1981)337
- Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7(1982)201
- Studies on fermentation of sorghum silage during storage, and its effect on milch animals 9(1983)143
- Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197
- Feeding value of alkali-treated whole rye (*Secale cereale* L.) grain for lactating cows and its digestibility for sheep 10(1983/1984)77
- The prediction of the net energy value of compound feedstuffs for dairy cattle 11(1984)137
- The effect of intake and hay: concentrate ratio upon digestibility and digesta passage 11(1984)167
- Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279
- The use of barley straw as a feed 14(1986)29
- Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81
- Sugar beet pulp products for dairy cows 15(1986)123
- Concentrates for dairy cows: effects of feeding method, proportion in diet and type 15(1986)167
- Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammer-milled lupin grain in oats hay or oats pasture based diets 16(1986/1987)143

- Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261
- A review of the nutritive value and utilization of whole cottonseed, cottonseed meal and associated by-products by dairy cattle 18(1987)89
- A note on the use of sugar-beet pulp silage and molasses in the diet of lactating dairy cows 20(1988)13
- The substitution of grass silage by a straw and concentrate mixture for dairy cows 20(1988)135
- The intake and digestion of maize silage-based diets by dairy cows and sheep 20(1988)299
- Feed intake and milk yield at increasing supplement of a palmitic and stearic acid-rich type of fat in comparison with animal fat 22(1988)179
- Feed intake, milk yield and milk composition by replacing unprotected fat by Ca-soaps for dairy cows 22(1988)193
- Efficacy of supplemental dietary neutralizing agents for lactating dairy cows. A review 23(1989)277
- Response of high-producing dairy cows to high levels of dried corn gluten feed 24(1989)191
- Discriminating power of the nylon bag technique and the pepsin-cellulase method to detect less commonly used components included in concentrates for dairy cows 25(1989)1
- Rumen ingesta kinetics of cell wall components in dairy cows 25(1989)89
- Cattle-general**
- The potato (*Solanum tuberosum*) as a source of nutrients for pigs, calves and fowl - a review 2(1977)171
- The intake of steers and heifers fed on mixtures of alkali-treated or untreated straw with grass silage 2(1977)315
- Differences between cattle and sheep in their digestion and relative intake of a mature tropical grass hay 3(1978)41
- Estimation of the digestibility of low-quality hays by cattle from measurements made with sheep 3(1978)51
- Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73
- Digestibility by cattle of diets containing dried pig faeces 3(1978)155
- Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3(1978)227
- Digestibility of wet cage layer excreta and an excreta-supplemented diet by cattle as determined by two methods 5(1980)77
- Nutritive value of ensiled broiler litter for cattle 8(1983)19
- Comparative digestibility of feedstuffs by sheep and cows 12(1984/1985)47
- Effects of molasses/urea and bacterial inoculant additives on silage quality, dry matter recovery, and feeding value for cattle 12(1984/1985)205
- The effect of nitrogen supplements to maize silage on the amount and composition of the nitrogen fraction entering the intestine of cattle 13(1985)25
- Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47
- Studies on the feeding value of agro-industrial by-products. III. Digestibility of cocoa-pod and cocoa-pod-based diets by ruminants 13(1985)249
- The energetic efficiency of rumen microbial protein synthesis in cattle given silage-based diets 13(1985)255
- Hyperexcitability in cattle fed ammoniated roughages 17(1987)121
- Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19
- The effect of grinding and pelleting hay on digestibility, fermentation rate, digesta passage and rumen and faecal particle size in cows 19(1988)145
- Rumen fermentation, urease activity and performance of cattle given palm kernel cake-based diet 20(1988)79
- The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988)259
- Colonization of guinea grass by anaerobic rumen fungi in swamp buffalo and cattle 22(1988)161
- Comparative fiber degradation by mixed rumen fungi from Australian and U.S.A. cattle 23(1989)305
- Distribution of liquid supplement injected into large bales of roughage and comparison of nitrogen supplements for stocker cattle fed low-quality roughages 24(1989)13
- Effect of urea treatment on rumen degradation characteristics of rice straws 24(1989)83
- Effect of soft white wheat addition to alfalfa-grass forage on heifer gain, diet digestibility and in vitro digestion kinetics 24(1989)141

- Utilization of sugar beet molasses and monensin for finishing dairy bullocks 25(1989)11
- Intake, digestion and daily gain by cattle consuming bermudagrass (*Cynodon dactylon*) and supplemented with different combinations of ground corn, vegetable oil, urea and corn gluten and blood meals 25(1989)99
- Cell content**
- Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607
- Cell wall**
- Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607
- Use of processing to explore the factors affecting the digestion of forage cell walls 6(1981)387
- Suitability of cell-wall constituents as predictors of organic matter digestibility in some tropical and subtropical by-products 7(1982)191
- Effect of treatment of wheat straw with ammonia and peracetic acid on digestibility in vitro and cell wall composition 7(1982)325
- Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197
- Electron microscopy of bacteria involved in the digestion of plant cell walls 10(1983/1984)93
- The role of ciliate protozoa and fungi in the rumen digestion of plant cell walls 10(1983/1984)121
- Lignin-poly-saccharide complexes of the plant cell wall and their effect on microbial degradation in the rumen 21(1988)219
- SO₂-treated straw as a silage additive: the participation of soluble and cell wall monosaccharide residues in lucerne silage fermentation 22(1988)255
- Oven and stack ammoniation of grass hays. 1. Changes in chemical composition in relation to digestibility in vitro and cell-wall degradability 24(1989)299
- Rumen ingesta kinetics of cell wall components in dairy cows 25(1989)89
- Cellulase**
- Prediction of the digestibility of ruminant feeds from their solubility in enzyme solutions 2(1977)153
- Enzyme solubility - a method for evaluating the digestibility of alkali-treated straw 2(1977)205
- Prediction of the dry matter digestibility of tropical grasses using a pepsin-cellulase assay 2(1977)247
- Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3(1978)227
- A note on Onozuka 3S cellulase as a replacement for Onozuka SS (P1500) cellulase when estimating forage digestibility in vitro 5(1980)347
- Accuracy of predicting digestibility by the cellulase technique; the effect of pretreatment of forage samples with neutral detergent or acid pepsin 7(1982)83
- Comparison of forage digestibility in vitro with enzymic solubility 11(1984)125
- A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum-pepsin treatment for predicting digestibility and intake of hays in sheep 12(1984/1985)101
- The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14(1986)203
- The use of a cellulase technique to predict digestibility, metabolizable and net energy of forages 19(1988)247
- Effect of sodium silicate on forage digestion with rumen fluid of goats or cellulase using culture solutions adjusted for pH 24(1989)173
- Cellulolysis**
- Manipulation of rumen fluid pH and its influence on cellulolysis in sacco, dry matter degradation and the rumen microflora of sheep offered either hay or concentrate 10(1983/1984)1
- Associative effects of mixed feeds. I. Effects of type and level of supplementation and the influence of the rumen fluid pH on cellulolysis in vivo and dry matter digestion of various roughages 10(1983/1984)15
- Cellulose**
- The digestibility and nutritive value for dairy heifers of rations containing cellulose fibre 2(1977)287
- Some remarks on the determination of acid-detergent fibre in feeds with cetyltrimethylammonium bromide versus determination of detergent fibre with alkylarylsulfonic acid 6(1981)91
- The effects of sulphur supplementation on cellulose digestion in vitro and on nutrient digestion, nitrogen metabolism and rumen characteristics of lambs fed on good quality

- fescue and tropical star grass hays 8(1983)129
- Effect of defaunation on the loss of energy in wethers fed different quantities of cellulose and normal or steamflaked maize starch 16(1986/1987)233
- In-reactor cellulose conversion 21(1988)99
- The physiological effects of cellulose in the human large intestine 23(1989)241
- Cephalosporium eichhorniae***
- Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977)7
- Protein quality studies on rats fed on cassava-root meal enriched with *Cephalosporium eichhorniae* 152 or with conventional plant-protein supplements 6(1981)197
- Cereals, general**
- A comparison of bioavailable energy values of ground cereal grains measured with adult cockerels and growing pigs 19(1988)105
- Chaetomium cellulolyticum***
- Utilization of cattle manure for single-cell protein production with *Chaetomium cellulolyticum* 4(1979)199
- Nutritive value of single-cell protein produced by *Chaetomium cellulolyticum* grown on maize stover and pulp-mill sludge 11(1984)211
- Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. I. Composition and nutritive evaluation 22(1988)1
- Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. II. Effect of method of drying and response to amino acid supplementation 22(1988)13
- Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. III. Nutritive value for chicks and piglets 22(1988)23
- Chaparral**
- Seasonal changes in composition and intake of chaparral by Spanish goats 8(1983)47
- Chenopodium quinoa***
- The quality of the green fraction of leaf protein concentrate from *Chenopodium quinoa* Willd. grown at different levels of fertilizer nitrogen 11(1984)239
- Chicken-broiler**
- Nutritive value of spruce muka (foliage) for the growing chick 3(1978)63
- Utilization of processed pig hair in poultry diets 3(1978)243
- Influence of steam pelleting on the feeding value of tower and candle rapeseed products for broiler chickens 3(1978)251
- Trypsin inhibitor, mineral availability, and performance of broiler chickens fed on diets based on rice bran 4(1979)133
- Feeding value of sunflower and groundnut cakes for broilers 4(1979)143
- House fly larvae meal grown on municipal organic waste as a source of protein in poultry diets 4(1979)227
- Distillery effluents as animal feed: the use of condensed beet molasses stillage (CBMS) in broiler feeding 5(1980)233
- Nitrogen metabolism in broiler chickens consuming the bacterial strain *Alcaligenes eutrophus* 5(1980)241
- Evaluation of brewer's dried grains in the diets of broiler chickens 6(1981)179
- Influence of age on the nutrient utilization of diets for broilers 6(1981)405
- Influence of increasing levels of β -glucanase on the productive value of barley diets for broiler chickens 7(1982)351
- Nutritional value of waxy barleys for broiler chickens 8(1983)25
- The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303
- A note on the nutritive value of dried instant coffee residue for broiler chickens and turkey poults 10(1983/1984)285
- Substitution of finger millet (*Eleusine coracana*) and bulrush millet (*Pennisetum typhoides*) for maize in broiler feeds 10(1983/1984)291
- Meat and bone and fish meals in balancer feeds for choice-fed broilers 11(1984)221
- The proximate composition and nutritive value of the winged bean *Psophocarpus tetragonolobus* (L.) DC for broilers 11(1984)231
- The utilization by broiler chickens of poultry offal hydrolysed in formic acid 11(1984)247
- The effect of pelleting on the utilisation of yeast grown on n-alkanes by broiler chicks 12(1984/1985)285
- Influence of supplemental fat and steam pelleting on the response of broilers to diets varying in energy density 13(1985)57
- The effect of β -glucanase on the utilization of starch and nitrogen by broiler chickens fed on barley of low- or high-viscosity 15(1986)83
- Effect of dietary ascorbic acid (vitamin C) supplementation on the performance of broiler chickens in a tropical environment 16(1986/1987)17

- Influence of nutrient density on feed consumption, weight gain and gut capacity of broilers, Leghorns and turkeys reared to 26 days of age 16(1986/1987)129
- Palm kernel meal as a feed for poultry. 3. Replacement of groundnut cake by palm kernel meal in broiler diets 16(1986/1987)195
- Nutritive value of irradiated and β -glucanase-treated wild oat groats (*Avena fatua* L.) for broiler chickens 16(1986/1987)243
- The effect of adding white fish meal containing enzyme digested or untreated stickwater solids to diets for broiler chickens 16(1986/1987)253
- Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Proteen 17(1987)1
- Oil palm slurry (OPS) as a partial replacement for maize in the diets of broiler chickens 17(1987)157
- The energy content of full-fat soya beans in meal and pelleted feeds for adult cocks and broilers 19(1988)97
- Feeding and nutritive value of hominy and corn grits for poultry 19(1988)313
- The metabolisable energy (AME_m) of fats for broilers in relation to their chemical composition 20(1988)45
- Treatment of low-tannin sorghum grain for broiler feed 20(1988)69
- Utilization of aquatic plants *Elodea canadensis* and *Hydrilla verticillata* in broiler chick diets 20(1988)155
- Effects of enzyme supplementation of diets based on wheat, rye or triticale on their productive value for broiler chickens 20(1988)313
- The nutritional value for broilers of fats characterized by short-chain fatty acids as affected by level of inclusion and age 22(1988)105
- Nutritional value of triticale (Carman) for broiler diets 22(1988)119
- Effects of wood ash extract treatment on the feeding value and utilization of high-tannin sorghums by broiler chicks 22(1988)131
- The nutritive value of dried coffee pulp (DCP) in broiler chickens's diets 22(1988)139
- The effect of fat level and type on the utilization of triticale (cultivar Carman) by broiler chicks 22(1988)273
- The nutritive value of tamarind seeds for broiler chicks 22(1988)285
- Effects of oven-dried cassava peel meal on the performance and carcass characteristics of broiler chickens 24(1989)247
- The effect of β -glucanase supplementation on nutrient digestibility and growth in broilers given diets containing barley, oat groats or wheat 25(1989)193
- Chicken-chick**
- Amino acid supplementation of methanol-grown dried microbial cells in diets for young chicks 4(1979)63
- Nutritive value of leaf-protein concentrate from potato haulm for rats and chicks 6(1981)413
- Growth performance of, and mimosine excretion by, young chicks fed on *Leucaena leucocephala* 7(1982)247
- Metabolisable energy value of fats in chicks and adult cockerels 7(1982)365
- Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983)99
- The effect of nutrient supplementation on the feeding value of kapok seed for poultry 9(1983)263
- A comparative study of metabolisable energy in ducklings and domestic chicks 11(1984)199
- A comparison of metabolizable energy content of eight common feed ingredients determined with young guinea fowls (keets) and pullet chicks 15(1986)1
- Palm kernel meal as a feed for poultry. 1. Composition of palm kernel meal and availability of its amino acids to chicks 16(1986/1987)179
- Palm kernel meal as a feed for poultry. 2. Diets containing palm kernel meal for starter and grower pullets 16(1986/1987)187
- Utilization of tannin-containing and tannin-free faba beans (*Vicia faba*) by young chicks: effects of pelleting feeds on energy, protein and starch digestibility 20(1988)59
- Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. III. Nutritive value for chicks and piglets 22(1988)23
- Canavanine content and toxicity of sesbania leaf meal for growing chicks 25(1989)137
- The effect of kernels of karanja (*Pongamia glabra* Vent) on growth and feed efficiency in broiler chicks to 4 weeks of age 25(1989)201

Chicken-cockerel

Metabolisable energy value of fats in chicks and adult cockerels 7(1982)365

Comparison of bioavailable energy values of diets measured with poultry and swine 8 (1983)191

Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983) 99

Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin) 11(1984)293

Feeding value of hydroponically sprouted barley for poultry and pigs 13(1985)183

A comparison of bioavailable energy values of ground cereal grains measured with adult cockerels and growing pigs 19(1988)105

Effect of enzyme supplementation on the nutritive value of hullless barley in chicken diets 24(1989)233

Chicken-general

The potato (*Solanum tuberosum*) as a source of nutrients for pigs, calves and fowl - a review 2(1977)171

A comparison of three fibre methods for predicting the metabolizable energy content of sorghum grain for poultry 2(1977)197

Estimation of lysine in compounded poultry diets after enzymic digestion 3(1978)57

Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73

Rapeseed meal and its use in poultry diets. A review 5(1980)255

Proximate analysis, amino acid composition and tannin content of improved Nigerian sorghum varieties and their potential in poultry feeds 7(1982)359

Chemical composition and nutritional value of para-rubber seed and its products for chickens 10(1983/1984)257

A new biological method for determining amino acid digestibility in poultry feed-stuffs using a simple cannula, and the influence of dietary fibre on endogenous amino acid output 12(1984/1985)29

Nutritional value for chickens of dried distillers-spent-grain from barley and dehulled barley 17(1987)145

Nutritional value of pigeonpea (*Cajanus cajan*) meal in poultry diets 25(1989)123

Chicken-laying hen

Feeding value of sunflower- and groundnut-cakes for laying hens 6(1981)63

Effects of the addition of silica to the diet of the laying hen, with particular reference to egg-shell quality 6(1981)157

Evaluation of brewer's dried grains in the diets of laying hens 6(1981)169

Rapeseed meal tannins and egg taint 6(1981) 421

The utilization of torula yeast in diets for White Leghorn birds during growth and laying periods 7(1982)185

The involvement of trimethylamine oxide in fish meal in the production of egg taint 8 (1983)119

Influence of nutrient density on feed consumption, weight gain and gut capacity of broilers, Leghorns and turkeys reared to 26 days of age 16(1986/1987)129

The influence of oat groats and dietary level of lysine on the laying performance of cross-bred hens 17(1987)271

Evaluation of triticale given to pigs, poultry and rats 19(1988)359

Palm kernel meal as a feed for poultry. 4. Use of palm kernel meal by laying birds 20 (1988)279

Dry matter yield and chemical composition of pigeon pea (*C. cajan*) leaf meal and the nutritive value of pigeon pea leaf meal and grain meal for laying hens 24(1989)111

Chopping

Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227

Chunies

Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315

Citrus pulp

Composition of citrus pulp 5(1980)1

Citrus pulp in diets for fattening lambs 5 (1980)11

Citrus pulp in diets for fattening rabbits 5 (1980)23

Clover

Losses of beta-carotene in red clover in an acid medium during ensiling 4(1979)81

The enzymic nature of the degradation of beta-carotene in red clover and in other forage crops during silage-making with acid additives 5(1980)59

Degradation in vitro and in sacco of red clover leaves and stems 17(1987)295

Cobalt

Treatment of dairy sheep with soluble glass boluses containing copper, cobalt and selenium 19(1988)79

The effect of soluble-glass boluses containing copper, cobalt and selenium on the blood composition of ewes 21(1988)23

Cocoa-pod

Studies on the feeding value of agro-industrial by-products: strategies for improving the utilisation of cocoa-pod-based diets by ruminants 20(1988)189

Coffee pulp

Changes in nutrient intake and performance by goats fed coffee pulp-based diets followed by a commercial concentrate 14(1986)1

The nutritional value of dried coffee pulp (DCP) in broiler chickens's diets 22(1988)139

Coffee residue

A note on the nutritive value of dried instant coffee residue for broiler chickens and turkey poults 10(1983/1984)285

Colour value

Influence of pericarp and endosperm colour and type on digestibility of grain sorghum by pigs 2(1977)219

Relationships between tristimulus colour values and digestibility of grain sorghum by swine 2(1977)327

Compound feeds

Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3(1978)227

Some factors affecting the future composition of UK compound animal feeds 6(1981)1

Energy evaluation of feeds and compounded diets for pigs - a review 7(1982)387

The prediction of the net energy value of compound feedstuffs for dairy cattle 11(1984)137

The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14(1986)203

The prediction of the energy value of compounded pig foods from chemical analysis 17(1987)81

Cooking

Effect of sodium hydroxide treatment and/or extrusion cooking on the nutritive value of peanut hulls 12(1984/1985)187

Copper

Reducing the potential copper toxicity of concentrates to sheep by the use of molybdenum and sulphur supplements 2(1977)235

Utilization by sheep of dried pig faeces containing a high concentration of copper 3(1978)31

Olaquinox as performance-promoting feed additive for growing pigs 4(1979)117

Treatment of dairy sheep with soluble glass boluses containing copper, cobalt and selenium 19(1988)79

The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171

The effect of soluble-glass boluses containing copper, cobalt and selenium on the blood composition of ewes 21(1988)23

Copper and iodine in pig diets with high glucosinolate rapeseed meal. I. Performance and thyroid hormone status of growing pigs fed on a diet with rapeseed meal treated with copper sulphate solution or untreated and supplements of iodine, copper or a quinoxaline derivative 22(1988)33

Copper and iodine in pig diets with high glucosinolate rapeseed meal. II. Influence of iodine supplements for rations with rapeseed meal untreated or treated with copper ions on performance and thyroid hormone status of growing pigs 22(1988)45

Corn gluten feed

Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375

Response of high-producing dairy cows to high levels of dried corn gluten feed 24(1989)191

Cotton

Effect of ammonia treatment and processing of whole cotton plants as sole source of roughage in complete feeds for growing cross-bred calves 13(1985)93

Cotton straw

Effects of ammonia treatment and processing of cotton straw as the sole source of roughage in complete feeds on rumen characteristics with Murrah buffaloes 14(1986)95

Cottonseed

A review of the nutritive value and utilization of whole cottonseed, cottonseed meal and associated by-products by dairy cattle 18(1987)89

Cottonseed cake

Comparison of cottonseed cake and dried blood as nitrogen sources for growing Sudan Nubian goats 22(1988)333

Cottonseed meal

Effects of supplementing a wheat straw diet with urea or a urea-molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25

Nutrient composition of Greek cottonseed meal 18(1987)295

Evaluation of Greek cottonseed meal. 2. Use in rations for fattening pigs 18(1987)303

Chemical and biological evaluation of Nigerian cottonseed meal protein quality 20(1988)251

Cowpea

Variation in metabolizable energy content of raw or autoclaved white and brown varieties of three tropical grain legumes 13(1985)141

Crop residues

Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter disappearance in sacco 17(1987)179

Culinary wastes

Chemical composition of culinary wastes and their potential as a feed for ruminants 5(1980)205

Cyanide

The influence of processing and storage on hydrogen cyanide and tannin contents of para-rubber seed and its products 9(1983)319

Effect of cassava-based diets varying in cyanide content on the performance and physiopathology of the African giant rat (*Cricetomys gambianus* Waterhouse) 11(1984)1

Effect of variety and plant age on the cyanide content of whole-root cassava chips and its reduction by sun-drying 11(1984)57

Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227

Cystine

Meat meal in the diet of the early-weaned pig. II. Amino acid supplementation 4(1979)43

D**Defaunation**

Effect of defaunation on straw digestion in sacco and on particle retention in the rumen 10(1983/1984)165

Effect of defaunation on the loss of energy in wethers fed different quantities of cellulose and normal or steamflaked maize starch 16(1986/1987)233

Effect of defaunating the rumen 21(1988)229

Deoxynivalenol

Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin) 11(1984)293

Deterioration

A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3(1978)89

Losses of beta-carotene in red clover in an acid medium during ensiling 4(1979)81

Chemicals for preventing moulding in damp hay 6(1981)323

Losses of beta-carotene in unwilted forage crops during silage-making and feeding 9(1983)63

A comparison between degradation in vitro and in sacco of constituents of untreated and ammonia-treated barley straw 10(1983/1984)199

The effect of anhydrous ammonia and moisture content on the preservation and chemical composition of perennial ryegrass hay 11(1984)159

Effect of dietary nitrogen source on microbial protein synthesis, dietary protein degradation and nutrient digestion in steers 14(1986)117

Nutrient losses during deterioration of hay in relation to changes in biochemical composition and microbial growth 15(1986)149

Estimating yield and nutrient losses due to rainfall on field-drying alfalfa hay 16(1986/1987)7

Effect of formaldehyde treatment on the degradation of acid-preserved fish silage protein in vitro 16(1986/1987)297

Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro degradability during maturation 17(1987)15

Degradation in vitro and in sacco of red clover leaves and stems 17(1987)295

Microbial dynamics in barley grain stored under controlled atmosphere 19(1988)299

Fungal contamination of hydroponic forage 20(1988)163

Deterioration

The enzymic nature of the degradation of beta-carotene in red clover and in other forage crops during silage-making with acid additives 5(1980)59

Dicoumarol

Dicoumarol concentrations in sweetclover hay treated with preservatives and in spoiled hay of high- and low-coumarin cultivars of sweetclover 14(1986)221

Digesta particle size

The determination of digesta particle size in some herbivores 7(1982)35

Digesta retention

Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279

Digestibility-Agrocyron elongatum

Effect of lactate on the in vitro digestion of *Agrocyron elongatum* by rumen microorganisms 16(1986/1987)161

Digestibility-alfalfa

Composition and fibre digestion in morphological components of an alfalfa-timothy sward 19(1988)135

Digestibility-amino acids and proteins

Apparent ileal and faecal digestibility of amino acids in barley-based diets supplemented with soya bean meal or canola meal for growing pigs 14(1986)183

Digestibility-animal waste

Digestibility by cattle of diets containing dried pig faeces 3(1978)155

Digestibility of wet cage layer excreta and an excreta-supplemented diet by cattle as determined by two methods 5(1980)77

Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 17(1987)235

Digestibility-barley

The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Effect of storage and processing of barley on its utilization by pigs and rats 9(1983)89

The digestion by sheep of barley and maize-based diets containing either meat and bone meal or soya bean meal 15(1986)301

Digestibility-barley meal

The influence of feeding level and level of dietary inclusion on the digestibility of barley meal in the pig 2(1977)41

Digestibility-bermudagrass

Intake, digestion and daily gain by cattle consuming bermudagrass (*Cynodon dactylon*) and supplemented with different combinations of ground corn, vegetable oil, urea and corn gluten and blood meals 25(1989)99

Digestibility-canola meal

Factors affecting the digestibility by pigs of energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253

Digestibility-cassava

Net energy for gain (NE_{gain}) of cassava tops 20(1988)97

Digestibility-cell contents

Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197

Digestibility-cell wall

Use of processing to explore the factors affecting the digestion of forage cell walls 6(1981)387

Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197

Digestibility-compound feeds

The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14(1986)203

Digestibility-crop residues

Improving utilization of crop residues. Evaluation of two chemicals used to improve digestibility of crop residues 9(1983)205

Digestibility-desert grass

Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by goats 17(1987)219

Digestibility-dry matter

The accuracy of the pepsin-cellulase technique for estimating the dry matter digestibility in vivo of grasses and legumes 3(1978)277

Bias when predicting crude protein, dry matter digestibility and voluntary intake of tropical grasses by near-infrared reflectance 9(1983)221

Variations and prevision of the in sacco dry matter digestion of concentrates and by-products 13(1985)7

Digestibility-faba bean

Utilization of tannin-containing and tannin-free faba beans (*Vicia faba*) by young chicks: effects of pelleting feeds on energy, protein and starch digestibility 20(1988)59

Digestibility-fibre

The effects of different sources of nitrogen supplementation on the digestion of fibre components in the rumen of steers 17(1987)65

Approach to the energetic importance of fibre digestion in pigs. I. Importance of fermentation in the overall energy supply 23(1989)141

Approach to the energetic importance of fibre digestion in pigs. II. An experimental approach to hindgut digestion 23(1989)169

Digestion and fermentation pattern of various dietary fiber sources in the rat 23(1989)195

Digestibility-fish meal

Fish protein concentrates for pre-ruminant calves: effects of processing temperatures and source of fish on protein digestibility and biological value 18(1987)181

Digestibility-fodder beet

The chemical composition, digestibility and energy value of fodder-beet roots 18(1987)225

Digestibility-forage, general

Some remarks on the analytical procedure of Van Soest for the prediction of forage digestibility 3(1978)309

Variability in structural carbohydrates and in vitro digestibility of forages. 4. Fenugreek (*Trigonella foenum-graecum*) 4(1979)17

A note on Onozuka 3S cellulase as a replacement for Onozuka SS (P1500) cellulase when estimating forage digestibility in vitro 5(1980)347

Effect of high-pressure steam treatment on chemical composition and digestibility in vitro of roughages 7(1982)61

Accuracy of predicting digestibility by the cellulase technique; the effect of pretreatment of forage samples with neutral detergent or acid pepsin 7(1982)83

Relative intakes, digestibility and bacterial protein synthesis by sheep and cattle fed high-roughage diets 16(1986/1987)75

Prediction of forage digestibility by principal component analysis of near infrared reflectance spectra 16(1986/1987)215

Digestibility-forages, general

Comparison of laboratory methods for predicting the organic matter digestibility of forages 2(1977)337

Digestibility-general

Prediction of the digestibility of ruminant feeds from their solubility in enzyme solutions 2(1977)153

Comparative digestibility of feedstuffs by sheep and cows 12(1984/1985)47

Digestibility-grapevine

Digestibility in vivo of ensiled grapevines (branches and leaves): influence of the system of analysis in the detergent fibre scheme on the prediction of digestibility 22(1988)173

Digestibility-grass

Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607

Prediction of the dry matter digestibility of tropical grasses using a pepsin-cellulase assay 2(1977)247

Comparison of forage digestibility in vitro with enzymic solubility 11(1984)125

Effect of sodium silicate on forage digestion with rumen fluid of goats or cellulase using culture solutions adjusted for pH 24(1989)173

Digestibility-groundnut

The effect of grinding groundnut hulls either with or without alkali treatment on digestibility of diet and on ruminal and blood components 15(1986)69

Digestibility-hay

Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607

Differences between cattle and sheep in their digestion and relative intake of a mature tropical grass hay 3(1978)41

Estimation of the digestibility of low-quality hays by cattle from measurements made with sheep 3(1978)51

The effect of intake and hay: concentrate ratio upon digestibility and digesta passage 11(1984)167

Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279

The voluntary intake and digestibility of combinations of cereal crop residues and legume hay for sheep 12(1984/1985)241

Chemical composition and digestion in vitro of orchardgrass hay ammoniated by different techniques 17(1987)109

The effect of grinding and pelleting hay on digestibility, fermentation rate, digesta passage and rumen and faecal particle size in cows 19(1988)145

Oven and stack ammoniation of grass hays. 1. Changes in chemical composition in relation to digestibility in vitro and cell-wall degradability 24(1989)299

Oven and stack ammoniation of grass hays.

2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313

Effects of urea treatment on chemical composition and in vitro digestibility on meadow hays of Northern Portugal 25(1989)157

Digestibility-jackbean

Ripe beans of *Canavalia ensiformis* (jackbean) as feed ingredient for monogastric animals 20(1988)325

Digestibility-legumes

Comparison of forage digestibility in vitro with enzymic solubility 11(1984)125

Digestibility-Leucaena leucocephala

The intake and digestibility in goats of *Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989)45

Digestibility-lupin

Digestibility and energy value of sweet lupin seed (*Lupinus albus* var. multolupa) in pigs 12(1984/1985)171

Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammer-milled lupin grain in oat hay or oat straw based diets 16(1986/1987)143

Digestibility-maize

Rate of growth, feed utilisation and digestibility observed with diets containing ensiled maize grain or maize ears fed alone or in mixtures 1(1976)401

The digestion and metabolism of the energy and protein of maize 1(1976)455

Measurement and prediction of digestibility of forage maize in The Netherlands 10(1983/1984)301

Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91

High temperature and pressure processing of maize cobs. I. Digestibility in vitro of processed cobs 12(1984/1985)151

The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99

Digestibility-manioc

The digestion of diets containing manioc by young growing pigs 12(1984/1985)119

Digestibility-mixed feeds

Associative effects of mixed feeds. I. Effects of type and level of supplementation and the influence of the rumen fluid pH on cellulolysis in vivo and dry matter digestion of various roughages 10(1983/1984)15

Associative effects of mixed feeds. II. The effect of dietary addition of bicarbonate salts on the voluntary intake and digestibility of diets containing various proportions of hay and barley 10(1983/1984)31

Digestibility-molasse

Differences in chemical composition and digestibility of beet and cane molasses 13(1985)83

Digestibility-nitrogen

The apparent and true ileal digestibility of nitrogen and amino acids in lactic casein for the growing pig 16(1986/1987)169

Digestibility-olive cake

Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139

Digestibility-organic matter

Suitability of cell-wall constituents as predictors of organic matter digestibility in some tropical and subtropical by-products 7(1982)191

The effect of high levels of molasses in combinations with hay on digestibility of organic matter, microbial protein synthesis and volatile fatty acid production in vitro 7(1982)277

A comparison between fermentation heat of forages and organic matter digestibility determined by in vitro incubation with rumen fluid 23(1989)333

Digestibility-paper diets

Intake, digestibility and energy utilization of corrugated paper diets 17(1987)57

Digestibility-pasture plants

The relationship between the intake of herbage grazed by dairy cows and its digestibility 1(1976)53

The analytical and biological accuracy of estimating the dry matter digestibility of different legume species 1(1976)61

Solubility of the protein of tropical pasture species and the rate of its digestion in the rumen 5(1980)183

Digestibility-pea

Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal tract of pigs 17(1987)33

Digestibility-plantain sheath

In vitro studies and short-term feeding trial in lambs to evaluate plantain sheath (*Musa sapientum*) as a feed for ruminants 20 (1988)343

Digestibility-potato

The effect of different patterns of thermal processing of potatoes on their digestibility by growing pigs 4(1979)295

Digestibility-prediction

Comparison of methods for predicting digestibility of feeds 20(1988)203

Digestibility-protected fat

The effects on feed intake, growth and digestibility of nutrients of including calcium soaps of fat in diets for young calves 14 (1986)103

Digestibility-proteins and amino acids

Digestibility of protein and amino acids of a fermentation single-cell protein for veal calves 2(1977)377

The apparent and true ileal digestibility of nitrogen and amino acids in lactic casein for the growing pig 16(1986/1987)169

Digestibility-rapeseed

Degradation of crude protein and quality of undegradable protein in untreated or formaldehyde-treated rapeseed meal 12 (1984/1985)19

Effect of graded levels of raw and processed rapeseed on feed digestibility and nutrient utilization in young pigs 24(1989)275

Digestibility-rapeseed meal

The digestion in the small intestine of young bulls of the protein of rapeseed meal treated or untreated with formaldehyde 7 (1982)225

Digestibility-ryegrass

The effect of urea on the preservation and digestibility in vitro of perennial ryegrass 10(1983/1984)49

Digestibility-silage

Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279

The effect of supplementing grass silage with sodium hydroxide-treated chopped or milled straw on intake and digestibility by wether sheep 12(1984/1985)215

The intake and digestion of maize silage-based diets by dairy cows and sheep 20 (1988)299

The digestibility and metabolisable energy content of grass silage and their prediction from laboratory measurements 24(1989)27

Digestibility-sisal waste

Feed value of Tanzanian sisal wastes: chemical composition and digestibility in vitro 14 (1986)255

Digestibility-sorghum grain

Influence of pericarp and endosperm colour and type on digestibility of grain sorghum by pigs 2(1977)219

Relationships between tristimulus colour values and digestibility of grain sorghum by swine 2(1977)327

Digestibility-soya bean

Effects of micronization on the digestibility of whole soya beans and rapeseeds for the growing pig 3(1978)179

Effect of formaldehyde-treated soya bean meal and urea in starters on nitrogen quality, degradability in sacco, sheep digestibility and calf performance 16(1986/1987)287

Digestibility-straw

The effect on digestibility of a new technique for alkali treatment of straw 1(1976)73

Effect on digestibility and nitrogen content of barley straw of different ammonia treatments 2(1977)131

Enzyme solubility - a method for evaluating the digestibility of alkali-treated straw 2 (1977)205

Ensiled alkali-treated straw. I. Effect of level and type of alkali on the composition and digestibility in vitro of ensiled barley straw 3(1978)117

Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381

Titrateable groups and soluble phenolic compounds as indicators of the digestibility of chemically treated roughages 6(1981)123

Studies on ammonia-treated straw. I. The effects of type and level of ammonia, moisture content and treatment time on the digestibility in vitro and enzyme soluble organic matter of oat straw 7(1982)45

Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71

The effect of calcium hydroxide treatment on the digestibility of chopped oat straw supplemented with a solution containing urea, calcium, phosphorus, sodium, trace elements and vitamins 7(1982)93

Intake and digestibility in sheep fed on alkali-treated straw and concentrates, as complete diets or separate feeds 7(1982)271

Effect of treatment of wheat straw with ammonia and peracetic acid on digestibility in vitro and cell wall composition 7(1982)325

Digestibility of diets containing increasing levels of NaOH-treated or untreated wheat straw 7(1982)331

The effect of prolonged storage on the digestibility and nitrogen content of ammonia-treated barley straw 8(1983)147

Composition and enzymatic digestibility of Oregon grass straws 9(1983)283

Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65

Effect of defaunation on straw digestion in sacco and on particle retention in the rumen 10(1983/1984)165

Digestibility studies on ammonia-treated straw 10(1983/1984)213

The digestion of untreated and ammonia-treated barley straw in an artificial rumen 10(1983/1984)229

Ammonia treatment of straw via the hydrolysis of urea. II. Additions of soya bean (urease), sodium hydroxide and molasses; effects on the digestibility of urea-treated straw 11(1984)115

The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149

The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12(1984/1985)141

Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81

The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99

The effect of dietary barley on carbohydrate digestibility of sulphur dioxide-treated wheat straw by sheep 18(1987)55

Digestibility-sugar beet

Dried sugar beet pulp as a high energy feed for beef cattle 1(1976)643

Digestion of diets containing molassed or plain sugar-beet pulp by growing pigs 23(1989)67

Digestibility-tallow

Digestibility of a fully hydrogenated tallow for growing pigs 5(1980)87

Digestibility-Tilley and Terry test

Improving the reliability of the two-stage in vitro assay for ruminant feed digestibility by calibration against in vivo data from a wide range of sources 18(1987)233

Digestibility-timothy

Composition and fibre digestion in morphological components of an alfalfa-timothy sward 19(1988)135

Digestibility-wheat

The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Effect of soft white wheat addition to alfalfa-grass forage on heifer gain, diet digestibility and in vitro digestion kinetics 24(1989)141

Drechslera sorokiniana

Effects of *Drechslera sorokiniana* infection on the yield and quality of tall fescue forage 15(1986)41

Dried bakery product

Variation in nutrient content of samples of dried bakery product 7(1982)419

Dried microbial cells

The replacement of skim milk by dried microbial cells and whey in milk diets for calves 2(1977)85

Limiting amino acids in dried microbial cells given to young turkeys 4(1979)209

Drying methods

Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381

Influence of the drying process on the composition of brewer's dried grains 6(1981)163

Effect of variety and plant age on the cyanide content of whole-root cassava chips and its reduction by sun-drying 11(1984)57

Microwave drying of alfalfa compared to field- and oven-drying: effects on forage quality 16(1986/1987)1

Estimating yield and nutrient losses due to rainfall on field-drying alfalfa hay 16(1986/1987)7

Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227

The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20(1988)335

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. II. Effect of method of drying and response to amino acid supplementation 22(1988)13

Drying treatment drastically affects feed evaluation and feed quality with certain tropical forage species 22(1988)69

Duck

A comparative study of metabolisable energy in ducklings and domestic chicks 11(1984) 199

Dye absorption test

The dye absorption test as an indicator of the nutritive quality of heat-processed navy bean 9(1983)71

E**Economics**

Economic aspects of high-plane feeding of bulls with maize silage 1(1976)513

Energetics of stockfeed production 3(1978)1

Assessment of the production costs of fish protein hydrolysates 7(1982)147

Editorial-general

The dilemma of animal feeds and nutrition 1 (1976)1

Egg taint

Rapeseed meal tannins and egg taint 6(1981) 421

The involvement of trimethylamine oxide in fish meal in the production of egg taint 8 (1983)119

Elodea

Utilization of aquatic plants *Elodea canadensis* and *Hydrilla verticillata* in broiler chick diets 20(1988)155

Energy-metabolizable

A comparative study of metabolisable energy in ducklings and domestic chicks 11(1984) 199

Effect of prolonged wilting in poor conditions on the fermentation quality, metabolisability and net energy value of silage given to sheep 12(1984/1985)109

Variation in metabolizable energy content of raw or autoclaved white and brown varieties of three tropical grain legumes 13 (1985)141

The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14 (1986)203

A comparison of metabolizable energy content of eight common feed ingredients determined with young guinea fowls (keets) and pullet chicks 15(1986)1

The metabolisable energy (AME_n) of fats for broilers in relation to their chemical composition 20(1988)45

The digestibility and metabolisable energy content of grass silage and their prediction from laboratory measurements 24(1989)27

Energy-net

The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14 (1986)203

The use of a cellulase technique to predict digestibility, metabolizable and net energy of forages 19(1988)247

Net energy for gain (NE_{gain}) of cassava tops 20(1988)97

Energy balance

Energy and nitrogen balance, intake and growth rate of lambs fed on unwilted grass silages treated with a formaldehyde-acid mixture or untreated 9(1983)107

Dietary fiber and energy balance: integration of the human ileostomy and in vitro fermentation models 23(1989)261

Energy density

Influence of supplemental fat and steam pelleting on the response of broilers to diets varying in energy density 13(1985)57

Energy intake

Dry matter, protein, energy and fibre intake by dairy heifers grazing a Rhodes grass (*Chloris gayana*) pasture 6(1981)15

Energy metabolism

Energy metabolism in lambs fed silage diets with sucrose or a growth promoter 16 (1986/1987)203

Effect of defaunation on the loss of energy in wethers fed different quantities of cellulose and normal or steamflaked maize starch 16 (1986/1987)233

Energy systems

The energy systems used in the EEC countries: their application to 24 commercial pig-feeds and relationships between them 6(1981)211

A flexible, two-component feeding system derived from Blaxter's three-component, metabolisable-energy system 6(1981)223

Prediction of the energy value of commercial pig feeds, according to the various energy systems used in the European Community, that are based on simple analytical parameters of the feed 6(1981)367

Energy utilisation

Protein and energy utilisation in rats of diets based on lupin seed (*Lupinus albus* var. multolupa) 15(1986)33

Intake, digestibility and energy utilization of corrugated paper diets 17(1987)57

Energy value

- The digestion and metabolism of the energy and protein of maize 1(1976)455
- Dried sugar beet pulp as a high energy feed for beef cattle 1(1976)643
- A comparison of three fibre methods for predicting the metabolizable energy content of sorghum grain for poultry 2(1977)197
- Waxy versus regular maize: energy value for growing pigs and utilization in piglet diets 4(1979)109
- Protein and energy value of vinasse for pigs 5(1980)221
- Predicting the energetic value of forage from digestibility of organic matter in vitro or solubility in potassium hydroxide 6(1981)235
- Metabolizable energy value of fats in chicks and adult cockerels 7(1982)365
- Energy evaluation of feeds and compounded diets for pigs - a review 7(1982)387
- Comparison of bioavailable energy values of diets measured with poultry and swine 8(1983)191
- Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983)99
- The prediction of the net energy value of compound feedstuffs for dairy cattle 11(1984)137
- Digestibility and energy value of sweet lupin seed (*Lupinus albus* var. multolupa) in pigs 12(1984/1985)171
- Utilization of whole rape (Canola) seed and sunflower seeds as sources of energy and protein in calf starter diets 15(1986)101
- The prediction of the energy value of compounded pig foods from chemical analysis 17(1987)81
- Comparative bioavailable energy values of five ground cereal grains measured with growing rats and pigs 18(1987)143
- The chemical composition, digestibility and energy value of fodder-beet roots 18(1987)225
- The energy content of full-fat soya beans in meal and pelleted feeds for adult cocks and broilers 19(1988)97
- A comparison of bioavailable energy values of ground cereal grains measured with adult cockerels and growing pigs 19(1988)105
- The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173

Approach to the energetic importance of fibre digestion in pigs. I. Importance of fermentation in the overall energy supply 23(1989)141

Approach to the energetic importance of fibre digestion in pigs. II. An experimental approach to hindgut digestion 23(1989)169

Enzymes, general

The enzymic nature of the degradation of beta-carotene in red clover and in other forage crops during silage-making with acid additives 5(1980)59

Use of enzymes for improvement of feed 6(1981)105

Composition and enzymatic digestibility of Oregon grass straws 9(1983)283

The effect of enzymatic treatment on amino acid content and nitrogen characteristics of feather meal 16(1986/1987)151

Factors affecting the formation of polysaccharide-degrading enzymes by rumen micro-organisms 21(1988)191

Effect of enzyme supplementation on the nutritive value of hullless barley in chicken diets 24(1989)233

Ethanol

Production of ethanol from lignocellulosic materials 21(1988)175

Ethanolamine

Effects of ethanolamine and ethylenediamine treatments on the composition and degradability of wheat straw 13(1985)131

Ethylenediamine

Effects of ethanolamine and ethylenediamine treatments on the composition and degradability of wheat straw 13(1985)131

F

Faba bean

Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261

Utilization of tannin-containing and tannin-free faba beans (*Vicia faba*) by young chicks: effects of pelleting feeds on energy, protein and starch digestibility 20(1988)59

Faecal particle size

The effect of grinding and pelleting hay on digestibility, fermentation rate, digesta passage and rumen and faecal particle size in cows 19(1988)145

Fats

Poly-unsaturated fat in young ruminants 2 (1977)93

Digestibility of a fully hydrogenated tallow for growing pigs 5(1980)87

Metabolisable energy value of fats in chicks and adult cockerels 7(1982)365

Variability for protein and oil quality in *Lupinus albus* 9(1983)271

The effect of level and source of protein, fibre and fat in the diet on the energy value of compounded pig feeds 11(1984)11

Effects of pelleting, added sodium bentonite and fat in the diet on the energy value of performance and carcass characteristics of small white turkeys 12(1984/1985)223

Effect of processing on the utilization by young pigs of the fat in soya beans and sunflower seeds 12(1984/1985)267

Influence of supplemental fat and steam pelleting on the response of broilers to diets varying in energy density 13(1985)57

Performance and carcass characteristics of Sudan desert sheep and goats on high roughage diets with added fat 13(1985)147

Effect of dietary protein and fat levels on the utilization of the fat in sunflower seeds by the young pig 13(1985)159

Thermofractographic determination of monocarbonyl compounds in animal waste fats used as feed fat 13(1985)227

Fat supplementation of diets for growing rabbits 16(1986/1987)109

Influence of nutrient density on feed consumption, weight gain and gut capacity of broilers, Leghorns and turkeys reared to 26 days of age 16(1986/1987)129

High-fat maize in diets for pigs and sows 17 (1987)201

Isoacids in ruminant nutrition: their role in ruminal and intermediary metabolism and possible influences on performances - a review 18(1987)169

Investigations on the nutritional value for ruminants of various fats 19(1988)111

Use of fat in diets of sheep in hot environments. I. Effects on performance, carcass characteristics and lipid composition of plasma 19(1988)327

Use of fat in diets of sheep in hot environments. II. Effect on rumen metabolism 19 (1988)343

The metabolisable energy (AME_n) of fats for broilers in relation to their chemical composition 20(1988)45

Feed intake and milk yield at increasing supplement of a palmitic and stearic acid-rich type of fat in comparison with animal fat 22(1988)179

Feed intake, milk yield and milk composition by replacing unprotected fat by Ca-soaps for dairy cows 22(1988)193

The effect of fat level and type on the utilization of triticale (cultivar Carman) by broiler chicks 22(1988)273

Fatty acids

Absorption of short-chain fatty acids and their role in the hindgut of monogastric animals 23(1989)43

Feather meal

Protein quality of mixtures of animal protein meals 9(1983)79

Effects of processing time and moisture content on amino acid composition and nitrogen characteristics of feather meal 14(1986)279

The effect of enzymatic treatment on amino acid content and nitrogen characteristics of feather meal 16(1986/1987)151

Feeding standards

Who needs feeding standards? 6(1981)101

Feeding standards are necessary 7(1982)423

Fenugreek

Variability in structural carbohydrates and in vitro digestibility of forages. 4. Fenugreek (*Trigonella foenum-graecum*) 4(1979)17

Fermentation-animal waste

Fermentation characteristics and feeding value of ensiled poultry litter containing wheat straw, bagasse or sawdust 7(1982)341

Slaughterhouse by-products preserved by *Lactobacillus plantarum* fermentation as feed for mink and foxes 20(1988)287

Fermentation-colon

In vitro fermentation of dietary fiber by human fecal organisms 23(1989)121

Approach to the energetic importance of fibre digestion in pigs. I. Importance of fermentation in the overall energy supply 23(1989)141

Digestion and fermentation pattern of various dietary fiber sources in the rat 23(1989)195

The rumen as a model of the microbiology of fibre digestion in the human intestine 23 (1989)227

Fermentation-forage

A comparison between fermentation heat of forages and organic matter digestibility determined by in vitro incubation with rumen fluid 23(1989)333

Fermentation-fresh crops

The chemical composition and nutritive value for ruminants of four possible energy crops and their residues of anaerobic fermentation 15(1986)197

Fermentation-illum

Dietary fiber and energy balance: integration of the human ileostomy and in vitro fermentation models 23(1989)261

Fermentation-in vitro

The effect of high levels of molasses in combinations with hay on digestibility of organic matter, microbial protein synthesis and volatile fatty acid production in vitro 7(1982)277

Fermentation-rumen

Nitrogen levels in low-roughage diets for efficient rumen microbial protein production 2(1977)255

Inhibition of ruminal deamination in vitro by formaldehyde treatment of sunflower-seed, soya bean and fish meals: response curves to protective treatment 2(1977)267

Rumen and post-rumen fermentation of spent tea leaf protein and other protein sources studied by the nylon bag method 7(1982)221

The energetic efficiency of rumen microbial protein synthesis in cattle given silage-based diets 13(1985)255

The effect of monensin on rumen fermentation in sheep fed on all-roughage and barley/roughage diets 15(1986)7

In vitro effects of avoparcin on protein degradability and rumen fermentation 15(1986)215

Evaluation of a dual flow continuous culture system for estimating bacterial fermentation in vivo of mixed diets containing various soya bean products 16(1986/1987)51

Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261

Effect of straw treatment and laboratory of incubation on degradation of neutral detergent residue in the rumen 18(1987)75

The effect of grinding and pelleting hay on digestibility, fermentation rate, digesta passage and rumen and faecal particle size in cows 19(1988)145

Fibre degradation in the rumens of animals receiving hay, untreated or ammonia-treated straw 19(1988)277

The effect of five different supplements on the degradation of straw in sheep given untreated barley straw 19(1988)289

Rumen fermentation, urease activity and performance of cattle given palm kernel cake-based diet 20(1988)79

Formation and utilization of end products of ligno cellulose degradation in ruminants 21(1988)141

The rumen as a model of the microbiology of fibre digestion in the human intestine 23(1989)227

Fermentation and utilization by lambs of sugarcane harvested fresh and ensiled with and without NaOH. 4. Ruminant kinetics 23(1989)323

Effect of urea treatment on rumen degradation characteristics of rice straws 24(1989)83

Fermentation-silage

Effect of maize silage harvest stage on yield, plant composition and fermentation losses 1(1976)313

Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299

The effect of silage additives containing formaldehyde on the fermentation of ryegrass ensiled at different dry matter levels and on the nutritive value of direct-cut silage 7(1982)303

Studies on the fermentation of chopped sugarcane 7(1982)411

The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Studies on fermentation of sorghum silage during storage, and its effect on milch animals 9(1983)143

Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185

The effects of ammonia on the fermentation of chopped sugarcane 9(1983)291

Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91

The effects of inoculation, addition of glucose and mincing on fermentation and proteolysis in ryegrass ensiled in laboratory silos 19(1988)85

SO₂-treated straw as a silage additive: fermentation data on lucerne with particular reference to protein degradation 22(1988)247

SO₂-treated straw as a silage additive: the participation of soluble and cell wall monosaccharide residues in lucerne silage fermentation 22(1988)255

Inoculation of corn silage and high-moisture corn with lactic acid bacteria and its effects on the subsequent fermentations and on feedlot performance of beef steers 25(1989) 23

Fertilizers

The quality of the green fraction of leaf protein concentrate from *Chenopodium quinoa* Willd. grown at different levels of fertilizer nitrogen 11(1984)239

The nutritional value of crop residue components from several wheat cultivars grown at different fertilizer levels 11(1984)301

Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Fibre

A comparison of three fibre methods for predicting the metabolizable energy content of sorghum grain for poultry 2(1977)197

The digestibility and nutritive value for dairy heifers of rations containing cellulose fibre 2(1977)287

Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3 (1978)227

Some remarks on the determination of acid-detergent fibre in feeds with cetyltrimethyl-ammonium bromide versus determination of detergent fibre with alkylarylsulfonic acid 6(1981)91

The study of several modifications of the neutral detergent fibre procedure 9(1983)19

Chemical properties of fibre in relation to nutritive quality of ammonia-treated forages 10(1983/1984)155

The effect of level and source of protein, fibre and fat in the diet on the energy value of compounded pig feeds 11(1984)11

A new biological method for determining amino acid digestibility in poultry feed-stuffs using a simple cannula, and the influence of dietary fibre on endogenous amino acid output 12(1984/1985)29

Effects of dietary concentrates on rumen digestion of fibrous feedstuffs 14(1986)193

The response of screened manure fibre to ozone and sodium hydroxide treatments 15 (1986)47

Influence of nutrient density on feed consumption, weight gain and gut capacity of

broilers, Leghorns and turkeys reared to 26 days of age 16(1986/1987)129

The effects of different sources of nitrogen supplementation on the digestion of fibre components in the rumen of steers 17(1987) 65

Composition and fibre digestion in morphological components of an alfalfa-timothy sward 19(1988)135

Dietary fibre and nitrogen excretion and retention by pigs 19(1988)185

Fibre degradation in the rumens of animals receiving hay, untreated or ammonia-treated straw 19(1988)277

The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988) 259

Effect of the nature of dietary fibre on transit time and faecal excretion in the growing pig 20(1988)327

Digestibility in vivo of ensiled grapevines (branches and leaves): influence of the system of analysis in the detergent fibre scheme on the prediction of digestibility 22 (1988)173

Measuring dietary fiber 23(1989)1

Determination of soluble fiber components: (1-<3; 1-<4)- β -D-glucans and pectins 23 (1989)15

Classification and measurement of plant polysaccharides 23(1989)27

Secretory response of the pig gut to non-starch polysaccharides 23(1989)55

Fibre-degrading microorganisms in the monogastric digestive tract 23(1989)91

In vitro fermentation of dietary fiber by human fecal organisms 23(1989)121

Approach to the energetic importance of fibre digestion in pigs. I. Importance of fermentation in the overall energy supply 23(1989) 141

Approach to the energetic importance of fibre digestion in pigs. II. An experimental approach to hindgut digestion 23(1989)169

Digestion and fermentation pattern of various dietary fiber sources in the rat 23(1989)195

Plant cell walls and monogastric diets 23 (1989)205

The rumen as a model of the microbiology of fibre digestion in the human intestine 23 (1989)227

The physiological effects of cellulose in the human large intestine 23(1989)241

Dietary fiber and energy balance: integration of the human ileostomy and in vitro fermentation models 23(1989)261

Comparative fiber degradation by mixed rumen fungi from Australian and U.S.A. cattle 23(1989)305

Response of high-producing dairy cows to high levels of dried corn gluten feed 24(1989)191

The effect of a protein energy supplement on pasture protein and fibre digestion in the rumen of grazing steers 25(1989)39

Fish meal

Inhibition of ruminal deamination in vitro by formaldehyde treatment of sunflower-seed, soya bean and fish meals: response curves to protective treatment 2(1977)267

The involvement of trimethylamine oxide in fish meal in the production of egg taint 8(1983)119

Substitution of finger millet (*Eleusine coracana*) and bulrush millet (*Pennisetum typhoides*) for maize in broiler feeds 10(1983/1984)291

Meat and bone and fish meals in balancer feeds for choice-fed broilers 11(1984)221

The amino acid and mineral composition of white fish meal containing enzyme-digested or untreated stickwater solids 14(1986)291

The effect of adding white fish meal containing enzyme digested or untreated stickwater solids to diets for broiler chickens 16(1986/1987)253

Fish protein concentrates for pre-ruminant calves: effects of processing temperatures and source of fish on protein digestibility and biological value 18(1987)181

Fish protein hydrolylates

Assessment of the production costs of fish protein hydrolysates 7(1982)147

Fish protein hydrolysates

General review of fish protein hydrolysates 7(1982)113

Preparation of fish protein hydrolysates 7(1982)125

Use of fish protein hydrolysate in milk replacers 7(1982)135

Performance of calves fed on milk replacers containing fish protein hydrolysate 7(1982)141

Fish silage - preparation, properties and uses 7(1982)153

Fish silage

Nutritional value and mercury content of fish silage for growing pigs 9(1983)169

Thiaminase activity in fish silage and moist fish feed 13(1985)39

Effect of formaldehyde treatment on the degradation of acid-preserved fish silage protein in vitro 16(1986/1987)297

Fish silage as a protein supplement for early-weaned calves 17(1987)165

Examination of stability, and its effect on the nutritive value, of fish silage in diets for growing pigs 21(1988)43

Fodder beet roots

The chemical composition, digestibility and energy value of fodder-beet roots 18(1987)225

Formaldehyde treatment

The digestion in the small intestine of young bulls of the protein of rapeseed meal treated or untreated with formaldehyde 7(1982)225

Energy and nitrogen balance, intake and growth rate of lambs fed on unwilted grass silages treated with a formaldehyde-acid mixture or untreated 9(1983)107

Effect of formaldehyde treatment on the degradation of acid-preserved fish silage protein in vitro 16(1986/1987)297

Complexes of urea and formaldehyde as non-protein nitrogen compounds in ruminant rations: a review 20(1988)1

Effect of formaldehyde treatment on the rumen digestion of barley and wheat grain and its influence on hay intake and digestion by sheep 25(1989)55

Formalin

The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21

Fox

Slaughterhouse by-products preserved by *Lactobacillus plantarum* fermentation as feed for mink and foxes 20(1988)287

Fractionation

Possibilities for multipurpose fractionation and alternative uses of plant biomass 21(1988)267

Fungi

Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977)7

The value of 'Pekilo protein' for growing pigs 2(1977)161

Production of fungal protein from rasped fresh cassava roots using 200- and 3000-litre fermentors 8(1983)313

The role of ciliate protozoa and fungi in the rumen digestion of plant cell walls 10(1983/1984)121

Effects of *Drechslera sorokiniana* infection on the yield and quality of tall fescue forage 15(1986)41

Nutrient losses during deterioration of hay in relation to changes in biochemical composition and microbial growth 15(1986)149

Association of rumen fungi with various forage grasses 16(1986/1987)273

Colonization of thick-walled plant tissues by anaerobic fungi 19(1988)25

Fungal contamination of hydroponic forage 20(1988)163

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. I. Composition and nutritive evaluation 22(1988)1

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. II. Effect of method of drying and response to amino acid supplementation 22(1988)13

Colonization of guinea grass by anaerobic rumen fungi in swamp buffalo and cattle 22(1988)161

Comparative fiber degradation by mixed rumen fungi from Australian and U.S.A. cattle 23(1989)305

G

Garden orach

The chemical composition and nutritive value for ruminants of four possible energy crops and their residues of anaerobic fermentation 15(1986)197

Genetics

Genetic variation in the feeding value of cereal straw 21(1988)127

Germination

The role of germination in sorghum reconstitution 14(1986)243

Giant star grass

The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171

(1-<3; 1-<4)- β -D-glucans

Determination of soluble fiber components: (1-<3; 1-<4)- β -D-glucans and pectins 23(1989)15

Glucose

The effects of inoculation, addition of glucose and mincing on fermentation and proteolysis in ryegrass ensiled in laboratory silos 19(1988)85

Glucosinolate

Low-glucosinolate rapeseed products for pigs - a review 9(1983)239

Glutamic acid

Amino acid composition of maize silage produced in the United Kingdom 4(1979)163

Gluten

Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375

Goat

Seasonal changes in composition and intake of chaparral by Spanish goats 8(1983)47

Utilization of alkali-treated and neutralized wheat straw-based rations for growing goat kids 9(1983)155

The value of urea-treated straw in diets of lactating goats 11(1984)67

Performance and carcass characteristics of Sudan desert sheep and goats on high roughage diets with added fat 13(1985)147

Changes in nutrient intake and performance by goats fed coffee pulp-based diets followed by a commercial concentrate 14(1986)1

The value of banana stalk as a feed for goats 15(1986)95

Effect of sodium hydroxide treatment of desert grass on carbohydrate digestibility by goats 17(1987)219

Cassava meal as supplement to napier grass diets for growing sheep and goats 18(1987)197

Comparison of cottonseed cake and dried blood as nitrogen sources for growing Sudan Nubian goats 22(1988)333

The intake and digestibility in goats of *Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989)45

Upgrading barley straw for goats: the effectiveness of a sodium hydroxide and urea dip method 24(1989)57

Grain

High grain content maize silage in fattening diets of young male cattle 16(1986/1987)63

Gram

Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315

Grapevine

Digestibility in vivo of ensiled grapevines (branches and leaves): influence of the system of analysis in the detergent fibre scheme on the prediction of digestibility 22 (1988)173

Grass, general

The accuracy of the pepsin-cellulase technique for estimating the dry matter digestibility in vivo of grasses and legumes 3(1978)277

Feed consumption and performance of dairy cows with alternate feedings of grass silage and maize silage 6(1981)337

Chemical composition and animal performance of grass forage of varying maturity stored as hay or silage 11(1984)35

Association of rumen fungi with various forage grasses 16(1986/1987)273

Grass, tropical

Bias when predicting crude protein, dry matter digestibility and voluntary intake of tropical grasses by near-infrared reflectance 9(1983)221

Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistani napier grass), *Pennisetum* sp. (bana grass), *Impomea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985)1

Grass, tropical pasture

Prediction of the dry matter digestibility of tropical grasses using a pepsin-cellulase assay 2(1977)247

Solubility of the protein of tropical pasture species and the rate of its digestion in the rumen 5(1980)183

Grass juice

Value of lucerne juice and grass juice as sources of protein for the growing pig 4 (1979)233

Grass silage

The use of effluent from grass silage in the diet of finishing pigs. I. Variation in composition of effluents 4(1979)263

The use of effluent from grass silage in the diet of finishing pigs. II. Assessment of nutritive value of fresh and stored effluent 4(1979)275

Effect of soaking, ensilage and hydrogen peroxide treatment of barley straw on rumen degradability 22(1988)295

Green berseem

Ensiling characteristics of cattle waste fermented with wheat straw and green berseem (*Trifolium alexandrinum*) 25(1989)149

Green panic

Preservation and storage of green panic (*Panicum maximum*) as moist hay with urea 20 (1988)87

Grinding

The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303

Ground mesquite

Pilot plant production of sulphur dioxide-treated mesquite wood. II. Digestibility and feedlot performance by lambs 22(1988)61

Groundnut cake

Feeding value of sunflower and groundnut cakes for broilers 4(1979)143

Feeding value of sunflower- and groundnut-cakes for laying hens 6(1981)63

Groundnut hulls

The effect of grinding groundnut hulls either with or without alkali treatment on digestibility of diet and on ruminal and blood components 15(1986)69

Groundnut meal

Detoxification of aflatoxins in groundnut meal 2(1977)67

Growth promoters

Energy metabolism in lambs fed silage diets with sucrose or a growth promoter 16 (1986/1987)203

Guinea fowl

A comparison of metabolizable energy content of eight common feed ingredients determined with young guinea fowls (keets) and pullet chicks 15(1986)1

Guinea grass

The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171

Colonization of guinea grass by anaerobic rumen fungi in swamp buffalo and cattle 22 (1988)161

H**Hair**

Utilization of processed pig hair in poultry diets 3(1978)243

Protein digestibility of hydrolyzed hog hair meal for pigs 3(1978)323

Harvesting time

Preservation of sorghum plant portions harvested, processed and ensiled at ten stages of maturity 7(1982)257

Influence of harvest date on the nutritive value and ensiling characteristics of maize stover 14(1986)11

The chemical composition of pea plant parts as related to harvesting time 16(1986/1987) 305

Yield, composition and feeding value of irrigated *Sesbania sesban* var. *Nubia* leaves harvested at latitude 38°N during a Mediterranean summer 18(1987)247

Treatment of whole-crop cereals with alkali. I. The influence of sodium hydroxide and ensiling on the chemical composition and in vitro digestibility of rye, barley and wheat crops harvested at increasing maturity and dry matter content 18(1987)257

The nutritive value of ensiled forages cut at three stages of growth 22(1988)237

Hay

Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607

Differences between cattle and sheep in their digestion and relative intake of a mature tropical grass hay 3(1978)41

Estimation of the digestibility of low-quality hays by cattle from measurements made with sheep 3(1978)51

Laboratory application of preservatives to hay and the effects of irregular distribution on mould development 6(1981)73

Chemicals for preventing moulding in damp hay 6(1981)323

Utilization of alkali-treated grain. 2. Utilization by steers of diets based on hay or straw and mixed with either NaOH-treated or rolled barley 6(1981)347

The effect of high levels of molasses in combinations with hay on digestibility of organic matter, microbial protein synthesis and volatile fatty acid production in vitro 7(1982)277

Laboratory tests of 8-quinolinol as an additive to volatile fatty acids for preserving hay and other feed-stuffs 7(1982)287

Ammoniation of wheat straw and native grass hay during baling of large round bales 8(1983)35

The effects of sulphur supplementation on cellulose digestion in vitro and on nutrient digestion, nitrogen metabolism and rumen characteristics of lambs fed on good quality

fescue and tropical star grass hays 8(1983) 129

Methods of improving the utilization of cereal straw by ruminants. I. Supplements of urea, molasses and dried grass and treatment with sodium hydroxide 8(1983)221

Methods of improving the utilization of cereal straw by ruminants. II. Further studies on dried grass as a supplement for lambs 8(1983)229

Problems of testing novel chemicals for the preservation of damp hay 8(1983)283

Manipulation of rumen fluid pH and its influence on cellulolysis in sacco, dry matter degradation and the rumen microflora of sheep offered either hay or concentrate 10(1983/1984)1

Associative effects of mixed feeds. II. The effect of dietary addition of bicarbonate salts on the voluntary intake and digestibility of diets containing various proportions of hay and barley 10(1983/1984)31

Chemical composition and animal performance of grass forage of varying maturity stored as hay or silage 11(1984)35

The effect of anhydrous ammonia and moisture content on the preservation and chemical composition of perennial ryegrass hay 11(1984)159

The effect of intake and hay: concentrate ratio upon digestibility and digesta passage 11(1984)167

Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279

A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum-pepsin treatment for predicting digestibility and intake of hays in sheep 12(1984/1985)101

The voluntary intake and digestibility of combinations of cereal crop residues and legume hay for sheep 12(1984/1985)241

Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistan napier grass), *Pennisetum* sp. (bana grass), *Impomea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985) 1

Storage of large round bales of hay harvested at different moisture concentrations and treated with sodium diacetate and/or covered with plastic 13(1985)69

Effect of ammonia level and time of exposure to ammonia on nutritional and preservative characteristics of dry and high-moisture coastal bermuda grass hay 14(1986)55

Dicoumarol concentrations in sweetclover hay treated with preservatives and in spoiled hay of high- and low-coumarin cultivars of sweetclover 14(1986)221

The effect of monensin on rumen fermentation in sheep fed on all-roughage and barley/roughage diets 15(1986)7

Nutrient losses during deterioration of hay in relation to changes in biochemical composition and microbial growth 15(1986)149

Estimating yield and nutrient losses due to rainfall on field-drying alfalfa hay 16(1986/1987)7

Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammer-milled lupin grain in oat hay or oat pasture based diets 16(1986/1987)143

Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and stored at different moisture contents 17(1987)45

Chemical composition and digestion in vitro of orchardgrass hay ammoniated by different techniques 17(1987)109

Effects of whole or ground corn with different forms of hay in 85% concentrate diets on digestion and passage rate in beef heifers 18(1987)151

The effect of grinding and pelleting hay on digestibility, fermentation rate, digesta passage and rumen and faecal particle size in cows 19(1988)145

Fibre degradation in the rumens of animals receiving hay, untreated or ammoniated straw 19(1988)277

Evaluation of mixed lucerne and caucasian bluestem hay diets fed to sheep 20(1988)19

Nitrogen utilization, ruminal fermentation and abomasal nitrogen flow in sheep fed caucasian bluestem hay supplemented with lucerne or urea 20(1988)33

Preservation and storage of green panic (*Panicum maximum*) as moist hay with urea 20(1988)87

Effect of level and frequency of maize supplementation on the utilization of low-quality grass hay by beef cows 24(1989)129

Oven and stack ammoniation of grass hays.
1. Changes in chemical composition in relation to digestibility in vitro and cell-wall degradability 24(1989)209

Oven and stack ammoniation of grass hays.
2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313

Effect of formaldehyde treatment on the rumen digestion of barley and wheat grain

and its influence on hay intake and digestion by sheep 25(1989)55

Water-drinking frequency and quantity of water drunk by steers fed hay or fresh forage 25(1989)207

Hominy

Feeding and nutritive value of hominy and corn grits for poultry 19(1988)313

House fly larvae

House fly larvae meal grown on municipal organic waste as a source of protein in poultry diets 4(1979)227

Hydrilla

Utilization of aquatic plants *Elodea canadensis* and *Hydrilla verticillata* in broiler chick diets 20(1988)155

Hydrolysis

Estimation of lysine in compounded poultry diets after enzymic digestion 3(1978)57

Hydroponic forage

Fungal contamination of hydroponic forage 20(1988)163

Hyperexcitability

Hyperexcitability in cattle fed ammoniated roughages 17(1987)121

I

Immunoglobulins

A continuous process for the production of purified porcine γ -globulin for use in pig-milk replacer supplements 17(1987)213

Intake

Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979)177

Effect of dry-matter concentration of milk substitute and method of feeding on intake and performance by lambs 5(1980)321

Dry matter, protein, energy and fibre intake by dairy heifers grazing a Rhodes grass (*Chloris gayana*) pasture 6(1981)15

Intake and digestibility in sheep fed on alkali-treated straw and concentrates, as complete diets or separate feeds 7(1982)271

Seasonal changes in composition and intake of chaparral by Spanish goats 8(1983)47

Energy and nitrogen balance, intake and growth rate of lambs fed on unwilted grass silages treated with a formaldehyde-acid mixture or untreated 9(1983)107

- Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197
- Bias when predicting crude protein, dry matter digestibility and voluntary intake of tropical grasses by near-infrared reflectance 9(1983)221
- Associative effects of mixed feeds. II. The effect of dietary addition of bicarbonate salts on the voluntary intake and digestibility of diets containing various proportions of hay and barley 10(1983/1984)31
- The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149
- The effect of intake and hay: concentrate ratio upon digestibility and digesta passage 11(1984)167
- A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum-pepsin treatment for predicting digestibility and intake of hays in sheep 12(1984/1985)101
- Improving the intake and nutritive value of wheat straw for beef cows 12(1984/1985)125
- The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12(1984/1985)141
- The effect of supplementing grass silage with sodium hydroxide-treated chopped or milled straw on intake and digestibility by wether sheep 12(1984/1985)215
- The voluntary intake and digestibility of combinations of cereal crop residues and legume hay for sheep 12(1984/1985)241
- Methods of improving the utilization of cereal straw by ruminants. IV. The effect on intake of level of supplementation with urea and dried grass 13(1985)281
- Changes in nutrient intake and performance by goats fed coffee pulp-based diets followed by a commercial concentrate 14(1986)1
- Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81
- Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139
- Effects of supplementing a wheat straw diet with urea or a urea-molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25
- The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99
- Intake, digestibility and energy utilization of corrugated paper diets 17(1987)57
- Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271
- Treatment of whole-crop cereals with alkali. 3. Voluntary intake and digestibility studies in sheep given immature wheat ensiled with sodium hydroxide, urea or ammonia 18(1987)283
- Treatment of whole-crop cereals with alkali. 4. Voluntary intake and growth in steers given wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55
- The intake and digestion of maize silage-based diets by dairy cows and sheep 20(1988)299
- Feed intake and milk yield at increasing supplement of a palmitic and stearic acid-rich type of fat in comparison with animal fat 22(1988)179
- Feed intake, milk yield and milk composition by replacing unprotected fat by Ca-soaps for dairy cows 22(1988)193
- The intake and digestibility in goats of *Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989)45
- Oven and stack ammoniation of grass hays. 2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313
- Effect of formaldehyde treatment on the rumen digestion of barley and wheat grain and its influence on hay intake and digestion by sheep 25(1989)55
- Intake, digestion and daily gain by cattle consuming bermudagrass (*Cynodon dactylon*) and supplemented with different combinations of ground corn, vegetable oil, urea and corn gluten and blood meals 25(1989)99
- ### Iodine
- Copper and iodine in pig diets with high glucosinolate rapeseed meal. I. Performance and thyroid hormone status of growing pigs fed on a diet with rapeseed meal treated with copper sulphate solution or untreated and supplements of iodine, copper or a quinoxaline derivative 22(1988)33
- Copper and iodine in pig diets with high glucosinolate rapeseed meal. II. Influence of iodine supplements for rations with rapeseed meal untreated or treated with copper ions on performance and thyroid hormone status of growing pigs 22(1988)45
- ### Iron
- The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171

Irrigation

Cultivation of maize in irrigated areas for silage and grain production and their use by ruminants 1(1976)237

Isoacids

Isoacids in ruminant nutrition: their role in ruminal and intermediary metabolism and possible influences on performances - a review 18(1987)169

Isopropyl alcohol treatment

Effects of isopropyl alcohol treatment of soya bean meal on nitrogen utilization by ruminants 14(1986)127

J**Jackbean**

Ripe beans of *Canavalia ensiformis* (jackbean) as feed ingredient for monogastric animals 20(1988)325

K**Kapok seed**

The effect of nutrient supplementation on the feeding value of kapok seed for poultry 9(1983)263

Karanja

The effect of kernels of karanja (*Pongamia glabra* Vent) on growth and feed efficiency in broiler chicks to 4 weeks of age 25(1989)201

Keratin

Utilization of processed pig hair in poultry diets 3(1978)243

Protein digestibility of hydrolyzed hog hair meal for pigs 3(1978)323

Chemical preparation of NaOH-keratin hydrolysate for improving the nutritive value of wheat straw 13(1985)215

Kidney bean

The use of kidney beans (*Phaseolus vulgaris*) as protein supplements in diets for young rapidly growing beef steers 12(1984/1985)1

L**Lactate**

Effect of lactate on the in vitro digestion of *Agropyron elongatum* by rumen microorganisms 16(1986/1987)161

Lactobacillus

The chemical composition of grass silages prepared with and without pre-treatment with inoculants containing *Lactobacillus plantarum* 13(1985)269

Slaughterhouse by-products preserved by *Lactobacillus plantarum* fermentation as feed for mink and foxes 20(1988)287

Lactobacillus

Efficacy of virginiamycin and a commercially-available lactobacillus probiotic in swine diets 8(1983)69

Stability of viable lactobacillus products 11(1984)261

Lantana seeds

Can lantana seeds be utilized in lamb rations? 9(1983)29

Leaf protein concentrate

Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189

Nutritive value of leaf-protein concentrate from potato haulm for rats and chicks 6(1981)413

Yield, composition and nutritive value of a leaf protein concentrate obtained by fractionating the shelled aerial part of *Pisum sativum* L. 8(1983)155

The quality of the green fraction of leaf protein concentrate from *Chenopodium quinoa* Willd. grown at different levels of fertilizer nitrogen 11(1984)239

The effect of different precipitating agents on quality of leaf protein concentrate from lucerne 12(1984/1985)11

The effect of toxic substances present in some leaf protein concentrates on their nutritive value 24(1989)151

Legumes

The analytical and biological accuracy of estimating the dry matter digestibility of different legume species 1(1976)61

The accuracy of the pepsin-cellulase technique for estimating the dry matter digestibility in vivo of grasses and legumes 3(1978)277

Ensilage of leguminous green forages in combination with different dry roughages and molasses 12(1984/1985)133

The voluntary intake and digestibility of combinations of cereal crop residues and legume hay for sheep 12(1984/1985)241

Rumen degradability of nitrogen in typical cultivated grasses and leguminous fodders 22(1988)79

Leucaena leucocephala

A note on the nutrient and mineral composition of *Leucaena leucocephala* in Western Nigeria 4(1979)221

Growth performance of, and mimosine excretion by, young chicks fed on *Leucaena leucocephala* 7(1982)247

Investigations into the effects of processing on the retention of the carotenoid fractions of *Leucaena leucocephala* during storage, and the effects of processing on mimosine concentration 9(1983)307

Nutrient and amino acid composition of *Leucaena leucocephala* (Lam.) de Wit 15(1986)183

The intake and digestibility in goats of *Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989)45

Lignification

Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71

Lignocellulose

Formation and utilization of end products of ligno cellulose degradation in ruminants 21(1988)141

Production of ethanol from lignocellulosic materials 21(1988)175

Chemistry of lignocellulose: methods of analysis and consequences of structure 21(1988)279

Biological structure of lignocellulose and its degradation in the rumen 21(1988)295

Listeria monocytogenes

Inhibition of *Listeria monocytogenes* by defined lactic microflora in gnotobiotic silages of lucerne, fescue, ryegrass and maize - influence of dry matter and temperature 2(1977)297

Lucerne

Nutritive value of leaf protein coagulum and cereals mixtures 4(1979)91

The effect of different precipitating agents on quality of leaf protein concentrate from lucerne 12(1984/1985)11

The chemical composition and nutritive value for ruminants of four possible energy crops and their residues of anaerobic fermentation 15(1986)197

Nitrogen utilization, ruminal fermentation and abomasal nitrogen flow in sheep fed caucasian bluestem hay supplemented with lucerne or urea 20(1988)33

Lucerne juice

Value of lucerne juice and grass juice as sources of protein for the growing pig 4(1979)233

Value of freshly produced lucerne juice as a source of supplemental protein for the growing pig 5(1980)215

Lucerne juice as a protein supplement for growing pigs: effects of mineral content of the diet and of the water supply 6(1981)35

Lucerne straw

SO₂-treated straw as a silage additive: the participation of soluble and cell wall monosaccharide residues in lucerne silage fermentation 22(1988)255

Lupin

Variability for protein and oil quality in *Lupinus albus* 9(1983)271

Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261

Lupin grain

Milling barley and lupin grain in diets for cattle 12(1984/1985)57

Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammermilled lupin grain in oat hay or oat pasture based diets 16(1986/1987)143

Lupin seed

The alkaloid content of sweet lupin seed used in feeding trials on pigs and rats 2(1977)59

Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189

Rumen degradation of protein in sunflower meal, rapeseed meal and lupin seed placed in nylon bags 11(1984)87

Digestibility and energy value of sweet lupin seed (*Lupinus albus* var. multolupa) in pigs 12(1984/1985)171

Protein and energy utilisation in rats of diets based on lupin seed (*Lupinus albus* var. multolupa) 15(1986)33

Lupin seed meal

A comparison between meals prepared from the seeds of different varieties of lupin as protein supplements to barley-based diets for growing pigs 2(1977)49

Lupinus angustifolius cv. Unicrop as a protein- and energy-source for weaner pigs 6(1981)27

Lupin-seed meal (*Lupinus albus* cv. Hamburg) as a source of protein for growing pigs 6(1981)285

Lupin-seed meal

Lupin-seed meal (*Lupinus angustifolius* cv. Uniwhite) as a protein supplement to barley-based diets for growing pigs 1(1976) 631

Lysine

Adjustments of the protein level for slaughter pigs and supplementation with lysine and methionine 1(1976)619

The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21

Estimation of lysine in compounded poultry diets after enzymic digestion 3(1978)57

Meat meal in the diet of the early-weaned pig. II. Amino acid supplementation 4(1979)43

Amino acid composition of maize silage produced in the United Kingdom 4(1979)163

The determination of available lysine in barley using *Tetrahymena pyriformis* W. 5 (1980)33

A comparison of three methods for the determination of available lysine in barley 5 (1980)43

Meat meal in the diet of the early-weaned pig. IV. The supplementation of diets with tryptophane, lysine and methionine 7 (1982)27

Lysine requirements of imported European pigs in the humid lowland tropics from 23 to 62 kg liveweight 7(1982)233

Proximate analysis, amino acid composition and tannin content of improved Nigerian sorghum varieties and their potential in poultry feeds 7(1982)359

Evaluation of optimum lysine and threonine supplements to a wheat and barley-based diet in rats 8(1983)163

Lysine requirement of weanling European pigs fed on yellow maize and groundnut cake diets in the humid lowland tropics 8 (1983)197

The absorption of dietary L-lysine and extruded L-lysine in pigs 15(1986)141

The influence of oat groats and dietary level of lysine on the laying performance of cross-bred hens 17(1987)271

The utilization by beef steers of a soft high lysine and a hard normal size maize cultivar in whole grain feedlot diets 21(1988)31

Lysinoalanine

Note on the lysinoalanine content of alkali-treated barley 7(1982)217

M**Magnesium**

Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Maize-cobs

Effect of level of sodium hydroxide treatment and volume of solution on the nutritive value of maize cobs 4(1979)1

Performance of heifers fed on diets based on NaOH-treated maize cobs and the effect of supplementary urea and source of carbohydrates 4(1979)97

Chemical composition, digestibility and feeding value of maize cobs 9(1983)121

High temperature and pressure processing of maize cobs. II. Animal response and effect of sodium metabisulphite 14(1986)161

Maize-ear

Maize grain or ears in concentrate diets for young fattening bulls 1(1976)393

Maize-general

Survey of the maize situation in Italy 1(1976) 109

Review of production and utilisation of maize for animal nutrition in Yugoslavia 1(1976) 115

Use of maize for ruminants and pig feeding in Greece 1(1976)125

Survey of the use of maize for livestock feeding in Spain 1(1976)129

Survey of the use of maize for livestock feeding in Portugal 1(1976)137

Maize and animal production in France 1 (1976)141

Survey of the use of maize for livestock feeding in Switzerland 1(1976)149

Survey of the use of maize for livestock feeding in Austria 1(1976)153

Use of maize for livestock feeding in the Federal Republic of Germany 1(1976)157

Survey of the use of maize for livestock feeding in Belgium 1(1976)167

Use of maize for livestock feeding in the Netherlands 1(1976)177

Prospects of growing maize in Finland 1 (1976)187

The current situation regarding the use of maize in Ireland 1(1976)195

The use of maize for livestock feeding in the United Kingdom 1(1976)199

- Discussion on session 1 1(1976)215
- The use of techniques of maize cultivation to maximise feed for beef production 1(1976)229
- Cultivation of maize in irrigated areas for silage and grain production and their use by ruminants 1(1976)237
- Selection objectives for maximum yields of feed with hybrid maize 1(1976)245
- Effect of agronomic factors on the yield of maize and on its composition in relation to its use by ruminants 1(1976)251
- Agronomic factors affecting the growth and composition of the maize plant 1(1976)263
- Discussion on session 2 1(1976)273
- Efficiency of harvesting and conservation 1 (1976)289
- Discussion on session 3 1(1976)327
- Maize silage with or without NPN, dehydrated whole-crop maize pellets or high moisture maize grain for finishing bulls 1 (1976)347
- Utilising the maize crop in various forms for beef production 1(1976)417
- Nutritive value of dehydrated whole-crop maize 1(1976)429
- The digestion and metabolism of the energy and protein of maize 1(1976)455
- Discussion on session 4 1(1976)485
- For beef cattle, all forms of maize can be used 1(1976)521
- Bull-beef production from the maize crop in Italy in large and small units 1(1976)531
- Discussion on session 5 1(1976)545
- Requirements for future research 1(1976)555
- Abomasal and plasma nitrogen constituents of steers fed on different physical forms of shelled maize 2(1977)225
- Waxy versus regular maize:energy value for growing pigs and utilization in piglet diets 4(1979)109
- Preservation of high-moisture maize - a comparison of gaseous and liquid anhydrous ammonia with methylene-bis-propionate 5 (1980)337
- Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983)99
- Measurement and prediction of digestibility of forage maize in The Netherlands 10 (1983/1984)301
- The evaluation of fresh sweet liquid whey as a protein supplement to maize for growing pigs 14(1986)171
- Response of different sorghum grain and maize varieties to reconstitution 15(1986)231
- The rate of dry matter disappearance from nylon bags of maize silage/grain diets as influenced by basal ration and processing of test samples 16(1986/1987)225
- High-fat maize in diets for pigs and sows 17 (1987)201
- The maize replacement value of cassava flour in rations for growing ewe lambs 20(1988)233
- Nutritive value of diets containing triticale and varying mixtures of triticale and maize for growing-finishing swine 22(1988)217
- Effect of level and frequency of maize supplementation on the utilization of low-quality grass hay by beef cows 24(1989)129
- Inoculation of corn silage and high-moisture corn with lactic acid bacteria and its effects on the subsequent fermentations and on feedlot performance of beef steers 25(1989)23
- Maize-grain**
- Maize grain or ears in concentrate diets for young fattening bulls 1(1976)393
- Feeding value of ensiled high-moisture maize grain with cob for growing-finishing pigs 20 (1988)145
- Maize-silage**
- Forage maize in the rations of dairy cows and fattening cattle in Bulgaria 1(1976)45
- Maize as a forage crop in Denmark 1(1976)183
- Influence of ensiling maize ears and grain on chemical composition, conservation losses and digestibility 1(1976)301
- Effect of maize silage harvest stage on yield, plant composition and fermentation losses 1(1976)313
- Energy supplementation of maize silage harvested at different maturity stages 1(1976)369
- Factors influencing the composition and nutritive value of ensiled whole-crop maize 1 (1976)381
- Rate of growth, feed utilisation and digestibility observed with diets containing ensiled maize grain or maize ears fed alone or in mixtures 1(1976)401
- Voluntary intake and efficiency of utilisation of whole-crop maize silage 1(1976)441
- Levels, sources and methods of nitrogen supplementation of maize silage for beef production 1(1976)465

- The nutritive value (energy) of maize silage 1 (1976)481
- Effect of weight at slaughter on the efficiency of fattening young cattle 1(1976)505
- Economic aspects of high-plane feeding of bulls with maize silage 1(1976)513
- Maize silage and whole maize plant pellets as the basic forages in complete feeds for milk cows 3(1978)23
- A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3 (1978)89
- Laboratory evaluation of ensiled mixtures of alkali-treated cattle excreta and whole-crop maize 3(1978)335
- Amino acid composition of maize silage produced in the United Kingdom 4(1979)163
- Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185
- Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91
- The effect of nitrogen supplements to maize silage on the amount and composition of the nitrogen fraction entering the intestine of cattle 13(1985)25
- High grain content maize silage in fattening diets of young male cattle 16(1986/1987)63
- The intake and digestion of maize silage-based diets by dairy cows and sheep 20 (1988)299
- Maize-solubles**
- Effects of dietary maize solubles and minerals on albumen quality of fresh and stored brown-shelled eggs 8(1983)177
- Maize-starch**
- Effect of defaunation on the loss of energy in wethers fed different quantities of cellulose and normal or steamflaked maize starch 16 (1986/1987)233
- Maize-stem**
- Colonization of thick-walled plant tissues by anaerobic fungi 19(1988)25
- Maize-stover**
- Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979) 177
- Nutritive value of single-cell protein produced by *Chaetomium cellulolyticum* grown on maize stover and pulp-mill sludge 11(1984) 211
- Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistan napier grass), *Pennisetum* sp. (bana grass), *Impomea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985) 1
- Influence of harvest date on the nutritive value and ensiling characteristics of maize stover 14(1986)11
- The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99
- Effects of moisture level, treatment time and soya bean addition on the nutritive value of urea-treated maize stover 19(1988)67
- Effects of ruminal degradability of the protein supplement on the utilization of maize-stover silage by growing beef cattle 20 (1988)241
- Effects of added dietary urea on the utilization of maize stover silage by growing beef cattle 22(1988)227
- Maize-whole**
- The utilization by beef steers of a soft high lysine and a hard normal size maize cultivar in whole grain feedlot diets 21(1988)31
- Maize-whole stillage**
- Maize whole stillage as a supplemental protein source for growing lambs and finishing steers 11(1984)181
- Maize silage**
- Effects of stage of maturity and addition of molasses on nutritive value of maize stover silage 12(1984/1985)65
- Maltase**
- Proximate chemical analysis of final cane molasses and effect of feeding 30% molasses on intestinal sucrose and maltase activities in the rat 3(1978)15
- Man**
- Assessing the importance of host-derived polysaccharides as carbon sources for bacteria growing in the human colon 23(1989) 109
- In vitro fermentation of dietary fiber by human fecal organisms 23(1989)121
- The physiological effects of cellulose in the human large intestine 23(1989)241
- Dietary fiber and energy balance: integration of the human ileostomy and in vitro fermentation models 23(1989)261

Manganese

The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171

Manioc

The digestion of diets containing manioc by young growing pigs 12(1984/1985)119

Manure

Utilization by sheep of dried pig faeces containing a high concentration of copper 3 (1978)31

Digestibility by cattle of diets containing dried pig faeces 3(1978)155

Laboratory evaluation of ensiled mixtures of alkali-treated cattle excreta and whole-crop maize 3(1978)335

The effects of air dried poultry manure on performance, organ weights and haematological measurements of weaner pigs reared in the tropics 4(1979)23

Utilization of cattle manure for single-cell protein production with *Chaetomium cellulolyticum* 4(1979)199

Digestibility of wet cage layer excreta and an excreta-supplemented diet by cattle as determined by two methods 5(1980)77

Utilization of cattle manure for Torula yeast production from straw hydrolysates 5(1980)175

Performance of fattening cattle and lactating beef cows on diets high in chemically treated wheat straw and poultry litter 5 (1980)299

Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51

The determination of digesta particle size in some herbivores 7(1982)35

Fermentation characteristics and feeding value of ensiled poultry litter containing wheat straw, bagasse or sawdust 7(1982)341

Nutritive value of ensiled broiler litter for cattle 8(1983)19

Corrugated cardboard and dried poultry manure as dietary ingredients for sheep 14 (1986)265

The response of screened manure fibre to ozone and sodium hydroxide treatments 15 (1986)47

Digestibility and beef performance after feeding NaOH- or KOH-treated pig wastes 17 (1987)235

Evaluation of cardboard and dried poultry waste as feed ingredients for ruminants 18 (1987)209

Animal excreta as ruminant feed - scope and limitations under Indian conditions 19 (1988)1

Effect of the nature of dietary fibre on transit time and faecal excretion in the growing pig 20(1988)327

Nutritive value of rice straw (*Oryza sativa*) ensiled with animal excreta and rumen digesta 24(1989)69

Mastication

The effect of method of preservation of forages on their resistance to reduction in particle size with an artificial masticator 24 (1989)185

Maturation

Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro degradability during maturation 17(1987)15

Meat and bone meal

Protein quality of mixtures of animal protein meals 9(1983)79

The digestion by sheep of barley and maize-based diets containing either meat and bone meal or soya bean meal 15(1986)301

Meat meal

Meat meal in the diet of the early-weaned pig. I. A comparison of meat meal and soya bean meal 4(1979)33

Meat meal in the diet of the early-weaned pig. II. Amino acid supplementation 4(1979)43

Meat meal in the diet of the early-weaned pig. III. Meat meal quality and the processing of meat meals 4(1979)53

Meat meal in the diet of the early-weaned pig. IV. The supplementation of diets with tryptophane, lysine and methionine 7 (1982)27

Meat and bone and fish meals in balancer feeds for choice-fed broilers 11(1984)221

Meat quality

Effect of avoparcin on performance, body composition and meat quality of finishing bulls 8(1983)237

Menadione

Comparison of the relative biological availability and stability of commercial sources of vitamin K activity 3(1978)329

Mercury

Nutritional value and mercury content of fish silage for growing pigs 9(1983)169

Metabolic faecal output

Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279

Methane digester effluent

Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47

Methionine

Adjustments of the protein level for slaughter pigs and supplementation with lysine and methionine 1(1976)619

Meat meal in the diet of the early-weaned pig. II. Amino acid supplementation 4(1979)43

Limiting amino acids in dried microbial cells given to young turkeys 4(1979)209

Meat meal in the diet of the early-weaned pig. IV. The supplementation of diets with tryptophane, lysine and methionine 7(1982)27

Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Proteen 17(1987)1

Methylene-bis-propionate

Preservation of high-moisture maize - a comparison of gaseous and liquid anhydrous ammonia with methylene-bis-propionate 5(1980)337

Micro-ingredients

The effect of dilution levels in premixes on micro-ingredient dispersion in animal feeds 8(1983)139

Microbial protein synthesis

Effect of dietary nitrogen source on microbial protein synthesis, dietary protein degradation and nutrient digestion in steers 14(1986)117

Relative intakes, digestibility and bacterial protein synthesis by sheep and cattle fed high-roughage diets 16(1986/1987)75

Micronization

Effects of micronization on the digestibility of whole soya beans and rapeseeds for the growing pig 3(1978)179

Milk substitutes

The replacement of skim milk by dried microbial cells and whey in milk diets for calves 2(1977)85

Digestibility of protein and amino acids of a fermentation single-cell protein for veal calves 2(1977)377

Effect of dry-matter concentration of milk substitute and method of feeding on intake and performance by lambs 5(1980)321

Use of fish protein hydrolysate in milk replacers 7(1982)135

Performance of calves fed on milk replacers containing fish protein hydrolysate 7(1982)141

The effects on feed intake, growth and digestibility of nutrients of including calcium soaps of fat in diets for young calves 14(1986)103

A continuous process for the production of purified porcine γ -globulin for use in pig-milk replacer supplements 17(1987)213

Fish protein concentrates for pre-ruminant calves: effects of processing temperatures and source of fish on protein digestibility and biological value 18(1987)181

The effects of method of processing of starters, tallow inclusion and roughage supplementation on the performance of early-weaned calves 19(1988)231

An evaluation of skim milk powders subjected to different heating conditions during processing as dietary protein sources for the young pig 22(1988)203

Evaluation of pea protein concentrate as a source of protein in milk replacers for Holstein calves 24(1989)267

Milling

Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983)99

The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149

Milling barley and lupin grain in diets for cattle 12(1984/1985)57

The effect of mechanical treatment on the nutritive value of high-moisture barley and the performance of growing-finishing pigs 24(1989)121

Mimosine

Growth performance of, and mimosine excretion by, young chicks fed on *Leucaena leucocephala* 7(1982)247

Investigations into the effects of processing on the retention of the carotenoid fractions of *Leucaena leucocephala* during storage, and the effects of processing on mimosine concentration 9(1983)307

Minerals, general

The use of whole barley diets fortified with solutions of urea, minerals and vitamins for lambs 2(1977)307

Soil chemistry and mineral problems in farm livestock. A review 5(1980)95

Lucerne juice as a protein supplement for growing pigs: effects of mineral content of the diet and of the water supply 6(1981)35

Effects of dietary maize solubles and minerals on albumen quality of fresh and stored brown-shelled eggs 8(1983)177

Mink

Slaughterhouse by-products preserved by *Lactobacillus plantarum* fermentation as feed for mink and foxes 20(1988)287

Modelling

Evaluation of mathematical models to describe neutral detergent residue in terms of its susceptibility to degradation in the rumen 15(1986)249

Sources of error in predicting digestible dry matter from the acid-detergent fiber content of forages 21(1988)205

Molasse

The nutritional value for sheep of molasses and condensed molasses solubles 2(1977)143

Proximate chemical analysis of final cane molasses and effect of feeding 30% molasses on intestinal sucrase and maltase activities in the rat 3(1978)15

Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299

Distillery effluents as animal feed: the use of condensed beet molasses stillage (CBMS) in broiler feeding 5(1980)233

Methods of improving the utilization of cereal straw by ruminants. I. Supplements of urea, molasses and dried grass and treatment with sodium hydroxide 8(1983)221

Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185

Effects of stage of maturity and addition of molasses on nutritive value of maize stover silage 12(1984/1985)65

Ensiling of leguminous green forages in combination with different dry roughages and molasses 12(1984/1985)133

Effects of molasses/urea and bacterial inoculant additives on silage quality, dry matter recovery, and feeding value for cattle 12(1984/1985)205

Effect of sugarcane condensed molasses solubles, sugarcane molasses and monensin on performance and volatile fatty acid production in finishing steers 12(1984/1985)275

Differences in chemical composition and digestibility of beet and cane molasses 13(1985)83

Methods of improving the utilization of cereal straw by ruminants. V. The effect of molasses and dried grass 13(1985)293

Effects of supplementing a wheat straw diet with urea or a urea-molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25

Sugar beet molasses for growing and fattening pigs 18(1987)131

Effect of sorghum straw alone or in combination with molasses and nitrogen sources on performance of Sudan desert lambs 19(1988)351

A note on the use of sugar-beet pulp silage and molasses in the diet of lactating dairy cows 20(1988)13

The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988)259

Digestion of diets containing molassed or plain sugar-beet pulp by growing pigs 23(1989)67

Utilization of sugar beet molasses and monensin for finishing dairy bullocks 25(1989)11

Molybdenum

Reducing the potential copper toxicity of concentrates to sheep by the use of molybdenum and sulphur supplements 2(1977)235

Monensin

Monensin-sodium as a performance-promoting additive for fattening bulls and its impact on carcass and meat quality characteristics 7(1982)401

Effect of sugarcane condensed molasses solubles, sugarcane molasses and monensin on performance and volatile fatty acid production in finishing steers 12(1984/1985)275

The effect of monensin on rumen fermentation in sheep fed on all-roughage and barley/roughage diets 15(1986)7

Utilization of sugar beet molasses and monensin for finishing dairy bullocks 25(1989)11

Monocarboxyl compounds

Thermofractographic determination of monocarboxyl compounds in animal waste fats used as feed fat 13(1985)227

Mycotoxins

In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations 25(1989)169

N

Napier grass

Cassava meal as supplement to napier grass diets for growing sheep and goats 18(1987)197

Navy bean

The dye absorption test as an indicator of the nutritive quality of heat-processed navy bean 9(1983)71

Neutral detergent

Effect of straw treatment and laboratory of incubation on degradation of neutral detergent residue in the rumen 18(1987)75

Neutral detergent fibre

The study of several modifications of the neutral detergent fibre procedure 9(1983)19

Neutral detergent residue

Evaluation of mathematical models to describe neutral detergent residue in terms of its susceptibility to degradation in the rumen 15(1986)249

Nitrogen

Effect on digestibility and nitrogen content of barley straw of different ammonia treatments 2(1977)131

Abomasal and plasma nitrogen constituents of steers fed on different physical forms of shelled maize 2(1977)225

Nitrogen levels in low-roughage diets for efficient rumen microbial protein production 2(1977)255

Nitrogen metabolism in broiler chickens consuming the bacterial strain *Alcaligenes eutrophus* 5(1980)241

Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375

The effects of sulphur supplementation on cellulose digestion in vitro and on nutrient digestion, nitrogen metabolism and rumen characteristics of lambs fed on good quality fescue and tropical star grass hays 8(1983)129

The effect of prolonged storage on the digestibility and nitrogen content of ammonia-treated barley straw 8(1983)147

The effect of some pre-ensiling treatments on silage composition and nitrogen disappearance in the rumen 8(1983)259

Energy and nitrogen balance, intake and growth rate of lambs fed on unwilted grass silages treated with a formaldehyde-acid mixture or untreated 9(1983)107

Note on high-nitrate para grass (*Brachiaria mutica*) silage given to rabbits 9(1983)325

Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65

The effect of nitrogen supplements to maize silage on the amount and composition of the nitrogen fraction entering the intestine of cattle 13(1985)25

Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47

Effect of dietary nitrogen source on microbial protein synthesis, dietary protein degradation and nutrient digestion in steers 14(1986)117

Effects of isopropyl alcohol treatment of soya bean meal on nitrogen utilization by ruminants 14(1986)127

Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Effects of processing time and moisture content on amino acid composition and nitrogen characteristics of feather meal 14(1986)279

Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21

The effect of β -glucanase on the utilization of starch and nitrogen by broiler chickens fed on barley of low- or high-viscosity 15(1986)83

Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammer-milled lupin grain in oaten hay or oaten pasture based diets 16(1986/1987)143

The effect of enzymatic treatment on amino acid content and nitrogen characteristics of feather meal 16(1986/1987)151

The apparent and true ileal digestibility of nitrogen and amino acids in lactic casein for the growing pig 16(1986/1987)169

The effects of different sources of nitrogen supplementation on the digestion of fibre components in the rumen of steers 17(1987)65

Dietary fibre and nitrogen excretion and retention by pigs 19(1988)185

Effect of sorghum straw alone or in combination with molasses and nitrogen sources on performance of Sudan desert lambs 19(1988)351

Complexes of urea and formaldehyde as non-protein nitrogen compounds in ruminant rations: a review 20(1988)1

The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988) 259

Rumen degradability of nitrogen in typical cultivated grasses and leguminous fodders 22(1988)79

Comparison of cottonseed cake and dried blood as nitrogen sources for growing Sudan Nubian goats 22(1988)333

Distribution of liquid supplement injected into large bales of roughage and comparison of nitrogen supplements for stocker cattle fed low-quality roughages 24(1989)13

Oven and stack ammoniation of grass hays. 2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313

Effect of blood meal processing method on performance and nitrogen utilization by growing cattle 24(1989)337

Noug meal

The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99

Nutritive value

Genetic variation in the feeding value of cereal straw 21(1988)127

Nutritive value-age influence

Influence of age on the nutrient utilization of diets for broilers 6(1981)405

Nutritive value-almond

Nutritive value of almond hulls for sheep 8(1983)63

Nutritive value-apple

Nutritive value of apple pomace for ruminants 16(1986/1987)311

Nutritive value-arhar

Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315

Nutritive value-bagaruwa

Biochemical evaluation of bagaruwa seeds (*Acacia nilotica*) for use as livestock feed 11(1984)45

Nutritive value-banana

Nutritive value of banana stalk (*Musa cavendishi*) as a feed for sheep 22(1988)327

Nutritive value-barley

Nutritional value of waxy barleys for broiler chickens 8(1983)25

Feeding value of hydroponically sprouted barley for poultry and pigs 13(1985)183

Nutrient content of hydroponically sprouted barley 13(1985)191

The nutritive value of barley for monogastric animals and for ruminants as affected by artificial drying and alkali and anaerobic storage 15(1986)285

Nutritional value for chickens of dried distillers-spent-grain from barley and dehulled barley 17(1987)145

The nutritive value of hulless barley for swine 19(1988)191

The effect of mechanical treatment on the nutritive value of high-moisture barley and the performance of growing-finishing pigs 24(1989)121

Effect of enzyme supplementation on the nutritive value of hulless barley in chicken diets 24(1989)233

Nutritive value-bean, winged

The proximate composition and nutritive value of the winged bean *Psophocarpus tetragonolobus* (L.) DC for broilers 11(1984)231

Nutritive value-brewer's dried grains

Evaluation of brewer's dried grains in the diets of laying hens 6(1981)169

Evaluation of brewer's dried grains in the diets of broiler chickens 6(1981)179

Nutritive value-buckwheat

A nutritional evaluation of buckwheat (*Fagopyrum esculentum*) 3(1978)95

Nutritive value-cabbage

Cabbage (*Brassica oleracea*) in the diet of growing-finishing pigs 5(1980)69

Nutritive value-canola meal

The nutritive value of canola meal for early-weaned pigs 18(1987)45

Nutritive value-Canola rapeseed

The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303

Nutritive value-cereals, general

Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6(1981)309

Nutritive value-cocoa

Studies on the feeding value of agro-industrial by-products. III. Digestibility of cocoa-pod and cocoa-pod-based diets by ruminants 13 (1985)249

Studies on the feeding value of agro-industrial by-products: strategies for improving the utilisation of cocoa-pod-based diets by ruminants 20(1988)189

Nutritive value-coffee pulp

A note on the nutritive value of dried instant coffee residue for broiler chickens and turkey poults 10(1983/1984)285

Nutritive value-cottonseed

A review of the nutritive value and utilization of whole cottonseed, cottonseed meal and associated by-products by dairy cattle 18 (1987)89

Nutritive value-fats

Investigations on the nutritional value for ruminants of various fats 19(1988)111

The nutritional value for broilers of fats characterized by short-chain fatty acids as affected by level of inclusion and age 22(1988) 105

Nutritive value-fibre

The digestibility and nutritive value for dairy heifers of rations containing cellulose fibre 2(1977)287

Nutritive value-fish silage

Examination of stability, and its effect on the nutritive value, of fish silage in diets for growing pigs 21(1988)43

Nutritive value-forages, general

A new technique for upgrading the nutritive value of roughages under field conditions 3 (1978)201

Principles governing the use of chemical methods for assessing the nutritive value of forages: a review 3(1978)265

Predicting the energetic value of forage from digestibility of organic matter in vitro or solubility in potassium hydroxide 6(1981) 235

Different sources of ammonia for improving the nutritive value of low quality roughages 6(1981)377

Chemical properties of fibre in relation to nutritive quality of ammonia-treated forages 10(1983/1984)155

Drying treatment drastically affects feed evaluation and feed quality with certain tropical forage species 22(1988)69

The nutritive value of ensiled forages cut at three stages of growth 22(1988)237

Nutritive value-fungal protein

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. I. Composition and nutritive evaluation 22(1988)1

Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. III. Nutritive value for chicks and piglets 22(1988)23

Nutritive value-gram

Nutritional evaluation of gram (*Cicer arietinum*) and arhar (*Cajanus cajan*) chunies for cattle 5(1980)315

Nutritive value-groundnut

Feeding value of sunflower and groundnut cakes for broilers 4(1979)143

Feeding value of sunflower- and groundnut-cakes for laying hens 6(1981)63

Nutritive value-hominy

Feeding and nutritive value of hominy and corn grits for poultry 19(1988)313

Nutritive value-kapok seed

The effect of nutrient supplementation on the feeding value of kapok seed for poultry 9 (1983)263

Nutritive value-leaf protein

Nutritive value of leaf protein coagulum and cereals mixtures 4(1979)91

Nutritive value-leaf protein concentrate

Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189

Nutritive value of leaf-protein concentrate from potato haulm for rats and chicks 6 (1981)413

The effect of toxic substances present in some leaf protein concentrates on their nutritive value 24(1989)151

Nutritive value-Leucaena leucocephala

A note on the nutrient and mineral composition of *Leucaena leucocephala* in Western Nigeria 4(1979)221

Nutritive value-lupin seed

Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189

Nutritive value-maize

Factors influencing the composition and nutritive value of ensiled whole-crop maize 1 (1976)381

Nutritive value of dehydrated whole-crop maize 1(1976)429

The nutritive value (energy) of maize silage 1 (1976)481

Effect of level of sodium hydroxide treatment and volume of solution on the nutritive value of maize cobs 4(1979)1

Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979)177

Nutritive value of thermoammoniated and steam-treated maize stover II. Rumen metabolites and rate of passage 4(1979)187

Chemical composition, digestibility and feeding value of maize cobs 9(1983)121

Influence of harvest date on the nutritive value and ensiling characteristics of maize stover 14(1986)11

Effects of moisture level, treatment time and soya bean addition on the nutritive value of urea-treated maize stover 19(1988)67

Feeding value of ensiled high-moisture maize grain with cob for growing-finishing pigs 20(1988)145

Nutritive value-manure

Fermentation characteristics and feeding value of ensiled poultry litter containing wheat straw, bagasse or sawdust 7(1982)341

Nutritive value of ensiled broiler litter for cattle 8(1983)19

Nutritive value-molasse

The nutritional value for sheep of molasses and condensed molasses solubles 2(1977)143

Nutritive value-navy bean

The dye absorption test as an indicator of the nutritive quality of heat-processed navy bean 9(1983)71

Nutritive value-oat

The effects of cereal particle size and pelleting on the nutritive value of oat-based diets for the growing pig 8(1983)91

Nutritive value of irradiated and β -glucanase-treated wild oat groats (*Avena fatua* L.) for broiler chickens 16(1986/1987)243

Nutritive value of naked oats for ruminants 18(1987)83

Nutritive value-olive pulp

Nutritional value for rabbits of olive pulp and the effects on their visceral organs 25(1989)79

Nutritive value-para-rubber seed

Chemical composition and nutritional value of para-rubber seed and its products for chickens 10(1983/1984)257

Nutritive value-pea

Yield, composition and nutritive value of a leaf protein concentrate obtained by fractionating the shelled aerial part of *Pisum sativum* L. 8(1983)155

Nutritional and growth studies with pea-crop meals and peas for growing-finishing pigs 16(1986/1987)119

The nutritive value of white- and dark-flowered cultivars of pea for growing-finishing pigs 17(1987)245

Dry matter yield and chemical composition of pigeon pea (*C. cajan*) leaf meal and the nutritive value of pigeon pea leaf meal and grain meal for laying hens 24(1989)111

Nutritive value-peanut

Effect of sodium hydroxide treatment and/or extrusion cooking on the nutritive value of peanut hulls 12(1984/1985)187

Nutritive value-pigeon pea

Nutritional value of pigeonpea (*Cajanus cajan*) meal in poultry diets 25(1989)123

Nutritive value-potato

Liquid potato feed for finishing pigs: feeding value, inclusion rate and storage properties 15(1986)129

Nutritive value-proteins and amino acids

Composition and nutritive value of single-cell protein (SCP) 1(1976)9

Nutritive value-pungam cake

Nutritive value of pungam (*Pongamia glabra* Vent) cake for sheep 22(1988)321

Nutritive value-rapeseed

The nutritive value of tower rapeseed meal for swine 3(1978)235

Influence of steam pelleting on the feeding value of tower and candle rapeseed products for broiler chickens 3(1978)251

Nutritive value-rubber seed

Nutritive value of Nigerian rubber seed (*Hevea brasiliensis*) meal and oil. III. Performance characteristics, relative organ weights, hematocrit and plasma metabolites of growing female rats fed corn diets containing rubber seed meal, soya bean meal or casein 20(1988)125

Nutritive value-rumen contents

Nutritive value of rumen contents for monogastric animals 2(1977)351

Nutritive value-rye grain

Feeding value of alkali-treated whole rye (*Secale cereale* L.) grain for lactating cows and its digestibility for sheep 10(1983/1984)77

Nutritive value-Sesbania sesban

Yield, composition and feeding value of irrigated *Sesbania sesban* var. Nubia leaves harvested at latitude 38°N during a Mediterranean summer 18(1987)247

Nutritive value-silage

A calorimetric investigation of the nutritive value of formic acid- and formaldehyde-formic acid-treated grass silages 6(1981)245

The effect of silage additives containing formaldehyde on the fermentation of ryegrass ensiled at different dry matter levels and on the nutritive value of direct-cut silage 7(1982)303

Nutritional value and mercury content of fish silage for growing pigs 9(1983)169

Chemical composition and animal performance of grass forage of varying maturity stored as hay or silage 11(1984)35

Nutritive value-silage effluent

The use of effluent from grass silage in the diet of finishing pigs. II. Assessment of nutritive value of fresh and stored effluent 4(1979)275

Nutritive value-single-cell protein

Nutritive value of single-cell protein produced by *Chaetomium cellulolyticum* grown on maize stover and pulp-mill sludge 11(1984)211

Nutritive value-sorghum

Comparison of the nutritional value of low and medium tannin sorghum grains for pigs 8(1983)85

Nutritive value-soya bean

Nutrient and trypsin inhibitor content of hydroponically sprouted soya beans 13(1985)203

Nutritive value-spruce

Nutritive value of spruce muka (foliage) for the growing chick 3(1978)63

Nutritive value-straw

Ensilaged alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensiled or frozen alkali-treated straw and ryegrass silage 3(1978)133

Alkali treatment of straw for ruminants. II. Nutritive value of straw ensiled after alkali treatment. 3(1978)289

The effect of ammoniation on the intake and nutritive value of straw 4(1979)169

The effect of urease enzyme on treatment time and the nutritive value of straw treated with ammonia as urea 8(1983)271

The value of urea-treated straw in diets of lactating goats 11(1984)67

Improving the intake and nutritive value of wheat straw for beef cows 12(1984/1985)125

A comparison of alkali treatment methods to improve the nutritive value of straw. I. Digestibility and metabolizability 12(1984/1985)295

Nutritive evaluation of uromol-wheat straw as a basal ration for buffaloes 13(1985)103

Chemical preparation of NaOH-keratin hydrolysate for improving the nutritive value of wheat straw 13(1985)215

Urea as a source of ammonia for improving the nutritive value of wheat straw 14(1986)67

A comparison of alkali treatment methods used to improve the nutritive value of straw. II. In sacco and in vitro degradation relative to in vivo digestibility 14(1986)215

Evaluation of the increase in feeding value of barley straw following mild treatment with sulphur dioxide 15(1986)13

Yield and potential feeding value of straws of grain sorghum cultivars developed in Nigeria 17(1987)285

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility in vitro and cell wall degradability 19(1988)159

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173

The effect of variation in the water content of wheat straw before ammoniation on its nutritive value for beef cattle 20(1988)111

The feeding value of ammoniated flax straw, wheat straw and wheat chaff for beef cattle 21(1988)57

Nutritive value of rice straw (*Oryza sativa*) ensiled with animal excreta and rumen digesta 24(1989)69

Nutritive value-sugar cane

Nutritive value for ruminants of sugar cane bagasse ensiled after spray treatment with different levels of NaOH 9(1983)1

Nutritive value-sunflower

Feeding value of sunflower and groundnut cakes for broilers 4(1979)143

Feeding value of sunflower- and groundnut-cakes for laying hens 6(1981)63

Nutritive value-tamarind

The nutritive value of tamarind seeds for broiler chicks 22(1988)285

Nutritive value-triticale

A nutritional evaluation of triticale with pigs 9(1983)49

Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19

Nutritional value of triticale (Carman) for broiler diets 22(1988)119

Nutritive value of diets containing triticale and varying mixtures of triticale and maize for growing-finishing swine 22(1988)217

Nutritive value-vinasse

Protein and energy value of vinasse for pigs 5 (1980)221

Nutritive value-wheat

Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin) 11(1984)293

Chemical composition and nutritive value of Canadian winter wheats (*Triticum aestivum* L.) given to laboratory rats 12(1984/1985)75

Feeding value of sprouted wheat (*Triticum aestivum*) for beef cattle finishing diets 15 (1986)113

Nutritive value-wheat residue

The nutritional value of crop residue components from several wheat cultivars grown at different fertilizer levels 11(1984)301

Nutritive value-whey

The nutritive value of liquid whey, either sour or sweet, when given in restricted amounts to the growing pig 3(1978)163

Nutritive value-yeast

Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6(1981)309

O**Oak browse**

Composition of kermes oak browse as affected by shade and stage of maturity 21(1988)1

Oat grain

The effects of cereal particle size and pelleting on the nutritive value of oat-based diets for the growing pig 8(1983)91

Oat groats

The influence of oat groats and dietary level of lysine on the laying performance of cross-bred hens 17(1987)271

Oat straw

Studies on ammonia-treated straw. I. The effects of type and level of ammonia, moisture content and treatment time on the digestibility in vitro and enzyme soluble organic matter of oat straw 7(1982)45

Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71

The effect of calcium hydroxide treatment on the digestibility of chopped oat straw supplemented with a solution containing urea, calcium, phosphorus, sodium, trace elements and vitamins 7(1982)93

The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility in vitro and cell wall degradability 19(1988)159

The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173

Oats

Utilization of alkali-treated grain. 1. Alkali-treated grain in complete diets for steers and lambs 6(1981)273

The feeding value for ruminants of straw and whole-crop barley and oats treated with anhydrous or aqueous ammonia or urea 8 (1983)247

Anaerobically stored high moisture cereals for growing pigs 12(1984/1985)159

Digestibility of dry matter, nitrogen and energy by dairy cows fed whole or hammer-milled lupin grain in oaten hay or oaten pasture based diets 16(1986/1987)143

Nutritive value of irradiated and β -glucanase-treated wild oat groats (*Avena fatua* L.) for broiler chickens 16(1986/1987)243

Nutritive value of naked oats for ruminants 18(1987)83

The effect of β -glucanase supplementation on nutrient digestibility and growth in broilers given diets containing barley, oat groats or wheat 25(1989)193

Oil palm slurry

Oil palm slurry (OPS) as a partial replacement for maize in the diets of broiler chickens 17(1987)157

Olaquinox

Olaquinox as performance-promoting feed additive for growing pigs 4(1979)117

Olive cake

Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139

Olive pulp

Nutritional value for rabbits of olive pulp and the effects on their visceral organs 25(1989)79

Orange

Effect of urea on the ensiling process of orange pulp 12(1984/1985)233

The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20(1988)335

Organic matter, general

Comparison of laboratory methods for predicting the organic matter digestibility of forages 2(1977)337

Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3(1978)227

Ozone

The response of screened manure fibre to ozone and sodium hydroxide treatments 15(1986)47

Effects of ozonation on the chemical composition and in vitro degradation of triticale internodes and red clover stems 24(1989)159

P**Palm kernel cake**

Rumen fermentation, urease activity and performance of cattle given palm kernel cake-based diet 20(1988)79

Palm kernel meal

Palm kernel meal as a feed for poultry. 1. Composition of palm kernel meal and availability of its amino acids to chicks 16(1986/1987)179

Palm kernel meal as a feed for poultry. 2. Diets containing palm kernel meal for starter and grower pullets 16(1986/1987)187

Palm kernel meal as a feed for poultry. 3. Replacement of groundnut cake by palm kernel meal in broiler diets 16(1986/1987)195

Palm kernel meal as a feed for poultry. 4. Use of palm kernel meal by laying birds 20(1988)279

Palm oil mill effluent

The intake and digestibility in goats of *Leucaena leucocephala* supplemented with dehydrated palm oil mill effluent 24(1989)45

Palmitic acid

Feed intake and milk yield at increasing supplement of a palmitic and stearic acid-rich type of fat in comparison with animal fat 22(1988)179

Paper diets

Intake, digestibility and energy utilization of corrugated paper diets 17(1987)57

Para grass

Note on high-nitrate para grass (*Brachiaria mutica*) silage given to rabbits 9(1983)325

Para-rubber seed

The influence of processing and storage on hydrogen cyanide and tannin contents of para-rubber seed and its products 9(1983)319

Chemical composition and nutritional value of para-rubber seed and its products for chickens 10(1983/1984)257

Particle size

The effect of method of preservation of forages on their resistance to reduction in particle size with an artificial masticator 24(1989)185

Particle size distribution

Particle size distribution of a feed premix containing amprolium 3(1978)211

The effects of cereal particle size and pelleting on the nutritive value of oat-based diets for the growing pig 8(1983)91

Pasture plants

The relationship between the intake of herbage grazed by dairy cows and its digestibility 1(1976)53

Pea

Yield, composition and nutritive value of a leaf protein concentrate obtained by fractionating the shelled aerial part of *Pisum sativum* L. 8(1983)155

Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistan

- napier grass), *Pennisetum* sp. (bana grass), *Impomoea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985) 1
- Fermentation of hammermilled barley, lupin, pea and faba bean grain in the rumen of dairy cows 16(1986/1987)261
- The chemical composition of pea plant parts as related to harvesting time 16(1986/1987) 305
- Whole-crop peas. I. Changes in botanical and chemical composition and rumen in vitro degradability during maturation 17(1987) 15
- Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal tract of pigs 17(1987)33
- The nutritive value of white- and dark-flowered cultivars of pea for growing-finishing pigs 17(1987)245
- Utilization of white-flowered smooth-seeded spring peas (*Pisum sativum hortense*, CV Amino) by the breeding sow 22(1988)91
- Pea-crop meals**
Nutritional and growth studies with pea-crop meals and peas for growing-finishing pigs 16(1986/1987)119
- Peanut hulls**
Effect of sodium hydroxide treatment and/or extrusion cooking on the nutritive value of peanut hulls 12(1984/1985)187
- Pectin**
Determination of soluble fiber components: (1-<3; 1-<4)- β -D-glucans and pectins 23 (1989)15
- The effects of starch and pectin-rich diets on quantitative aspects of digestion in sheep 24(1989)289
- 'Pekilo protein'**
The value of 'Pekilo protein' for growing pigs 2(1977)161
- Pellet quality**
Effect of pellet size and pellet quality on pig performance 10(1983/1984)277
- Pelleting**
Influence of steam pelleting on the feeding value of tower and candle rapeseed products for broiler chickens 3(1978)251
- The effects of cereal particle size and pelleting on the nutritive value of oat-based diets for the growing pig 8(1983)91
- The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303
- Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983) 99
- Effects of pelleting, added sodium bentonite and fat in a wheat-based diet on performance and carcass characteristics of small white turkeys 12(1984/1985)223
- The effect of pelleting on the utilisation of yeast grown on n-alkanes by broiler chicks 12(1984/1985)285
- Effect of ammonia treatment and processing of whole cotton plants as sole source of roughage in complete feeds for growing cross-bred calves 13(1985)93
- The functional properties of feed raw materials and their effect on the production and quality of feed pellets 18(1987)1
- Pentosanase**
Effects of enzyme supplementation of diets based on wheat, rye or triticale on their productive value for broiler chickens 20 (1988)313
- Pepsin**
Prediction of the digestibility of ruminant feeds from their solubility in enzyme solutions 2(1977)153
- Prediction of the dry matter digestibility of tropical grasses using a pepsin-cellulase assay 2(1977)247
- Comparison of forage digestibility in vitro with enzymic solubility 11(1984)125
- A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum-pepsin treatment for predicting digestibility and intake of hays in sheep 12(1984/1985)101
- The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14 (1986)203
- Pepsin-cellulase technique**
The accuracy of the pepsin-cellulase technique for estimating the dry matter digestibility in vivo of grasses and legumes 3(1978)277
- Pharmacology**
Long term feeding of concentrate fumigated with carbon tetrachloride to sheep 2(1977) 77
- A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3 (1978)89
- Particle size distribution of a feed premix containing amprolium 3(1978)211
- Efficacy of virginiamycin and a commercially-available lactobacillus probiotic in swine diets 8(1983)69

Phosphorus

Solubility and flow of calcium, magnesium and phosphorus in the digestive tract of sheep given maize or alfalfa silages 9(1983) 131

Biological availability of phosphorus from basic steel slag 11(1984)75

The relative availability of phosphorus in inorganic feed phosphates for young turkeys and pigs 13(1985)117

Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Compositional quality of certain unconventional calcium and phosphorus sources in India for use as mineral supplements for livestock 23(1989)343

Pig-finishing

The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21

Treatment of the haem fraction of abattoir blood with formic acid to produce a solid state, low dry matter protein supplement for use in finishing-pig diets 2(1977)367

The use of effluent from grass silage in the diet of finishing pigs. I. Variation in composition of effluents 4(1979)263

The use of effluent from grass silage in the diet of finishing pigs. II. Assessment of nutritive value of fresh and stored effluent 4(1979) 275

Liquid potato feed for finishing pigs: feeding value, inclusion rate and storage properties 15(1986)129

A comparison of wheat- or barley-based diets given ad libitum as meal or pellets to finishing pigs 22(1988)263

Pig-general

The influence of feeding level and level of dietary inclusion on the digestibility of barley meal in the pig 2(1977)41

The potato (*Solanum tuberosum*) as a source of nutrients for pigs, calves and fowl - a review 2(1977)171

Influence of pericarp and endosperm colour and type on digestibility of grain sorghum by pigs 2(1977)219

Relationships between tristimulus colour values and digestibility of grain sorghum by swine 2(1977)327

Use of unextracted sunflower seeds as a protein source 5(1980)51

The energy systems used in the EEC countries: their application to 24 commercial

pig-feeds and relationships between them 6(1981)211

Prediction of the energy value of commercial pig feeds, according to the various energy systems used in the European Community, that are based on simple analytical parameters of the feed 6(1981)367

Energy evaluation of feeds and compounded diets for pigs - a review 7(1982)387

Efficacy of virginiamycin and a commercially-available lactobacillus probiotic in swine diets 8(1983)69

Comparison of bioavailable energy values of diets measured with poultry and swine 8(1983)191

Effect of storage and processing of barley on its utilization by pigs and rats 9(1983)89

Low-glucosinolate rapeseed products for pigs - a review 9(1983)239

Digestibility and energy value of sweet lupin seed (*Lupinus albus* var. multolupa) in pigs 12(1984/1985)171

Waxy and normal grain sorghums with varying tannin contents in diets for young pigs 12(1984/1985)179

Feeding value of hydroponically sprouted barley for poultry and pigs 13(1985)183

Whole-crop peas. II. Digestion of early- and late-harvested crops in the gastrointestinal tract of pigs 17(1987)33

Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19

Secretory response of the pig gut to non-starch polysaccharides 23(1989)55

Characterization of the flora of the large bowel of pigs: a status report 23(1989)79

Approach to the energetic importance of fibre digestion in pigs. I. Importance of fermentation in the overall energy supply 23(1989) 141

Approach to the energetic importance of fibre digestion in pigs. II. An experimental approach to hindgut digestion 23(1989)169

Factors affecting the digestibility by pigs of energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253

Effect of graded levels of raw and processed rapeseed on feed digestibility and nutrient utilization in young pigs 24(1989)275

Pig-growing

Utilization of yeast grown on n -paraffins in the feeding of heavy pigs, from weaning to slaughter 1(1976)25

- Adjustments of the protein level for slaughter pigs and supplementation with lysine and methionine 1(1976)619
- Lupin-seed meal (*Lupinus angustifolius* cv. Uniwhite) as a protein supplement to barley-based diets for growing pigs 1(1976)631
- Swedes (*Brassica napus*) for growing pigs: chemical composition and use as a replacement for barley in the diet 2(1977)31
- A comparison between meals prepared from the seeds of different varieties of lupin as protein supplements to barley-based diets for growing pigs 2(1977)49
- The alkaloid content of sweet lupin seed used in feeding trials on pigs and rats 2(1977)59
- The value of 'Pekilo protein' for growing pigs 2(1977)161
- The effect of supplementary protein on the performance of pigs given swedes (*Brassica napus*) as a replacement for barley in the diet 2(1977)277
- The nutritive value of liquid whey, either sour or sweet, when given in restricted amounts to the growing pig 3(1978)163
- Effects of micronization on the digestibility of whole soya beans and rapeseeds for the growing pig 3(1978)179
- Blood meal as a source of protein for grower/finisher pigs 3(1978)191
- Protein digestibility of hydrolyzed hog hair meal for pigs 3(1978)323
- Waxy versus regular maize: energy value for growing pigs and utilization in piglet diets 4(1979)109
- Olaquinox as performance-promoting feed additive for growing pigs 4(1979)117
- Value of lucerne juice and grass juice as sources of protein for the growing pig 4(1979)233
- The effect of different patterns of thermal processing of potatoes on their digestibility by growing pigs 4(1979)295
- Cabbage (*Brassica oleracea*) in the diet of growing-finisher pigs 5(1980)69
- Digestibility of a fully hydrogenated tallow for growing pigs 5(1980)87
- Value of freshly produced lucerne juice as a source of supplemental protein for the growing pig 5(1980)215
- Protein and energy value of vinasse for pigs 5(1980)221
- Lucerne juice as a protein supplement for growing pigs: effects of mineral content of the diet and of the water supply 6(1981)35
- Lupin-seed meal (*Lupinus albus* cv. Hamburg) as a source of protein for growing pigs 6(1981)285
- Effects of different handling systems on pig feed quality 7(1982)11
- Lysine requirements of imported European pigs in the humid lowland tropics from 23 to 62 kg liveweight 7(1982)233
- Comparison of the nutritional value of low and medium tannin sorghum grains for pigs 8(1983)85
- The effects of cereal particle size and pelleting on the nutritive value of oat-based diets for the growing pig 8(1983)91
- Lysine requirement of weanling European pigs fed on yellow maize and groundnut cake diets in the humid lowland tropics 8(1983)197
- Cassava meal and cassava peel meal in diets for growing pigs 8(1983)211
- A nutritional evaluation of triticale with pigs 9(1983)49
- Nutritional value and mercury content of fish silage for growing pigs 9(1983)169
- Effect of pellet size and pellet quality on pig performance 10(1983/1984)277
- The effect of level and source of protein, fibre and fat in the diet on the energy value of compounded pig feeds 11(1984)11
- The digestion of diets containing manioc by young growing pigs 12(1984/1985)119
- Anaerobically stored high moisture cereals for growing pigs 12(1984/1985)159
- The relative availability of phosphorus in inorganic feed phosphates for young turkeys and pigs 13(1985)117
- Effect of dietary protein and fat levels on the utilization of the fat in sunflower seeds by the young pig 13(1985)159
- Tryptophan content of feeds, limitations in diets and requirement for growing pigs 13(1985)171
- The evaluation of fresh sweet liquid whey as a protein supplement to maize for growing pigs 14(1986)171
- Apparent ileal and faecal digestibility of amino acids in barley-based diets supplemented with soya bean meal or canola meal for growing pigs 14(1986)183
- The absorption of dietary L-lysine and extruded L-lysine in pigs 15(1986)141
- The utilisation by growing pigs of ruminant offal hydrolysed in formic acid 15(1986)273
- Nutritional and growth studies with pea-crop meals and peas for growing-finisher pigs 16(1986/1987)119

- The apparent and true ileal digestibility of nitrogen and amino acids in lactic casein for the growing pig 16(1986/1987)169
- The prediction of the energy value of compounded pig foods from chemical analysis 17(1987)81
- High-fat maize in diets for pigs and sows 17(1987)201
- The nutritive value of white- and dark-flowered cultivars of pea for growing-finishing pigs 17(1987)245
- Canola meal as a protein supplement for growing-finishing pigs 18(1987)37
- Sugar beet molasses for growing and fattening pigs 18(1987)131
- Comparative bioavailable energy values of five ground cereal grains measured with growing rats and pigs 18(1987)143
- Evaluation of Greek cottonseed meal. 2. Use in rations for fattening pigs 18(1987)303
- A comparison of bioavailable energy values of ground cereal grains measured with adult cockerels and growing pigs 19(1988)105
- Dietary fibre and nitrogen excretion and retention by pigs 19(1988)185
- The nutritive value of hulless barley for swine 19(1988)191
- Feeding value of ensiled high-moisture maize grain with cob for growing-finishing pigs 20(1988)145
- Effect of the nature of dietary fibre on transit time and faecal excretion in the growing pig 20(1988)327
- Examination of stability, and its effect on the nutritive value, of fish silage in diets for growing pigs 21(1988)43
- Copper and iodine in pig diets with high glucosinolate rapeseed meal. I. Performance and thyroid hormone status of growing pigs fed on a diet with rapeseed meal treated with copper sulphate solution or untreated and supplements of iodine, copper or a quinoxaline derivative 22(1988)33
- Copper and iodine in pig diets with high glucosinolate rapeseed meal. II. Influence of iodine supplements for rations with rapeseed meal untreated or treated with copper ions on performance and thyroid hormone status of growing pigs 22(1988)45
- Nutritive value of diets containing triticale and varying mixtures of triticale and maize for growing-finishing swine 22(1988)217
- Digestion of diets containing molassed or plain sugar-beet pulp by growing pigs 23(1989)67
- The effect of mechanical treatment on the nutritive value of high-moisture barley and the performance of growing-finishing pigs 24(1989)121
- Body composition and growth of young pigs as affected by protein source and a growth promoter 24(1989)219
- Pig-sow**
- The nutritive value of tower rapeseed meal for swine 3(1978)235
- Pruteen in the diet of breeding pigs: reproductive performance 6(1981)297
- High-fat maize in diets for pigs and sows 17(1987)201
- Utilization of white-flowered smooth-seeded spring peas (*Pisum sativum hortense*, CV Amino) by the breeding sow 22(1988)91
- Observations on the feeding of non-lactating sows on graded levels of liquid whey 24(1989)327
- Pig-starter**
- The effect of diet complexity and physical form on the performance of starter pigs and feed processing efficiency 8(1983)99
- A continuous process for the production of purified porcine γ -globulin for use in pig-milk replacer supplements 17(1987)213
- Pig-weaner**
- The effects of air dried poultry manure on performance, organ weights and haematological measurements of weaner pigs reared in the tropics 4(1979)23
- Meat meal in the diet of the early-weaned pig. I. A comparison of meat meal and soya bean meal 4(1979)33
- Meat meal in the diet of the early-weaned pig. II. Amino acid supplementation 4(1979)43
- Meat meal in the diet of the early-weaned pig. III. Meat meal quality and the processing of meat meals 4(1979)53
- Lupinus angustifolius* cv. Unicrop as a protein- and energy-source for weaner pigs 6(1981)27
- Meat meal in the diet of the early-weaned pig. IV. The supplementation of diets with tryptophane, lysine and methionine 7(1982)27
- Effect of processing on the utilization by young pigs of the fat in soya beans and sunflower seeds 12(1984/1985)267
- The nutritive value of canola meal for early-weaned pigs 18(1987)45
- Evaluation of triticale given to pigs, poultry and rats 19(1988)359
- Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. III. Nutritive value for chicks and piglets 22(1988)23

An evaluation of skim milk powders subjected to different heating conditions during processing as dietary protein sources for the young pig 22(1988)203

Alkali-treated soya flour for piglets 24(1989)1

Pigeon pea

Variation in metabolizable energy content of raw or autoclaved white and brown varieties of three tropical grain legumes 13 (1985)141

Dry matter yield and chemical composition of pigeon pea (*C. cajan*) leaf meal and the nutritive value of pigeon pea leaf meal and grain meal for laying hens 24(1989)111

Nutritional value of pigeonpea (*Cajanus cajan*) meal in poultry diets 25(1989)123

Pimaricin

A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3 (1978)89

Plant age

Cassava for animal feeding: effect of variety and plant age on production of leaves and roots 11(1984)49

Effect of variety and plant age on the cyanide content of whole-root cassava chips and its reduction by sun-drying 11(1984)57

Plant variety

Cassava for animal feeding: effect of variety and plant age on production of leaves and roots 11(1984)49

Effect of variety and plant age on the cyanide content of whole-root cassava chips and its reduction by sun-drying 11(1984)57

Plantain sheath

In vitro studies and short-term feeding trial in lambs to evaluate plantain sheath (*Musa sapientum*) as a feed for ruminants 20 (1988)343

Polysaccharides

Secretory response of the pig gut to non-starch polysaccharides 23(1989)55

Fibre-degrading microorganisms in the monogastric digestive tract 23(1989)91

Assessing the importance of host-derived polysaccharides as carbon sources for bacteria growing in the human colon 23(1989)109

Potassium hydroxide

Predicting the energetic value of forage from digestibility of organic matter in vitro or solubility in potassium hydroxide 6(1981)235

Potato

The potato (*Solanum tuberosum*) as a source of nutrients for pigs, calves and fowl - a review 2(1977)171

The effect of different patterns of thermal processing of potatoes on their digestibility by growing pigs 4(1979)295

Nutritive value of leaf-protein concentrate from potato haulm for rats and chicks 6 (1981)413

Liquid potato feed for finishing pigs: feeding value, inclusion rate and storage properties 15(1986)129

Poultry-general

Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19

Preservatives

Laboratory application of preservatives to hay and the effects of irregular distribution on mould development 6(1981)73

Chemicals for preventing moulding in damp hay 6(1981)323

Laboratory tests of 8-quinolinol as an additive to volatile fatty acids for preserving hay and other feed-stuffs 7(1982)287

Problems of testing novel chemicals for the preservation of damp hay 8(1983)283

Losses of beta-carotene in unwilted forage crops during silage-making and feeding 9 (1983)63

Dicoumarol concentrations in sweetclover hay treated with preservatives and in spoiled hay of high- and low-coumarin cultivars of sweetclover 14(1986)221

Efficacy of sulfur dioxide as a grain preservative 19(1988)203

Pressure treatment

High temperature and pressure processing of maize cobs. II. Animal response and effect of sodium metabisulphite 14(1986)161

Proline

Amino acid composition of maize silage produced in the United Kingdom 4(1979)163

Propionic acid

Problems of testing novel chemicals for the preservation of damp hay 8(1983)283

Protein-pig

Canola meal as a protein supplement for growing-finishing pigs 18(1987)37

Protein-nitrogen flow

Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47

Proteins-cattle

Methanol-fermentation protein in veal calf nutrition 1(1976)33

Levels, sources and methods of nitrogen supplementation of maize silage for beef production 1(1976)465

Solubility of the protein of tropical pasture species and the rate of its digestion in the rumen 5(1980)183

Dry matter, protein, energy and fibre intake by dairy heifers grazing a Rhodes grass (*Chloris gayana*) pasture 6(1981)15

Rumen and post-rumen fermentation of spent tea leaf protein and other protein sources studied by the nylon bag method 7(1982)221

The digestion in the small intestine of young bulls of the protein of rapeseed meal treated or untreated with formaldehyde 7(1982)225

The use of kidney beans (*Phaseolus vulgaris*) as protein supplements in diets for young rapidly growing beef steers 12(1984/1985)1

Effect of dietary nitrogen source on microbial protein synthesis, dietary protein degradation and nutrient digestion in steers 14(1986)117

Utilization of whole rape (Canola) seed and sunflower seeds as sources of energy and protein in calf starter diets 15(1986)101

Fish silage as a protein supplement for early-weaned calves 17(1987)165

Fish protein concentrates for pre-ruminant calves: effects of processing temperatures and source of fish on protein digestibility and biological value 18(1987)181

Fresh and dried brewer's grains as protein supplements to barley straw diets given to pregnant beef cows 19(1988)33

Effects of ruminal degradability of the protein supplement on the utilization of maize-stover silage by growing beef cattle 20(1988)241

Evaluation of pea protein concentrate as a source of protein in milk replacers for Holstein calves 24(1989)267

Effect of blood meal processing method on performance and nitrogen utilization by growing cattle 24(1989)337

The effect of a protein energy supplement on pasture protein and fibre digestion in the rumen of grazing steers 25(1989)39

Proteins-chicken

House fly larvae meal grown on municipal organic waste as a source of protein in poultry diets 4(1979)227

Rapeseed meal and its use in poultry diets. A review 5(1980)255

The utilization of torula yeast in diets for White Leghorn birds during growth and laying periods 7(1982)185

Proteins-fungal

Production of fungal protein from rasped fresh cassava roots using 200- and 3000-litre fermentors 8(1983)313

Proteins-general

The digestion and metabolism of the energy and protein of maize 1(1976)455

Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977)7

Nitrogen levels in low-roughage diets for efficient rumen microbial protein production 2(1977)255

Inhibition of ruminal deamination in vitro by formaldehyde treatment of sunflower-seed, soya bean and fish meals: response curves to protective treatment 2(1977)267

Utilization of cattle manure for single-cell protein production with *Chaetomium cel-lulolyticum* 4(1979)199

Amino acid composition of two varieties of sorghum grain 5(1980)169

Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189

Protein quality of mixtures of animal protein meals 9(1983)79

Variability for protein and oil quality in *Lupinus albus* 9(1983)271

The quality of the green fraction of leaf protein concentrate from *Chenopodium quinoa* Willd grown at different levels of fertilizer nitrogen 11(1984)239

In vitro effects of avoparcin on protein degradability and rumen fermentation 15(1986)215

The effect of enzymatic treatment on amino acid content and nitrogen characteristics of feather meal 16(1986/1987)151

Effect of formaldehyde treatment on the degradation of acid-preserved fish silage protein in vitro 16(1986/1987)297

Yield and composition of protein concentrate, press cake, green juice and solubles concentrate from wet fractionation of *Sophora japonica* L. foliage 20(1988)177

- Chemical and biological evaluation of Nigerian cottonseed meal protein quality 20 (1988)251
- Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. I. Composition and nutritive evaluation 22(1988)1
- Microbial biomass protein from the fungus *Chaetomium cellulolyticum*. II. Effect of method of drying and response to amino acid supplementation 22(1988)13
- Proteins-pig**
- Adjustments of the protein level for slaughter pigs and supplementation with lysine and methionine 1(1976)619
- Lupin-seed meal (*Lupinus angustifolius* cv. Uniwite) as a protein supplement to barley-based diets for growing pigs 1(1976)631
- The composition of processed abattoir blood and its use in finishing-pig diets 2(1977)21
- A comparison between meals prepared from the seeds of different varieties of lupin as protein supplements to barley-based diets for growing pigs 2(1977)49
- The value of 'Pekilo protein' for growing pigs 2(1977)161
- The effect of supplementary protein on the performance of pigs given swedes (*Brassica napus*) as a replacement for barley in the diet 2(1977)277
- Treatment of the haem fraction of abattoir blood with formic acid to produce a solid state, low dry matter protein supplement for use in finishing-pig diets 2(1977)367
- Blood meal as a source of protein for grower/finisher pigs 3(1978)191
- Protein digestibility of hydrolyzed hog hair meal for pigs 3(1978)323
- Value of lucerne juice and grass juice as sources of protein for the growing pig 4 (1979)233
- Use of unextracted sunflower seeds as a protein source 5(1980)51
- Value of freshly produced lucerne juice as a source of supplemental protein for the growing pig 5(1980)215
- Protein and energy value of vinasse for pigs 5 (1980)221
- Lupinus angustifolius* cv. Unicrop as a protein- and energy-source for weaner pigs 6(1981)27
- Lucerne juice as a protein supplement for growing pigs: effects of mineral content of the diet and of the water supply 6(1981)35
- Lupin-seed meal (*Lupinus albus* cv. Hamburg) as a source of protein for growing pigs 6(1981)285
- Pruteen in the diet of breeding pigs: reproductive performance 6(1981)297
- The effect of level and source of protein, fibre and fat in the diet on the energy value of compounded pig feeds 11(1984)11
- Effect of dietary protein and fat levels on the utilization of the fat in sunflower seeds by the young pig 13(1985)159
- An evaluation of skim milk powders subjected to different heating conditions during processing as dietary protein sources for the young pig 22(1988)203
- Body composition and growth of young pigs as affected by protein source and a growth promoter 24(1989)219
- Proteins-rat**
- Protein quality studies on rats fed on cassava-root meal enriched with *Cephalosporium eichhorniae* 152 or with conventional plant-protein supplements 6(1981)197
- Protein and energy utilisation in rats of diets based on lupin seed (*Lupinus albus* var. multolupa) 15(1986)33
- Proteins-sheep**
- Effects of treatment of barley straw with anhydrous ammonia and supplementation with heat-treated protein meals on feed intake and liveweight performance of growing lambs 6(1981)145
- Rumen degradation of protein in sunflower meal, rapeseed meal and lupin seed placed in nylon bags 11(1984)87
- Value of mechanically extracted rape and safflower oilseed meals as protein supplements for growing lambs 11(1984)269
- Degradation of crude protein and quality of undegradable protein in untreated or formaldehyde-treated rapeseed meal 12 (1984/1985)19
- Proteins-turkey**
- Turkey feeding regimes with different ages of introduction of a choice of foods 4(1979)125
- Protozoa**
- The role of ciliate protozoa and fungi in the rumen digestion of plant cell walls 10(1983/1984)121
- Effect of absence of ciliate protozoa on rumen fluid volume, flow rate and bacterial populations in sheep 10(1983/1984)145
- Fibre-degrading microorganisms in the monogastric digestive tract 23(1989)91
- In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations 25(1989)169

Pruteen

Digestibility of protein and amino acids of a fermentation single-cell protein for veal calves 2(1977)377

Pruteen in the diet of breeding pigs: reproductive performance 6(1981)297

Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen 17(1987)1

Publishing, general

Titles of scientific papers 2(1977)1

Citation decay in animal science 2(1977)195

Pungam cake

Nutritive value of pungam (*Pongamia glabra* Vent.) cake for sheep 22(1988)321

Q**Quinoxaline**

Copper and iodine in pig diets with high glucosinolate rapeseed meal. I. Performance and thyroid hormone status of growing pigs fed on a diet with rapeseed meal treated with copper sulphate solution or untreated and supplements of iodine, copper or a quinoxaline derivative 22(1988)33

R**Rabbit**

Citrus pulp in diets for fattening rabbits 5 (1980)23

Investigations of the treatment of sawdust for rabbit feeding. 1. Effect of sodium hydroxide treatment 6(1981)43

Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51

Note on high-nitrate para grass (*Brachiaria mutica*) silage given to rabbits 9(1983)325

Studies on the nutritive value of roots, cabbage and grass silage for growing commercial rabbits 13(1985)299

Fat supplementation of diets for growing rabbits 16(1986/1987)109

Nutritional value for rabbits of olive pulp and the effects on their visceral organs 25(1989)79

Rainfall

Estimating yield and nutrient losses due to rainfall on field-drying alfalfa hay 16(1986/1987)7

Rape seed

Utilization of whole rape (Canola) seed and sunflower seeds as sources of energy and protein in calf starter diets 15(1986)101

Rapeseed

Effects of micronization on the digestibility of whole soya beans and rapeseeds for the growing pig 3(1978)179

Influence of steam pelleting on the feeding value of tower and candle rapeseed products for broiler chickens 3(1978)251

Rapeseed meal and its use in poultry diets. A review 5(1980)255

Effect of graded levels of raw and processed rapeseed on feed digestibility and nutrient utilization in young pigs 24(1989)275

Rapeseed meal

The nutritive value of tower rapeseed meal for swine 3(1978)235

Rapeseed meal tannins and egg taint 6(1981)421

The digestion in the small intestine of young bulls of the protein of rapeseed meal treated or untreated with formaldehyde 7 (1982)225

Low-glucosinolate rapeseed products for pigs - a review 9(1983)239

Rumen degradation of protein in sunflower meal, rapeseed meal and lupin seed placed in nylon bags 11(1984)87

Value of mechanically extracted rape and sunflower oilseed meals as protein supplements for growing lambs 11(1984)269

Degradation of crude protein and quality of undegradable protein in untreated or formaldehyde-treated rapeseed meal 12 (1984/1985)19

Copper and iodine in pig diets with high glucosinolate rapeseed meal. I. Performance and thyroid hormone status of growing pigs fed on a diet with rapeseed meal treated with copper sulphate solution or untreated and supplements of iodine, copper or a quinoxaline derivative 22(1988)33

Copper and iodine in pig diets with high glucosinolate rapeseed meal. II. Influence of iodine supplements for rations with rapeseed meal untreated or treated with copper ions on performance and thyroid hormone status of growing pigs 22(1988)45

Rat

The alkaloid content of sweet lupin seed used in feeding trials on pigs and rats 2(1977)59

Proximate chemical analysis of final cane molasses and effect of feeding 30% molasses on

- intestinal sucrase and maltase activities in the rat 3(1978)15
- Protein quality studies on rats fed on cassava-root meal enriched with *Cephalosporium eichhorniae* 152 or with conventional plant-protein supplements 6(1981)197
- Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6(1981)309
- Nutritive value of leaf-protein concentrate from potato haulm for rats and chicks 6 (1981)413
- Evaluation of optimum lysine and threonine supplements to a wheat and barley-based diet in rats 8(1983)163
- Effect of storage and processing of barley on its utilization by pigs and rats 9(1983)89
- Chemical composition and nutritive value of Canadian winter wheats (*Triticum aestivum* L.) given to laboratory rats 12(1984/1985)75
- Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21
- Protein and energy utilisation in rats of diets based on lupin seed (*Lupinus albus* var. multolupa) 15(1986)33
- Comparative bioavailable energy values of five ground cereal grains measured with growing rats and pigs 18(1987)143
- Evaluation of triticale given to pigs, poultry and rats 19(1988)359
- Nutritive value of Nigerian rubber seed (*Hevea brasiliensis*) meal and oil. III. Performance characteristics, relative organ weights, hematocrit and plasma metabolites of growing female rats fed corn diets containing rubber seed meal, soya bean meal or casein 20(1988)125
- Digestion and fermentation pattern of various dietary fiber sources in the rat 23(1989)195
- Reconstitution**
- The role of germination in sorghum reconstitution 14(1986)243
- Response of different sorghum grain and maize varieties to reconstitution 15(1986)231
- Red clover**
- Effects of ozonation on the chemical composition and in vitro degradation of triticale internodes and red clover stems 24(1989)159
- Rhizopus chinensis**
- Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977)7
- Rhodes grass**
- Dry matter, protein, energy and fibre intake by dairy heifers grazing a Rhodes grass (*Chloris gayana*) pasture 6(1981)15
- Rice bran**
- Trypsin inhibitor, mineral availability, and performance of broiler chickens fed on diets based on rice bran 4(1979)133
- Rice straw**
- Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7(1982)201
- Supplementation of rice straw with crop by-products. I. Legume straw supplementation 19(1988)43
- Nutritive value of rice straw (*Oryza sativa*) ensiled with animal excreta and rumen digesta 24(1989)69
- Effect of urea treatment on rumen degradation characteristics of rice straws 24(1989)83
- Rubber-seed meal**
- Nutritive value of Nigerian rubber seed (*Hevea brasiliensis*) meal and oil. III. Performance characteristics, relative organ weights, hematocrit and plasma metabolites of growing female rats fed corn diets containing rubber seed meal, soya bean meal or casein 20(1988)125
- Rumen contents**
- Nutritive value of rumen contents for monogastric animals 2(1977)351
- Rumen degradation**
- Rumen degradation of protein in sunflower meal, rapeseed meal and lupin seed placed in nylon bags 11(1984)87
- Rumen degredation**
- Evaluation of mathematical models to describe neutral detergent residue in terms of its susceptibility to degradation in the rumen 15(1986)249
- Rumen fluid**
- Effect of absence of ciliate protozoa on rumen fluid volume, flow rate and bacterial populations in sheep 10(1983/1984)145
- Effect of sodium silicate on forage digestion with rumen fluid of goats or cellulase using culture solutions adjusted for pH 24(1989)173
- Rumen ingesta kinetics**
- Rumen ingesta kinetics of cell wall components in dairy cows 25(1989)89

Rumen liquor

Effects of feeding alkali-treated olive cake on intake, digestibility and rumen liquor parameters 14(1986)139

Rumen microbiology

Nitrogen levels in low-roughage diets for efficient rumen microbial protein production 2(1977)255

Manipulation of rumen fluid pH and its influence on cellulolysis in sacco, dry matter degradation and the rumen microflora of sheep offered either hay or concentrate 10(1983/1984)1

Effect of absence of ciliate protozoa on rumen fluid volume, flow rate and bacterial populations in sheep 10(1983/1984)145

Influence of heat sterilization and ammoniation on straw composition and degradation by pure cultures of cellulolytic rumen bacteria 12(1984/1985)195

Effect of lactate on the in vitro digestion of *Agropyron elongatum* by rumen microorganisms 16(1986/1987)161

Population changes of fibrolytic rumen bacteria in the presence of phenolic acids and plant extracts 19(1988)261

The establishment of rumen microbial communities 21(1988)69

Possibilities of the use of recombinant-DNA techniques with rumen micro-organisms 21(1988)161

Consecutive batch culture - a novel technique for the in vitro study of mixed microbial populations from the rumen 21(1988)183

Factors affecting the formation of polysaccharide-degrading enzymes by rumen micro-organisms 21(1988)191

Use of the rumen simulation technique (RUSITEC) to compare microbial digestion of various by-products 21(1988)197

Lignin-polysaccharide complexes of the plant cell wall and their effect on microbial degradation in the rumen 21(1988)219

Effect of defaunating the rumen 21(1988)229

Biological structure of lignocellulose and its degradation in the rumen 21(1988)295

Colonization of guinea grass by anaerobic rumen fungi in swamp buffalo and cattle 22(1988)161

Influence of stage of forage maturity and ammoniation of wheat straw on ruminal degradation of wheat forage tissues 24(1989)201

Evaluation of various methods for protecting soya bean protein from degradation by rumen bacteria 25(1989)111

Rumen pH

Manipulation of rumen fluid pH and its influence on cellulolysis in sacco, dry matter degradation and the rumen microflora of sheep offered either hay or concentrate 10(1983/1984)1

Associative effects of mixed feeds. I. Effects of type and level of supplementation and the influence of the rumen fluid pH on cellulolysis in vivo and dry matter digestion of various roughages 10(1983/1984)15

Use of fat in diets of sheep in hot environments. II. Effect on rumen metabolism 19(1988)343

Efficacy of supplemental dietary neutralizing agents for lactating dairy cows. A review 23(1989)277

Rye grain

Feeding value of alkali-treated whole rye (*Secale cereale* L.) grain for lactating cows and its digestibility for sheep 10(1983/1984)77

Ryegrass

Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607

Ensilaged alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensilaged or frozen alkali-treated straw and ryegrass silage 3(1978)133

The effect of silage additives containing formaldehyde on the fermentation of ryegrass ensilaged at different dry matter levels and on the nutritive value of direct-cut silage 7(1982)303

The effect of urea on the preservation and digestibility in vitro of perennial ryegrass 10(1983/1984)49

The effect of anhydrous ammonia and moisture content on the preservation and chemical composition of perennial ryegrass hay 11(1984)159

Effect of prolonged wilting in poor conditions on the fermentation quality, metabolisability and net energy value of silage given to sheep 12(1984/1985)109

The chemical composition of grass silages prepared with and without pre-treatment with inoculants containing *Lactobacillus plantarum* 13(1985)269

Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Treatment of whole-crop cereals with alkali.

1. The influence of sodium hydroxide and ensiling on the chemical composition and in vitro digestibility of rye, barley and wheat

crops harvested at increasing maturity and dry matter content 18(1987)257

Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271

The effects of inoculation, addition of glucose and mincing on fermentation and proteolysis in ryegrass ensiled in laboratory silos 19(1988)85

Effects of enzyme supplementation of diets based on wheat, rye or triticale on their productive value for broiler chickens 20(1988)313

S

Safflower oilseed meal

Value of mechanically extracted rape and safflower oilseed meals as protein supplements for growing lambs 11(1984)269

Sal seed meal

Tannins in sal seed (*Shorea robusta*) and sal seed meal limit their utilization as livestock feeds 7(1982)161

Sawdust

Investigations of the treatment of sawdust for rabbit feeding. 1. Effect of sodium hydroxide treatment 6(1981)43

Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51

Seasonality

Seasonal changes in composition and intake of chaparral by Spanish goats 8(1983)47

Chemical composition and animal performance of grass forage of varying maturity stored as hay or silage 11(1984)35

Seed dehulling

A simple device for dehulling seeds and grain 3(1978)109

Selenium

Treatment of dairy sheep with soluble glass boluses containing copper, cobalt and selenium 19(1988)79

The effect of soluble-glass boluses containing copper, cobalt and selenium on the blood composition of ewes 21(1988)23

Sesbania

Canavanine content and toxicity of sesbania leaf meal for growing chicks 25(1989)137

Sesbania sesban

Yield, composition and feeding value of irrigated *Sesbania sesban* var. *Nubia* leaves harvested at latitude 38°N during a Mediterranean summer 18(1987)247

Sheep-adult

Long term feeding of concentrate fumigated with carbon tetrachloride to sheep 2(1977)77

The nutritional value for sheep of molasses and condensed molasses solubles 2(1977)143

Utilization by sheep of dried pig faeces containing a high concentration of copper 3(1978)31

Differences between cattle and sheep in their digestion and relative intake of a mature tropical grass hay 3(1978)41

Estimation of the digestibility of low-quality hays by cattle from measurements made with sheep 3(1978)51

Spent tea leaf as a ruminant feed 3(1978)219

Chemical composition of culinary wastes and their potential as a feed for ruminants 5(1980)205

Intake and digestibility in sheep fed on alkali-treated straw and concentrates, as complete diets or separate feeds 7(1982)271

Solubility and flow of calcium, magnesium and phosphorus in the digestive tract of sheep given maize or alfalfa silages 9(1983)131

Effect of intake on digestibility of plant cell walls and cell contents of complete diets for ruminants 9(1983)197

Manipulation of rumen fluid pH and its influence on cellulolysis in sacco, dry matter degradation and the rumen microflora of sheep offered either hay or concentrate 10(1983/1984)1

Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65

Feeding value of alkali-treated whole rye (*Secale cereale* L.) grain for lactating cows and its digestibility for sheep 10(1983/1984)77

The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149

Feeding value of apple pomace silage for sheep 11(1984)189

Degradation of crude protein and quality of undegradable protein in untreated or formaldehyde-treated rapeseed meal 12(1984/1985)19

A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum-pepsin treatment

- for predicting digestibility and intake of hays in sheep 12(1984/1985)101
- Effect of prolonged wilting in poor conditions on the fermentation quality, metabolisability and net energy value of silage given to sheep 12(1984/1985)109
- The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12 (1984/1985)141
- The effect of supplementing grass silage with sodium hydroxide-treated chopped or milled straw on intake and digestibility by wether sheep 12(1984/1985)215
- The voluntary intake and digestibility of combinations of cereal crop residues and legume hay for sheep 12(1984/1985)241
- Performance and carcass characteristics of Sudan desert sheep and goats on high roughage diets with added fat 13(1985)147
- Studies on the feeding value of agro-industrial by-products. III. Digestibility of cocoa-pod and cocoa-pod-based diets by ruminants 13 (1985)249
- Corrugated cardboard and dried poultry manure as dietary ingredients for sheep 14 (1986)265
- The effect of monensin on rumen fermentation in sheep fed on all-roughage and barley/roughage diets 15(1986)7
- The effect of grinding groundnut hulls either with or without alkali treatment on digestibility of diet and on ruminal and blood components 15(1986)69
- The digestion by sheep of barley and maize-based diets containing either meat and bone meal or soya bean meal 15(1986)301
- Comparison between two methods of feeding concentrate to sheep given a diet high or low in concentrate with or without buffering substances 16(1986/1987)37
- Relative intakes, digestibility and bacterial protein synthesis by sheep and cattle fed high-roughage diets 16(1986/1987)75
- Composition of vine-branches with leaves of *Vitis vinifera* L. fresh and ensiled using different additives (previous results) 16(1986/1987)89
- The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99
- Effect of defaunation on the loss of energy in wethers fed different quantities of cellulose and normal or steamflaked maize starch 16 (1986/1987)233
- Effect of formaldehyde-treated soya bean meal and urea in starters on nitrogen quality, degradability in sacco, sheep digestibility and calf performance 16(1986/1987)287
- Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19
- The effect of dietary barley on carbohydrate digestibility of sulphur dioxide-treated wheat straw by sheep 18(1987)55
- Nutritive value of naked oats for ruminants 18(1987)83
- Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271
- Treatment of whole-crop cereals with alkali. 3. Voluntary intake and digestibility studies in sheep given immature wheat ensiled with sodium hydroxide, urea or ammonia 18(1987)283
- Treatment of dairy sheep with soluble glass boluses containing copper, cobalt and selenium 19(1988)79
- Use of fat in diets of sheep in hot environments. II. Effect on rumen metabolism 19 (1988)343
- Evaluation of mixed lucerne and caucasian bluestem hay diets fed to sheep 20(1988) 19
- The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171
- The intake and digestion of maize silage-based diets by dairy cows and sheep 20 (1988)299
- The effect of dietary inclusion of starch, sucrose and xylose on the utilization of dietary energy in sheep 21(1988)11
- The effect of soluble-glass boluses containing copper, cobalt and selenium on the blood composition of ewes 21(1988)23
- Amino acids flow to and absorption from the small intestine of sheep fed barley and sulfur dioxide-treated straw at different ratios 22(1988)147
- Digestibility in vivo of ensiled grapevines (branches and leaves): influence of the system of analysis in the detergent fibre scheme on the prediction of digestibility 22 (1988)173
- The nutritive value of ensiled forages cut at three stages of growth 22(1988)237
- Effect of soaking, ensilage and hydrogen peroxide treatment of barley straw on rumen degradability 22(1988)295
- Effect of ammoniation and energy supplementation on the utilization of wheat straw by sheep 22(1988)305
- Nutritive value of pungam (*Pongamia glabra* Vent) cake for sheep 22(1988)321

- The effects of starch and pectin-rich diets on quantitative aspects of digestion in sheep 24(1989)289
- Oven and stack ammoniation of grass hays. 2. Voluntary intake, digestibility and faecal nitrogen excretion in sheep in relation to hay composition 24(1989)313
- Effect of formaldehyde treatment on the rumen digestion of barley and wheat grain and its influence on hay intake and digestion by sheep 25(1989)55
- By-products from the canning industry as feedstuff for ruminants: digestibility and its prediction from chemical composition and laboratory bioassays 25(1989)67
- Digestion rates in and outflow rates from the rumen of sheep fed untreated or calcium hydroxide-treated wheat straw 25(1989)179
- Sheep-general**
- Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73
- Comparative digestibility of feedstuffs by sheep and cows 12(1984/1985)47
- The effect of five different supplements on the degradation of straw in sheep given untreated barley straw 19(1988)289
- Use of fat in diets of sheep in hot environments. I. Effects on performance, carcass characteristics and lipid composition of plasma 19(1988)327
- Sheep-lamb**
- Poly-unsaturated fat in young ruminants 2(1977)93
- Reducing the potential copper toxicity of concentrates to sheep by the use of molybdenum and sulphur supplements 2(1977)235
- The use of whole barley diets fortified with solutions of urea, minerals and vitamins for lambs 2(1977)307
- The effect of ammoniation on the intake and nutritive value of straw 4(1979)169
- Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979)177
- Nutritive value of thermoammoniated and steam-treated maize stover II. Rumen metabolites and rate of passage 4(1979)187
- Citrus pulp in diets for fattening lambs 5(1980)11
- Effect of dry-matter concentration of milk substitute and method of feeding on intake and performance by lambs 5(1980)321
- Effects of treatment of barley straw with anhydrous ammonia and supplementation with heat-treated protein meals on feed intake and liveweight performance of growing lambs 6(1981)145
- Utilization of alkali-treated grain. 1. Alkali-treated grain in complete diets for steers and lambs 6(1981)273
- Use of processing to explore the factors affecting the digestion of forage cell walls 6(1981)387
- Use of fish protein hydrolysate in milk replacers 7(1982)135
- Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375
- Nutritive value of almond hulls for sheep 8(1983)63
- The effects of sulphur supplementation on cellulose digestion in vitro and on nutrient digestion, nitrogen metabolism and rumen characteristics of lambs fed on good quality fescue and tropical star grass hays 8(1983)129
- Methods of improving the utilization of cereal straw by ruminants. II. Further studies on dried grass as a supplement for lambs 8(1983)229
- Can lantana seeds be utilized in lamb rations? 9(1983)29
- Energy and nitrogen balance, intake and growth rate of lambs fed on unwilted grass silages treated with a formaldehyde-acid mixture or untreated 9(1983)107
- Chemical composition, digestibility and feeding value of maize cobs 9(1983)121
- The use of Balanites kernel cake in a diet for fattening sheep 9(1983)301
- Effect of absence of ciliate protozoa on rumen fluid volume, flow rate and bacterial populations in sheep 10(1983/1984)145
- Biological availability of phosphorus from basic steel slag 11(1984)75
- Maize whole stillage as a supplemental protein source for growing lambs and finishing steers 11(1984)181
- Nitrogen balance and abomasal protein-nitrogen flow in growing ruminants fed methane digester effluent in combination with distillers dried grains 13(1985)47
- High temperature and pressure processing of maize cobs. II. Animal response and effect of sodium metabisulphite 14(1986)161
- Effects of supplementing a wheat straw diet with urea or a urea-molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25

- Energy metabolism in lambs fed silage diets with sucrose or a growth promoter 16 (1986/1987)203
- Cassava meal as supplement to napier grass diets for growing sheep and goats 18(1987) 197
- SO₂ treatment for converting straw into a concentrate-like feed: a growth study with lambs 19(1988)219
- Effect of sorghum straw alone or in combination with molasses and nitrogen sources on performance of Sudan desert lambs 19 (1988)351
- Nitrogen utilization, ruminal fermentation and abomasal nitrogen flow in sheep fed caucasian bluestem hay supplemented with lucerne or urea 20(1988)33
- The maize replacement value of cassava flour in rations for growing ewe lambs 20(1988) 233
- In vitro studies and short-term feeding trial in lambs to evaluate plantain sheath (*Musa sapientum*) as a feed for ruminants 20 (1988)343
- Pilot plant production of sulphur dioxide-treated mesquite wood. II. Digestibility and feedlot performance by lambs 22(1988) 61
- Nutritive value of banana stalk (*Musa cavendishi*) as a feed for sheep 22(1988)327
- Fermentation and utilization by lambs of sugarcane harvested fresh and ensiled with and without NaOH. 4. Ruminal kinetics 23 (1989)323
- Silage-alfalfa**
Solubility and flow of calcium, magnesium and phosphorus in the digestive tract of sheep given maize or alfalfa silages 9(1983) 131
- Silage-animal waste**
Ensiling of rumen contents and blood with wheat straw 18(1987)67
- Silage-apple pomace**
Feeding value of apple pomace silage for sheep 11(1984)189
- Some properties of apple pomace ensiled with and without additives 15(1986)57
- Silage-barley**
Ensiled alkali-treated straw. I. Effect of level and type of alkali on the composition and digestibility in vitro of ensiled barley straw 3(1978)117
- Ensiled alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensiled or frozen alkali-treated straw and ryegrass silage 3(1978)133
- Alkali treatment of straw for ruminants. II.**
Nutritive value of straw ensiled after alkali treatment. 3(1978)289
- Effect of soaking, ensilage and hydrogen peroxide treatment of barley straw on rumen degradability 22(1988)295
- Silage-beet**
Distillery effluents as animal feed: the use of condensed beet molasses stillage (CBMS) in broiler feeding 5(1980)233
- Silage-clover**
The enzymic nature of the degradation of beta-carotene in red clover and in other forage crops during silage-making with acid additives 5(1980)59
- Silage-fish**
Nutritional value and mercury content of fish silage for growing pigs 9(1983)169
- Thiaminase activity in fish silage and moist fish feed 13(1985)39
- Effect of formaldehyde treatment on the degradation of acid-preserved fish silage protein in vitro 16(1986/1987)297
- Fish silage as a protein supplement for early-weaned calves 17(1987)165
- Examination of stability, and its effect on the nutritive value, of fish silage in diets for growing pigs 21(1988)43
- Silage-forage, general**
Losses of beta-carotene in unwilted forage crops during silage-making and feeding 9 (1983)63
- Silage-general**
Treatment of whole-crop cereals with alkali.
1. The influence of sodium hydroxide and ensiling on the chemical composition and in vitro digestibility of rye, barley and wheat crops harvested at increasing maturity and dry matter content 18(1987)257
- Treatment of whole-crop cereals with alkali. 2. Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271
- The nutritive value of ensiled forages cut at three stages of growth 22(1988)237
- Silage-grass**
The intake of steers and heifers fed on mixtures of alkali-treated or untreated straw with grass silage 2(1977)315
- The use of effluent from grass silage in the diet of finishing pigs. I. Variation in composition of effluents 4(1979)263

- The use of effluent from grass silage in the diet of finishing pigs. II. Assessment of nutritive value of fresh and stored effluent 4(1979)275
- A calorimetric investigation of the nutritive value of formic acid- and formaldehyde-formic acid-treated grass silages 6(1981)245
- Feed consumption and performance of dairy cows with alternate feedings of grass silage and maize silage 6(1981)337
- The effect of silage additives containing formaldehyde on the fermentation of ryegrass ensiled at different dry matter levels and on the nutritive value of direct-cut silage 7(1982)303
- Investigations on the effect of sodium bentonite and other moisture absorbents on the production of effluent from grass silage 8(1983)107
- Energy and nitrogen balance, intake and growth rate of lambs fed on wilted grass silages treated with a formaldehyde-acid mixture or untreated 9(1983)107
- Chemical composition and animal performance of grass forage of varying maturity stored as hay or silage 11(1984)35
- Digestibility and digesta retention in dairy cows receiving hay or silage at varying concentrate levels 11(1984)279
- The effect of supplementing grass silage with sodium hydroxide-treated chopped or milled straw on intake and digestibility by wether sheep 12(1984/1985)215
- The chemical composition of grass silages prepared with and without pre-treatment with inoculants containing *Lactobacillus plantarum* 13(1985)269
- Studies on the nutritive value of roots, cabbage and grass silage for growing commercial rabbits 13(1985)299
- The substitution of grass silage by a straw and concentrate mixture for dairy cows 20(1988)135
- The digestibility and metabolisable energy content of grass silage and their prediction from laboratory measurements 24(1989)27
- Silage-legume**
- Ensiling of leguminous green forages in combination with different dry roughages and molasses 12(1984/1985)133
- Silage-lucerne**
- SO₂-treated straw as a silage additive: fermentation data on lucerne with particular reference to protein degradation 22(1988)247
- Silage-maize**
- Forage maize in the rations of dairy cows and fattening cattle in Bulgaria 1(1976)45
- Cultivation of maize in irrigated areas for silage and grain production and their use by ruminants 1(1976)237
- Influence of ensiling maize ears and grain on chemical composition, conservation losses and digestibility 1(1976)301
- Effect of maize silage harvest stage on yield, plant composition and fermentation losses 1(1976)313
- Maize silage with or without NPN, dehydrated whole-crop maize pellets or high moisture maize grain for finishing bulls 1(1976)347
- Energy supplementation of maize silage harvested at different maturity stages 1(1976)369
- Factors influencing the composition and nutritive value of ensiled whole-crop maize 1(1976)381
- Rate of growth, feed utilisation and digestibility observed with diets containing ensiled maize grain or maize ears fed alone or in mixtures 1(1976)401
- Voluntary intake and efficiency of utilisation of whole-crop maize silage 1(1976)441
- Levels, sources and methods of nitrogen supplementation of maize silage for beef production 1(1976)465
- The nutritive value (energy) of maize silage 1(1976)481
- Effect of weight at slaughter on the efficiency of fattening young cattle 1(1976)505
- Economic aspects of high-plane feeding of bulls with maize silage 1(1976)513
- Maize silage and whole maize plant pellets as the basic forages in complete feeds for milk cows 3(1978)23
- A note on the effect on the aerobic deterioration of maize silage of the manipulation of the microflora by means of antibiotics 3(1978)89
- Amino acid composition of maize silage produced in the United Kingdom 4(1979)163
- Feed consumption and performance of dairy cows with alternate feedings of grass silage and maize silage 6(1981)337
- Solubility and flow of calcium, magnesium and phosphorus in the digestive tract of sheep given maize or alfalfa silages 9(1983)131
- Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185

Effects of stage of maturity and addition of molasses on nutritive value of maize stover silage 12(1984/1985)65

Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91

The effect of nitrogen supplements to maize silage on the amount and composition of the nitrogen fraction entering the intestine of cattle 13(1985)25

Influence of harvest date on the nutritive value and ensiling characteristics of maize stover 14(1986)11

High grain content maize silage in fattening diets of young male cattle 16(1986/1987)63

Feeding value of ensiled high-moisture maize grain with cob for growing-finishing pigs 20(1988)145

The intake and digestion of maize silage-based diets by dairy cows and sheep 20(1988)299

Effects of added dietary urea on the utilization of maize stover silage by growing beef cattle 22(1988)227

Inoculation of corn silage and high-moisture corn with lactic acid bacteria and its effects on the subsequent fermentations and on feedlot performance of beef steers 25(1989)23

Silage-manure-maize

Laboratory evaluation of ensiled mixtures of alkali-treated cattle excreta and whole-crop maize 3(1978)335

Silage-orange peel

The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20(1988)335

Silage-orange pulp

Effect of urea on the ensiling process of orange pulp 12(1984/1985)233

Silage-para grass

Note on high-nitrate para grass (*Brachiaria mutica*) silage given to rabbits 9(1983)325

Silage-quality assessment

A titration method for silage assessment 22(1988)341

Silage-ryegrass

The effect of some pre-ensiling treatments on silage composition and nitrogen disappearance in the rumen 8(1983)259

Effect of prolonged wilting in poor conditions on the fermentation quality, metabolisability and net energy value of silage given to sheep 12(1984/1985)109

Energy metabolism in lambs fed silage diets with sucrose or a growth promoter 16(1986/1987)203

The effects of inoculation, addition of glucose and mashing on fermentation and proteolysis in ryegrass ensiled in laboratory silos 19(1988)85

Silage-sorghum

Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299

Preservation of sorghum plant portions harvested, processed and ensiled at ten stages of maturity 7(1982)257

Studies on fermentation of sorghum silage during storage, and its effect on milk animals 9(1983)143

Effects of molasses/urea and bacterial inoculant additives on silage quality, dry matter recovery, and feeding value for cattle 12(1984/1985)205

Silage-straw

Methods of improving the utilization of cereal straw by ruminants. III. A note on the effect of ensiling straw treated with urea 9(1983)181

Ensiling characteristics of cattle waste fermented with wheat straw and green berseem (*Trifolium alexandrinum*) 25(1989)149

Silage-sugar beet

A note on the use of sugar-beet pulp silage and molasses in the diet of lactating dairy cows 20(1988)13

Counteracting structure loss in pressed sugar beet pulp silage 24(1989)97

Silage-sugar cane

Nutritive value for ruminants of sugar cane bagasse ensiled after spray treatment with different levels of NaOH 9(1983)1

Fermentation and utilization by lambs of sugarcane harvested fresh and ensiled with and without NaOH. 4. Ruminant kinetics 23(1989)323

Silage-tropical forages

Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistan napier grass), *Pennisetum* sp. (bana grass), *Impomea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985)1

Silage-vine

Composition of vine-branches with leaves of *Vitis vinifera* L. fresh and ensiled using different additives (previous results) 16(1986/1987)89

Silage-wheat

Wheat silage for fattening young male cattle 12(1984/1985)253

Treatment of whole-crop cereals with alkali. 4. Voluntary intake and growth in steers given wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55

Silage disease

Inhibition of *Listeria monocytogenes* by defined lactic microflora in gnotobiotic silages of lucerne, fescue, ryegrass and maize - influence of dry matter and temperature 2(1977)297

Silage effluent

Investigations on the effect of sodium bentonite and other moisture absorbents on the production of effluent from grass silage 8(1983)107

Silica

Effects of the addition of silica to the diet of the laying hen, with particular reference to egg-shell quality 6(1981)157

Single cell protein

Utilization of cattle manure for Torula yeast production from straw hydrolysates 5(1980)175

Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen 17(1987)1

Single-cell protein

Composition and nutritive value of single-cell protein (SCP) 1(1976)9

Digestibility of protein and amino acids of a fermentation single-cell protein for veal calves 2(1977)377

Nutritive value of single-cell protein produced by *Chaetomium cellulolyticum* grown on maize stover and pulp-mill sludge 11(1984)211

Sisal

Feed value of Tanzanian sisal wastes: chemical composition and digestibility in vitro 14(1986)255

Sodium acrylate

The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Effect of sodium acrylate and calcium and sodium hydroxides on the fermentation, digestibility, chemical composition and aerobic deterioration of whole-crop maize silage 12(1984/1985)91

Sodium bentonite

Investigations on the effect of sodium bentonite and other moisture absorbents on the production of effluent from grass silage 8(1983)107

Effects of pelleting, added sodium bentonite and fat in a wheat-based diet on performance and carcass characteristics of small white turkeys 12(1984/1985)223

Sodium bicarbonate

Effects of sodium chloride and sodium bicarbonate in the diet on the performance of calves 5(1980)309

Efficacy of supplemental dietary neutralizing agents for lactating dairy cows. A review 23(1989)277

Sodium chloride

Effects of sodium chloride and sodium bicarbonate in the diet on the performance of calves 5(1980)309

Sodium diacetate

Storage of large round bales of hay harvested at different moisture concentrations and treated with sodium diacetate and/or covered with plastic 13(1985)69

Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and stored at different moisture contents 17(1987)45

Sodium hydroxide

The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Ammonia treatment of straw via the hydrolysis of urea. II. Additions of soya bean (urease), sodium hydroxide and molasses; effects on the digestibility of urea-treated straw 11(1984)115

The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149

Sodium metabisulphite

High temperature and pressure processing of maize cobs. II. Animal response and effect of sodium metabisulphite 14(1986)161

Sodium silicate

Effect of sodium silicate on forage digestion with rumen fluid of goats or cellulase using culture solutions adjusted for pH 24(1989)173

Soil chemistry

Soil chemistry and mineral problems in farm livestock. A review 5(1980)95

Sophora japonica

Yield and composition of protein concentrate, press cake, green juice and solubles concentrate from wet fractionation of *Sophora japonica* L. foliage 20(1988)177

Sorbic acid

The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20(1988)335

Sorghum

Influence of pericarp and endosperm colour and type on digestibility of grain sorghum by pigs 2(1977)219

Relationships between tristimulus colour values and digestibility of grain sorghum by swine 2(1977)327

Preservation of sorghum plant portions harvested, processed and ensiled at ten stages of maturity 7(1982)257

Proximate analysis, amino acid composition and tannin content of improved Nigerian sorghum varieties and their potential in poultry feeds 7(1982)359

Waxy and normal grain sorghums with varying tannin contents in diets for young pigs 12(1984/1985)179

The role of germination in sorghum reconstitution 14(1986)243

Yield and potential feeding value of straws of grain sorghum cultivars developed in Nigeria 17(1987)285

Effects of wood ash extract treatment on the feeding value and utilization of high-tannin sorghums by broiler chicks 22(1988)131

Factors affecting the digestibility by pigs of energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253

Sorghum grain

A comparison of three fibre methods for predicting the metabolizable energy content of sorghum grain for poultry 2(1977)197

Amino acid composition of two varieties of sorghum grain 5(1980)169

Comparison of the nutritional value of low and medium tannin sorghum grains for pigs 8(1983)85

Response of different sorghum grain and maize varieties to reconstitution 15(1986)231

Treatment of low-tannin sorghum grain for broiler feed 20(1988)69

Sorghum silage

Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299

Studies on fermentation of sorghum silage during storage, and its effect on milch animals 9(1983)143

Sorghum straw

Effect of sorghum straw alone or in combination with molasses and nitrogen sources on performance of Sudan desert lambs 19(1988)351

Soya bean

Effects of micronization on the digestibility of whole soya beans and rapeseeds for the growing pig 3(1978)179

Effect of processing on the utilization by young pigs of the fat in soya beans and sunflower seeds 12(1984/1985)267

Nutrient and trypsin inhibitor content of hydroponically sprouted soya beans 13(1985)203

Effects of moisture level, treatment time and soya bean addition on the nutritive value of urea-treated maize stover 19(1988)67

The energy content of full-fat soya beans in meal and pelleted feeds for adult cocks and broilers 19(1988)97

Alkali-treated soya flour for piglets 24(1989)1

Evaluation of various methods for protecting soya bean protein from degradation by rumen bacteria 25(1989)111

Soya bean hulls

Colonization of thick-walled plant tissues by anaerobic fungi 19(1988)25

Soya bean meal

Inhibition of ruminal deamination in vitro by formaldehyde treatment of sunflower-seed, soya bean and fish meals: response curves to protective treatment 2(1977)267

Alkali treatment of straw for ruminants. I. Utilization of complete diets containing straw by beef cattle 3(1978)143

Meat meal in the diet of the early-weaned pig. I. A comparison of meat meal and soya bean meal 4(1979)33

Ammonia treatment of straw via the hydrolysis of urea. II. Additions of soya bean (urease), sodium hydroxide and molasses; effects on the digestibility of urea-treated straw 11(1984)115

Effects of isopropyl alcohol treatment of soya bean meal on nitrogen utilization by ruminants 14(1986)127

- Apparent ileal and faecal digestibility of amino acids in barley-based diets supplemented with soya bean meal or canola meal for growing pigs 14(1986)183
- The digestion by sheep of barley and maize-based diets containing either meat and bone meal or soya bean meal 15(1986)301
- Effect of formaldehyde-treated soya bean meal and urea in starters on nitrogen quality, degradability in sacco, sheep digestibility and calf performance 16(1986/1987)287
- Nutritive value of Nigerian rubber seed (*Hevea brasiliensis*) meal and oil. III. Performance characteristics, relative organ weights, hematocrit and plasma metabolites of growing female rats fed corn diets containing rubber seed meal, soya bean meal or casein 20(1988)125
- Spruce muka**
Nutritive value of spruce muka (foliage) for the growing chick 3(1978)63
- Starch**
Further thermotolerant fungi for the conversion of cassava starch to protein 2(1977)7
Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73
The effect of β -glucanase on the utilization of starch and nitrogen by broiler chickens fed on barley of low- or high-viscosity 15(1986)83
The effect of dietary inclusion of starch, sucrose and xylose on the utilization of dietary energy in sheep 21(1988)11
The effects of starch and pectin-rich diets on quantitative aspects of digestion in sheep 24(1989)289
- Steam treatment**
Effect of high-pressure steam treatment on chemical composition and digestibility in vitro of roughages 7(1982)61
- Stearic acid**
Feed intake and milk yield at increasing supplement of a palmitic and stearic acid-rich type of fat in comparison with animal fat 22(1988)179
- Stockfeed, general**
Energetics of stockfeed production 3(1978)1
- Storage**
Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299
Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381
- Effects of different handling systems on pig feed quality 7(1982)11
- The effect of prolonged storage on the digestibility and nitrogen content of ammonia-treated barley straw 8(1983)147
- Effects of dietary maize solubles and minerals on albumen quality of fresh and stored brown-shelled eggs 8(1983)177
- Effect of storage and processing of barley on its utilization by pigs and rats 9(1983)89
- Studies on fermentation of sorghum silage during storage, and its effect on milch animals 9(1983)143
- Investigations into the effects of processing on the retention of the carotenoid fractions of *Leucaena leucocephala* during storage, and the effects of processing on mimosine concentration 9(1983)307
- The influence of processing and storage on hydrogen cyanide and tannin contents of para-rubber seed and its products 9(1983)319
- Anaerobically stored high moisture cereals for growing pigs 12(1984/1985)159
- Storage of large round bales of hay harvested at different moisture concentrations and treated with sodium diacetate and/or covered with plastic 13(1985)69
- Liquid potato feed for finishing pigs: feeding value, inclusion rate and storage properties 15(1986)129
- Nutrient losses during deterioration of hay in relation to changes in biochemical composition and microbial growth 15(1986)149
- The nutritive value of barley for monogastric animals and for ruminants as affected by artificial drying and alkali and anaerobic storage 15(1986)285
- Microfloral and compositional changes in alfalfa hay treated with sodium diacetate and stored at different moisture contents 17(1987)45
- Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227
- Microbial dynamics in barley grain stored under controlled atmosphere 19(1988)299
- Preservation and storage of green panic (*Panicum maximum*) as moist hay with urea 20(1988)87
- The effect of method of preservation of forages on their resistance to reduction in particle size with an artificial masticator 24(1989)185

Straw-barley

Effect on digestibility and nitrogen content of barley straw of different ammonia treatments 2(1977)131

Ensilaged alkali-treated straw. I. Effect of level and type of alkali on the composition and digestibility in vitro of ensilaged barley straw 3(1978)117

Ensilaged alkali-treated straw. II. The nutritive value for young beef cattle of mixtures of ensilaged or frozen alkali-treated straw and ryegrass silage 3(1978)133

Alkali treatment of straw for ruminants. I. Utilization of complete diets containing straw by beef cattle 3(1978)143

Alkali treatment of straw for ruminants. II. Nutritive value of straw ensilaged after alkali treatment. 3(1978)289

Effect of time of storage and method of drying on the digestibility in vitro of alkali-treated barley straw 3(1978)381

Effects of treatment of barley straw with anhydrous ammonia and supplementation with heat-treated protein meals on feed intake and liveweight performance of growing lambs 6(1981)145

Studies on ammonia-treated straw. II. Fixation of ammonia in treated straw by spraying with acids 7(1982)53

The feeding value for ruminants of straw and whole-crop barley and oats treated with anhydrous or aqueous ammonia or urea 8(1983)247

The effect of urease enzyme on treatment time and the nutritive value of straw treated with ammonia as urea 8(1983)271

Digestion of organic matter and nitrogen in ammoniated barley straw 10(1983/1984)65

A comparison between degradation in vitro and in sacco of constituents of untreated and ammonia-treated barley straw 10(1983/1984)199

Digestibility studies on ammonia-treated straw 10(1983/1984)213

The digestion of untreated and ammonia-treated barley straw in an artificial rumen 10(1983/1984)229

Ammonia-treated barley straw and rolled barley offered either together, in a mixed ration, or successively to beef steers 10(1983/1984)247

A comparison of alkali treatment methods to improve the nutritive value of straw. I. Digestibility and metabolizability 12(1984/1985)295

Treatment of barley straw with ammonia and sulphur dioxide gases under laboratory conditions 14(1986)41

A comparison of alkali treatment methods used to improve the nutritive value of straw. II. In sacco and in vitro degradation relative to in vivo digestibility 14(1986)215

Evaluation of the increase in feeding value of barley straw following mild treatment with sulphur dioxide 15(1986)13

Near infrared reflectance spectroscopy of untreated and ammoniated barley straw 15(1986)189

Effect of straw treatment and laboratory of incubation on degradation of neutral detergent residue in the rumen 18(1987)75

Fresh and dried brewer's grains as protein supplements to barley straw diets given to pregnant beef cows 19(1988)33

Effects of ammonia and sulphur dioxide gases on the composition and digestion of barley straw 19(1988)121

Fibre degradation in the rumens of animals receiving hay, untreated or ammonia-treated straw 19(1988)277

The effect of five different supplements on the degradation of straw in sheep given untreated barley straw 19(1988)289

Effect of soaking, ensilage and hydrogen peroxide treatment of barley straw on rumen degradability 22(1988)295

Upgrading barley straw for goats: the effectiveness of a sodium hydroxide and urea dip method 24(1989)57

Straw-cotton

Effects of ammonia treatment and processing of cotton straw as the sole source of roughage in complete feeds on rumen characteristics with Murrah buffaloes 14(1986)95

Straw-general

The effect on digestibility of a new technique for alkali treatment of straw 1(1976)73

Review article: the alkali treatment of straws 2(1977)105

Enzyme solubility - a method for evaluating the digestibility of alkali-treated straw 2(1977)205

The intake of steers and heifers fed on mixtures of alkali-treated or untreated straw with grass silage 2(1977)315

A new technique for upgrading the nutritive value of roughages under field conditions 3(1978)201

Cellulase insoluble fibre as a measure of unavailable organic matter in cattle compounds containing alkali-treated straw 3(1978)227

- The effect of ammoniation on the intake and nutritive value of straw 4(1979)169
- Utilization of alkali-treated grain. 2. Utilization by steers of diets based on hay or straw and mixed with either NaOH-treated or rolled barley 6(1981)347
- Intake and digestibility in sheep fed on alkali-treated straw and concentrates, as complete diets or separate feeds 7(1982)271
- The effect of prolonged storage on the digestibility and nitrogen content of ammonia-treated barley straw 8(1983)147
- Methods of improving the utilization of cereal straw by ruminants. I. Supplements of urea, molasses and dried grass and treatment with sodium hydroxide 8(1983)221
- Methods of improving the utilization of cereal straw by ruminants. II. Further studies on dried grass as a supplement for lambs 8(1983)229
- Methods of improving the utilization of cereal straw by ruminants. III. A note on the effect of ensiling straw treated with urea 9(1983)181
- A note on the OECD collaborative experiment on ammonia-treated straw 10(1983/1984)89
- Effect of defaunation on straw digestion in sacco and on particle retention in the rumen 10(1983/1984)165
- Ammonia treatment of straw: methods for treatment and feeding experience in Norway 10(1983/1984)173
- Ammonia versus sodium hydroxide treatment of straw for growing cattle 10(1983/1984)189
- Commercial experience of treating straw with ammonia 10(1983/1984)223
- Anhydrous ammonia-treated cereal straw for animal feeding. Some results from the Mediterranean area 10(1983/1984)239
- The value of urea-treated straw in diets of lactating goats 11(1984)67
- Ammonia treatment of straw via the hydrolysis of urea. I. Effects of dry matter and urea concentrations on the rate of hydrolysis of urea 11(1984)103
- Ammonia treatment of straw via the hydrolysis of urea. II. Additions of soya bean (urease), sodium hydroxide and molasses; effects on the digestibility of urea-treated straw 11(1984)115
- The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149
- Influence of heat sterilization and ammoniation on straw composition and degradation by pure cultures of cellulolytic rumen bacteria 12(1984/1985)195
- The effect of supplementing grass silage with sodium hydroxide-treated chopped or milled straw on intake and digestibility by wether sheep 12(1984/1985)215
- Methods of improving the utilization of cereal straw by ruminants. IV. The effect on intake of level of supplementation with urea and dried grass 13(1985)281
- Methods of improving the utilization of cereal straw by ruminants. V. The effect of molasses and dried grass 13(1985)293
- The use of barley straw as a feed 14(1986)29
- Urea as a source of ammonia for improving the nutritive value of wheat straw 14(1986)67
- Alkaline hydrogen peroxide treatments of crop residues to increase ruminal dry matter disappearance in sacco 17(1987)179
- Yield and potential feeding value of straws of grain sorghum cultivars developed in Nigeria 17(1987)285
- The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility in vitro and cell wall degradability 19(1988)159
- The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173
- The substitution of grass silage by a straw and concentrate mixture for dairy cows 20(1988)135
- The feeding value of ammoniated flax straw, wheat straw and wheat chaff for beef cattle 21(1988)57
- Genetic variation in the feeding value of cereal straw 21(1988)127
- Near infra-red spectra of ammonia-treated straw and of isolated cell walls 21(1988)209
- Upgrading of straw by ammoniation 21(1988)287
- SO₂-treated straw as a silage additive: the participation of soluble and cell wall monosaccharide residues in lucerne silage fermentation 22(1988)255
- Straw-grass**
- Titratable groups and soluble phenolic compounds as indicators of the digestibility of chemically treated roughages 6(1981)123
- Straw-grass seed**
- Composition and enzymatic digestibility of oregon grass straws 9(1983)283

Straw-oat

Studies on ammonia-treated straw. I. The effects of type and level of ammonia, moisture content and treatment time on the digestibility in vitro and enzyme soluble organic matter of oat straw 7(1982)45

Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71

The effect of calcium hydroxide treatment on the digestibility of chopped oat straw supplemented with a solution containing urea, calcium, phosphorus, sodium, trace elements and vitamins 7(1982)93

The intake and digestibility by sheep of oat straw and maize stover offered with different levels of noug (*Guizotia abyssinica*) meal 16(1986/1987)99

Straw-rice

Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7(1982)201

Supplementation of rice straw with crop by-products. I. Legume straw supplementation 19(1988)43

Nutritive value of rice straw (*Oryza sativa*) ensiled with animal excreta and rumen digesta 24(1989)69

Effect of urea treatment on rumen degradation characteristics of rice straws 24(1989)83

Straw-sorghum

Effect of sorghum straw alone or in combination with molasses and nitrogen sources on performance of Sudan desert lambs 19(1988)351

Straw-wheat

Performance of fattening cattle and lactating beef cows on diets high in chemically treated wheat straw and poultry litter 5(1980)299

Ammonia and sodium hydroxide treatment of wheat straw in diets for fattening steers 7(1982)1

Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71

Effect of treatment of wheat straw with ammonia and peracetic acid on digestibility in vitro and cell wall composition 7(1982)325

Digestibility of diets containing increasing levels of NaOH-treated or untreated wheat straw 7(1982)331

Ammoniation of wheat straw and native grass hay during baling of large round bales 8(1983)35

Utilization of alkali-treated and neutralized wheat straw-based rations for growing goat kids 9(1983)155

Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185

The response of wheat straw varieties to mild sulphur dioxide treatment 10(1983/1984)269

Improving the intake and nutritive value of wheat straw for beef cows 12(1984/1985)125

The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12(1984/1985)141

Nutritive evaluation of uromol-wheat straw as a basal ration for buffaloes 13(1985)103

Effects of ethanalamine and ethylenediamine treatments on the composition and degradability of wheat straw 13(1985)131

Chemical preparation of NaOH-keratin hydrolysate for improving the nutritive value of wheat straw 13(1985)215

Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81

Effects of supplementing a wheat straw diet with urea or a urea-molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25

Colonization of thick-walled plant tissues by anaerobic fungi 19(1988)25

SO₂ treatment for converting straw into a concentrate-like feed: a growth study with lambs 19(1988)219

Amino acids flow to and absorption from the small intestine of sheep fed barley and sulfur dioxide-treated straw at different ratios 22(1988)147

SO₂-treated straw as a silage additive: fermentation data on lucerne with particular reference to protein degradation 22(1988)247

Effect of ammoniation and energy supplementation on the utilization of wheat straw by sheep 22(1988)305

Influence of stage of forage maturity and ammoniation of wheat straw on ruminal degradation of wheat forage tissues 24(1989)201

Ensiling characteristics of cattle waste fermented with wheat straw and green berseem (*Trifolium alexandrinum*) 25(1989)149

Digestion rates in and outflow rates from the rumen of sheep fed untreated or calcium hydroxide-treated wheat straw 25(1989)179

Straw hydrolysates

Utilization of cattle manure for *Torula* yeast production from straw hydrolysates 5(1980) 175

Sucrase

Proximate chemical analysis of final cane molasses and effect of feeding 30% molasses on intestinal sucrase and maltase activities in the rat 3(1978)15

Sucrose

Energy metabolism in lambs fed silage diets with sucrose or a growth promoter 16 (1986/1987)203

The effect of dietary inclusion of starch, sucrose and xylose on the utilization of dietary energy in sheep 21(1988)11

Sugar beet

Sugar beet molasses for growing and fattening pigs 18(1987)131

Utilization of sugar beet molasses and monensin for finishing dairy bullocks 25(1989) 11

Sugar beet pulp

Dried sugar beet pulp as a high energy feed for beef cattle 1(1976)643

Sugar beet pulp - a review 8(1983)1

Sugar beet pulp products for dairy cows 15 (1986)123

A note on the use of sugar-beet pulp silage and molasses in the diet of lactating dairy cows 20(1988)13

The effects of barley, unmolassed sugar-beet pulp and molasses supplements on organic matter, nitrogen and fibre digestion in the rumen of cattle given a silage diet 20(1988) 259

The effect of dietary inclusion of starch, sucrose and xylose on the utilization of dietary energy in sheep 21(1988)11

Digestion of diets containing molassed or plain sugar-beet pulp by growing pigs 23 (1989)67

Sugar beet pulp silage

Counteracting structure loss in pressed sugar beet pulp silage 24(1989)97

Sugarcane

Studies on the fermentation of chopped sugarcane 7(1982)411

The effects of ammonia on the fermentation of chopped sugarcane 9(1983)291

Effect of sugarcane condensed molasses solubles, sugarcane molasses and monensin on performance and volatile fatty acid production in finishing steers 12(1984/1985)275

Fermentation and utilization by lambs of sugarcane harvested fresh and ensiled with and without NaOH. 4. Ruminant kinetics 23 (1989)323

Sugarcane bagasse

Nutritive value for ruminants of sugar cane bagasse ensiled after spray treatment with different levels of NaOH 9(1983)1

Sulphur

Reducing the potential copper toxicity of concentrates to sheep by the use of molybdenum and sulphur supplements 2(1977)235

Nutritive value of lupin-seed protein and leaf-protein concentrates supplemented with various sulphur sources 6(1981)189

The effects of sulphur supplementation on cellulose digestion *in vitro* and on nutrient digestion, nitrogen metabolism and rumen characteristics of lambs fed on good quality fescue and tropical star grass hays 8(1983) 129

Fractions of nitrogen, sulphur, phosphorus, calcium and magnesium in the herbage of perennial ryegrass as influenced by fertilizer nitrogen 14(1986)231

Sulphur dioxide

The response of wheat straw varieties to mild sulphur dioxide treatment 10(1983/1984) 269

Treatment of barley straw with ammonia and sulphur dioxide gases under laboratory conditions 14(1986)41

Evaluation of the increase in feeding value of barley straw following mild treatment with sulphur dioxide 15(1986)13

The effect of dietary barley on carbohydrate digestibility of sulphur dioxide-treated wheat straw by sheep 18(1987)55

Effects of ammonia and sulphur dioxide gases on the composition and digestion of barley straw 19(1988)121

Efficacy of sulfur dioxide as a grain preservative 19(1988)203

SO₂ treatment for converting straw into a concentrate-like feed: a growth study with lambs 19(1988)219

Pilot plant production of sulphur dioxide-treated mesquite wood. II. Digestibility and feedlot performance by lambs 22(1988) 61

Amino acids flow to and absorption from the small intestine of sheep fed barley and sulfur dioxide-treated straw at different ratios 22(1988)147

SO₂-treated straw as a silage additive: fermentation data on lucerne with particular reference to protein degradation 22(1988) 247

Sunflower cake

Feeding value of sunflower and groundnut cakes for broilers 4(1979)143

Feeding value of sunflower- and groundnut-cakes for laying hens 6(1981)63

Sunflower meal

Rumen degradation of protein in sunflower meal, rapeseed meal and lupin seed placed in nylon bags 11(1984)87

Sunflower seed

Inhibition of ruminal deamination in vitro by formaldehyde treatment of sunflower-seed, soya bean and fish meals: response curves to protective treatment 2(1977)267

Use of unextracted sunflower seeds as a protein source 5(1980)51

Effect of processing on the utilization by young pigs of the fat in soya beans and sunflower seeds 12(1984/1985)267

Effect of dietary protein and fat levels on the utilization of the fat in sunflower seeds by the young pig 13(1985)159

Utilization of whole rape (Canola) seed and sunflower seeds as sources of energy and protein in calf starter diets 15(1986)101

Supplementation-vine

Composition of vine-branches with leaves of *Vitis vinifera* L. fresh and ensiled using different additives (previous results) 16(1986/1987)89

Swedes

Swedes (*Brassica napus*) for growing pigs: chemical composition and use as a replacement for barley in the diet 2(1977)31

The effect of supplementary protein on the performance of pigs given swedes (*Brassica napus*) as a replacement for barley in the diet 2(1977)277

Sweet potato

Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73

Effects of ensiling or drying on five forage species in western Kenya: *Zea mays* (maize stover), *Pennisetum purpureum* (Pakistan napier grass), *Pennisetum* sp. (bana grass), *Impomea batata* (sweet potato vines) and *Cajanus cajan* (pigeon pea leaves) 13(1985)1

Sweetclover

Dicoumarol concentrations in sweetclover hay treated with preservatives and in spoiled hay of high- and low-coumarin cultivars of sweetclover 14(1986)221

T**Tall fescue**

Effects of *Drechslera sorokiniana* infection on the yield and quality of tall fescue forage 15(1986)41

Tallow

Digestibility of a fully hydrogenated tallow for growing pigs 5(1980)87

Tamarind seeds

The nutritive value of tamarind seeds for broiler chicks 22(1988)285

Tannin

Rapeseed meal tannins and egg taint 6(1981)421

Tannins in sal seed (*Shorea robusta*) and sal seed meal limit their utilization as livestock feeds 7(1982)161

Proximate analysis, amino acid composition and tannin content of improved Nigerian sorghum varieties and their potential in poultry feeds 7(1982)359

Comparison of the nutritional value of low and medium tannin sorghum grains for pigs 8(1983)85

The influence of processing and storage on hydrogen cyanide and tannin contents of para-rubber seed and its products 9(1983)319

Waxy and normal grain sorghums with varying tannin contents in diets for young pigs 12(1984/1985)179

Utilization of tannin-containing and tannin-free faba beans (*Vicia faba*) by young chicks: effects of pelleting feeds on energy, protein and starch digestibility 20(1988)59

Treatment of low-tannin sorghum grain for broiler feed 20(1988)69

Effects of wood ash extract treatment on the feeding value and utilization of high-tannin sorghums by broiler chicks 22(1988)131

Tea

Spent tea leaf as a ruminant feed 3(1978)219

Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7(1982)201

Rumen and post-rumen fermentation of spent tea leaf protein and other protein sources studied by the nylon bag method 7(1982)221

Techniques-amino acid evaluation

The determination of available lysine in barley using *Tetrahymena pyriformis* W. 5 (1980)33

A comparison of three methods for the determination of available lysine in barley 5 (1980)43

Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21

Techniques-ammonia treatment evaluation

Near infra-red spectra of ammonia-treated straw and of isolated cell walls 21(1988)209

Techniques-compound feed evaluation

The use of an enzymatic technique to predict digestibility, metabolizable and net energy of compound feedstuffs for ruminants 14 (1986)203

Techniques-delignification evaluation

Titratable groups and soluble phenolic compounds as indicators of the digestibility of chemically treated roughages 6(1981)123

Techniques-digestibility

Comparison of methods for predicting digestibility of feeds 20(1988)203

Techniques-digestibility evaluation

The analytical and biological accuracy of estimating the dry matter digestibility of different legume species 1(1976)61

Enzyme solubility - a method for evaluating the digestibility of alkali-treated straw 2 (1977)205

Prediction of the dry matter digestibility of tropical grasses using a pepsin-cellulase assay 2(1977)247

Principles governing the use of chemical methods for assessing the nutritive value of forages: a review 3(1978)265

Some remarks on the analytical procedure of Van Soest for the prediction of forage digestibility 3(1978)309

A note on Onozuka 3S cellulase as a replacement for Onozuka SS (P1500) cellulase when estimating forage digestibility in vitro 5(1980)347

Accuracy of predicting digestibility by the cellulase technique; the effect of pretreatment of forage samples with neutral detergent or acid pepsin 7(1982)83

Suitability of cell-wall constituents as predictors of organic matter digestibility in some tropical and subtropical by-products 7(1982)191

A new biological method for determining amino acid digestibility in poultry feedstuffs using a simple cannula, and the influence of dietary fibre on endogenous amino acid output 12(1984/1985)29

A comparison of laboratory methods involving single or two-stage cellulase and two-stage rumen inoculum-pepsin treatment for predicting digestibility and intake of hays in sheep 12(1984/1985)101

Improving the reliability of the two-stage in vitro assay for ruminant feed digestibility by calibration against in vivo data from a wide range of sources 18(1987)233

The use of a cellulase technique to predict digestibility, metabolizable and net energy of forages 19(1988)247

Comparison of methods for predicting digestibility of feeds 20(1988)203

A comparison of microbial, enzymatic, chemical and near-infrared reflectance spectroscopy methods in forage evaluation 20 (1988)219

Techniques-dry matter evaluation

Sources of error in predicting digestible dry matter from the acid-detergent fiber content of forages 21(1988)205

Techniques-dual flow continuous culture system

Evaluation of a dual flow continuous culture system for estimating bacterial fermentation in vivo of mixed diets containing various soya bean products 16(1986/1987)51

Techniques-dye absorption test

The dye absorption test as an indicator of the nutritive quality of heat-processed navy bean 9(1983)71

Techniques-energy availability evaluation

A comparison between fermentation heat of forages and organic matter digestibility determined by in vitro incubation with rumen fluid 23(1989)333

Techniques-fat and protein evaluation

Discriminating power of the nylon bag technique and the pepsin-cellulase method to detect less commonly used components included in concentrates for dairy cows 25 (1989)1

Techniques-fibre determination

A comparison of three fibre methods for predicting the metabolizable energy content of sorghum grain for poultry 2(1977)197

Some remarks on the determination of acid-detergent fibre in feeds with cetyltrimethylammonium bromide versus determination of detergent fibre with alkylarylsulfonic acid 6(1981)91

Techniques—forage digestibility prediction

Prediction of forage digestibility by principal component analysis of near infrared reflectance spectra 16(1986/1987)215

Techniques—monocarbonyl determination

Thermofractographic determination of monocarbonyl compounds in animal waste fats used as feed fat 13(1985)227

Techniques—near-infrared reflectance

Bias when predicting crude protein, dry matter digestibility and voluntary intake of tropical grasses by near-infrared reflectance 9(1983)221

Techniques—neutral detergent fibre evaluation

The study of several modifications of the neutral detergent fibre procedure 9(1983)19

Techniques—nutritive value

A new technique for upgrading the nutritive value of roughages under field conditions 3 (1978)201

Techniques—organic matter prediction

Comparison of laboratory methods for predicting the organic matter digestibility of forages 2(1977)337

Techniques—polysaccharide measurement

Classification and measurement of plant polysaccharides 23(1989)27

Techniques—preservative application

Laboratory application of preservatives to hay and the effects of irregular distribution on mould development 6(1981)73

Techniques—rumen microbiology

Possibilities of the use of recombinant-DNA techniques with rumen micro-organisms 21 (1988)161

Consecutive batch culture – a novel technique for the in vitro study of mixed microbial populations from the rumen 21(1988)183

Techniques—seed dehulling

A simple device for dehulling seeds and grain 3(1978)109

Techniques—silage assessment

A titration method for silage assessment 22 (1988)341

Techniques—straw spectroscopy

Near infrared reflectance spectroscopy of untreated and ammoniated barley straw 15 (1986)189

Tetrahymena pyriformis

The determination of available lysine in barley using *Tetrahymena pyriformis* W. 5 (1980)33

Thermal treatment

Nutritive value of thermoammoniated and steam-treated maize stover I. Intake, digestibility and nitrogen retention 4(1979) 177

Nutritive value of thermoammoniated and steam-treated maize stover II. Rumen metabolites and rate of passage 4(1979)187

The effect of different patterns of thermal processing of potatoes on their digestibility by growing pigs 4(1979)295

Investigations of the treatment of sawdust for rabbit feeding. 2. Effect of boiling and incubation with poultry manure 6(1981)51

The effect of steam-conditioning rate on the pelleting process 6(1981)83

The effect of some pre-ensiling treatments on silage composition and nitrogen disappearance in the rumen 8(1983)259

The influence of steam pelleting and grinding on the nutritive value of Canola rapeseed for poultry 8(1983)303

The dye absorption test as an indicator of the nutritive quality of heat-processed navy bean 9(1983)71

High temperature and pressure processing of maize cobs. I. Digestibility in vitro of processed cobs 12(1984/1985)151

Influence of heat sterilization and ammoniation on straw composition and degradation by pure cultures of cellulolytic rumen bacteria 12(1984/1985)195

Influence of supplemental fat and steam pelleting on the response of broilers to diets varying in energy density 13(1985)57

High temperature and pressure processing of maize cobs. II. Animal response and effect of sodium metabisulphite 14(1986)161

Thiaminase

Thiaminase activity in fish silage and moist fish feed 13(1985)39

Threonine

Evaluation of optimum lysine and threonine supplements to a wheat and barley-based diet in rats 8(1983)163

Thyroprotein

Effect of alkali-treated rice straw supplemented with spent tea leaf and thyroprotein on milk yield, milk composition and certain physiological parameters of dairy cows 7(1982)201

Timothy

Composition and fibre digestion in morphological components of an alfalfa-timothy sward 19(1988)135

Torula

Utilization of cattle manure for Torula yeast production from straw hydrolysates 5(1980) 175

The utilization of torula yeast in diets for White Leghorn birds during growth and laying periods 7(1982)185

Toxicology

Detoxification of aflatoxins in groundnut meal 2(1977)67

Long term feeding of concentrate fumigated with carbon tetrachloride to sheep 2(1977) 77

Reducing the potential copper toxicity of concentrates to sheep by the use of molybdenum and sulphur supplements 2(1977)235

Utilization by sheep of dried pig faeces containing a high concentration of copper 3 (1978)31

Soil chemistry and mineral problems in farm livestock. A review 5(1980)95

Can lantana seeds be utilized in lamb rations? 9(1983)29

Variability for protein and oil quality in *Lupinus albus* 9(1983)271

The use of Balanites kernel cake in a diet for fattening sheep 9(1983)301

The influence of processing and storage on hydrogen cyanide and tannin contents of para-rubber seed and its products 9(1983) 319

Note on high-nitrate para grass (*Brachiaria mutica*) silage given to rabbits 9(1983)325

Effect of cassava-based diets varying in cyanide content on the performance and physiopathology of the African giant rat (*Cricetomys gambianus* Waterhouse) 11 (1984)1

Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin) 11(1984)293

Hyperexcitability in cattle fed ammoniated roughages 17(1987)121

The effect of toxic substances present in some leaf protein concentrates on their nutritive value 24(1989)151

Canavanine content and toxicity of sesbania leaf meal for growing chicks 25(1989)137

In vitro metabolism of mycotoxins by bacterial, protozoal and ovine ruminal fluid preparations 25(1989)169

Trimethylamine oxide

The involvement of trimethylamine oxide in fish meal in the production of egg taint 8 (1983)119

Triticale

A nutritional evaluation of triticale with pigs 9(1983)49

Nutritive value of three cultivars of triticale for sheep, pigs and poultry 18(1987)19

Evaluation of triticale given to pigs, poultry and rats 19(1988)359

Effects of enzyme supplementation of diets based on wheat, rye or triticale on their productive value for broiler chickens 20 (1988)313

Nutritional value of triticale (Carman) for broiler diets 22(1988)119

Nutritive value of diets containing triticale and varying mixtures of triticale and maize for growing-finishing swine 22(1988)217

The effect of fat level and type on the utilization of triticale (cultivar Carman) by broiler chicks 22(1988)273

Effects of ozonation on the chemical composition and in vitro degradation of triticale internodes and red clover stems 24(1989) 159

Trypsin

Trypsin inhibitor, mineral availability, and performance of broiler chickens fed on diets based on rice bran 4(1979)133

Nutrient and trypsin inhibitor content of hydroponically sprouted soya beans 13(1985) 203

Tryptophan

Meat meal in the diet of the early-weaned pig. II. Amino acid supplementation 4(1979)43

Meat meal in the diet of the early-weaned pig. IV. The supplementation of diets with tryptophane, lysine and methionine 7 (1982)27

Tryptophan content of feeds, limitations in diets and requirement for growing pigs 13 (1985)171

Utilisation of L- and DL-amino acids by broiler chicks fed diets containing Pruteen 17(1987)1

Turkey

Turkey feeding regimes with different ages of introduction of a choice of foods 4(1979)125

Limiting amino acids in dried microbial cells given to young turkeys 4(1979)209

A note on the nutritive value of dried instant coffee residue for broiler chickens and turkey poults 10(1983/1984)285

Effects of pelleting, added sodium bentonite and fat in a wheat-based diet on performance and carcass characteristics of small white turkeys 12(1984/1985)223

The relative availability of phosphorus in inorganic feed phosphates for young turkeys and pigs 13(1985)117

Influence of nutrient density on feed consumption, weight gain and gut capacity of broilers, Leghorns and turkeys reared to 26 days of age 16(1986/1987)129

U

Urea

The use of whole barley diets fortified with solutions of urea, minerals and vitamins for lambs 2(1977)307

Studies on fermentation of sorghum silage during storage - effect of urea and molasses 3(1978)299

Performance of heifers fed on diets based on NaOH-treated maize cobs and the effect of supplementary urea and source of carbohydrates 4(1979)97

Evaluation of corn gluten meal with urea as a source of supplementary nitrogen for growing calves and lambs 7(1982)375

Methods of improving the utilization of cereal straw by ruminants. I. Supplements of urea, molasses and dried grass and treatment with sodium hydroxide 8(1983)221

The effect of urea on the preservation and digestibility in vitro of perennial ryegrass 10(1983/1984)49

The value of urea-treated straw in diets of lactating goats 11(1984)67

Ammonia treatment of straw via the hydrolysis of urea. I. Effects of dry matter and urea concentrations on the rate of hydrolysis of urea 11(1984)103

Effects of molasses/urea and bacterial inoculant additives on silage quality, dry matter recovery, and feeding value for cattle 12(1984/1985)205

Effect of urea on the ensiling process of orange pulp 12(1984/1985)233

Effects of supplementing a wheat straw diet with urea or a urea-molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25

Effect of formaldehyde-treated soya bean meal and urea in starters on nitrogen quality, degradability in sacco, sheep digestibility and calf performance 16(1986/1987)287

Complexes of urea and formaldehyde as non-protein nitrogen compounds in ruminant rations: a review 20(1988)1

Nitrogen utilization, ruminal fermentation and abomasal nitrogen flow in sheep fed

caucasian bluestem hay supplemented with lucerne or urea 20(1988)33

Preservation and storage of green panic (*Panicum maximum*) as moist hay with urea 20(1988)87

The effect of treatment with urea, sorbic acid, or dehydration on orange peel silage 20(1988)335

Effects of added dietary urea on the utilization of maize stover silage by growing beef cattle 22(1988)227

Urease

The effect of urease enzyme on treatment time and the nutritive value of straw treated with ammonia as urea 8(1983)271

Rumen fermentation, urease activity and performance of cattle given palm kernel cake-based diet 20(1988)79

Uromol

Nutritive evaluation of uromol-wheat straw as a basal ration for buffaloes 13(1985)103

V

Vinasse

Protein and energy value of vinasse for pigs 5(1980)221

Vine

Composition of vine-branches with leaves of *Vitis vinifera* L. fresh and ensiled using different additives (previous results) 16(1986/1987)89

Virginiamycin

Efficacy of virginiamycin and a commercially-available lactobacillus probiotic in swine diets 8(1983)69

Vitamin C

Effect of dietary ascorbic acid (vitamin C) supplementation on the performance of broiler chickens in a tropical environment 16(1986/1987)17

Vitamin K

Comparison of the relative biological availability and stability of commercial sources of vitamin K activity 3(1978)329

Vitamins, general

The use of whole barley diets fortified with solutions of urea, minerals and vitamins for lambs 2(1977)307

W

Water supply

Lucerne juice as a protein supplement for growing pigs: effects of mineral content of the diet and of the water supply 6(1981)35

Weeds

Utilization of aquatic plants *Elodea canadensis* and *Hydrilla verticillata* in broiler chick diets 20(1988)155

Weende components

Apparent and true digestibility of the Weende components, cell content and cell wall of ryegrass 1(1976)607

Wheat

Nutritive value of leaf protein coagulum and cereals mixtures 4(1979)91

Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6(1981)309

Evaluation of optimum lysine and threonine supplements to a wheat and barley-based diet in rats 8(1983)163

The effect of calcium and sodium hydroxides and of sodium acrylate on the fermentation and digestibility in vitro of ensiled whole-crop wheat and barley harvested at different stages of maturity 9(1983)37

Effects of milling and pelleting of maize, barley and wheat on their metabolizable energy value for cockerels and chicks 9(1983)99

The effect of milling and sodium hydroxide treatment on intake and digestibility of wheat straw by sheep 11(1984)149

Observations on the chemical and nutritive content of white winter and spring wheats contaminated with deoxynivalenol (vomitoxin) 11(1984)293

Chemical composition and nutritive value of Canadian winter wheats (*Triticum aestivum* L.) given to laboratory rats 12(1984/1985)75

Effects of pelleting, added sodium bentonite and fat in a wheat-based diet on performance and carcass characteristics of small white turkeys 12(1984/1985)223

Estimation of optimum supplements of essential amino acids to a wheat and barley-based diet in rats using a nitrogen balance technique 15(1986)21

Feeding value of sprouted wheat (*Triticum aestivum*) for beef cattle finishing diets 15(1986)113

Treatment of whole-crop cereals with alkali.

1. The influence of sodium hydroxide and ensiling on the chemical composition and in vitro digestibility of rye, barley and wheat crops harvested at increasing maturity and dry matter content 18(1987)257

Treatment of whole-crop cereals with alkali. 2.

Voluntary intake and digestibility by sheep of rye, barley and wheat crops ensiled with sodium hydroxide 18(1987)271

Treatment of whole-crop cereals with alkali. 3.

Voluntary intake and digestibility studies in sheep given immature wheat ensiled with sodium hydroxide, urea or ammonia 18(1987)283

Effects of enzyme supplementation of diets

based on wheat, rye or triticale on their productive value for broiler chickens 20(1988)313

A comparison of wheat- or barley-based diets

given ad libitum as meal or pellets to finishing pigs 22(1988)263

Effect of soft white wheat addition to alfalfa-

grass forage on heifer gain, diet digestibility and in vitro digestion kinetics 24(1989)141

Factors effecting the digestibility by pigs of

energy and protein in wheat, barley and sorghum diets supplemented with canola meal 24(1989)253

Effect of formaldehyde treatment on the rumen

digestion of barley and wheat grain and its influence on hay intake and digestion by sheep 25(1989)55

The effect of β -glucanase supplementation on

nutrient digestibility and growth in broilers given diets containing barley, oat groats or wheat 25(1989)193

Wheat grain

Turkey feeding regimes with different ages of introduction of a choice of foods 4(1979)125

Wheat silage

Wheat silage for fattening young male cattle 12(1984/1985)253

Treatment of whole-crop cereals with alkali. 4.

Voluntary intake and growth in steers given wheat ensiled with sodium hydroxide, urea or ammonia 19(1988)55

Wheat straw

Performance of fattening cattle and lactating beef cows on diets high in chemically treated wheat straw and poultry litter 5(1980)299

Ammonia and sodium hydroxide treatment of wheat straw in diets for fattening steers 7(1982)1

- Botanical composition and extent of lignification affecting digestibility of wheat and oat straw and paspalum hay 7(1982)71
- Effect of treatment of wheat straw with ammonia and peracetic acid on digestibility in vitro and cell wall composition 7(1982)325
- Digestibility of diets containing increasing levels of NaOH-treated or untreated wheat straw 7(1982)331
- Ammoniation of wheat straw and native grass hay during baling of large round bales 8(1983)35
- Utilization of alkali-treated and neutralized wheat straw-based rations for growing goat kids 9(1983)155
- Effect of wilting and the additives, straw, molasses and urea on the fermentation pattern of maize silage 9(1983)185
- The response of wheat straw varieties to mild sulphur dioxide treatment 10(1983/1984)269
- Improving the intake and nutritive value of wheat straw for beef cows 12(1984/1985)125
- The utilization of untreated and calcium hydroxide treated wheat straw by sheep 12(1984/1985)141
- Nutritive evaluation of uromol-wheat straw as a basal ration for buffaloes 13(1985)103
- Effects of ethanolamine and ethylenediamine treatments on the composition and degradability of wheat straw 13(1985)131
- Chemical preparation of NaOH-keratin hydrolysate for improving the nutritive value of wheat straw 13(1985)215
- Aqueous ammonia treatment of wheat straw: voluntary intake and digestibility in cattle 14(1986)81
- Effects of supplementing a wheat straw diet with urea or a urea-molasses block and/or cottonseed meal on intake and liveweight change of lambs 16(1986/1987)25
- Colonization of thick-walled plant tissues by anaerobic fungi 19(1988)25
- The effect of ammoniation on the nutritive value of wheat, barley and oat straws. I. Changes in chemical composition in relation to digestibility in vitro and cell wall degradability 19(1988)159
- The effect of ammoniation on the nutritive value of wheat, barley and oat straws. II. Digestibility and energy value measurements in vivo and their prediction from laboratory measurements 19(1988)173
- SO₂ treatment for converting straw into a concentrate-like feed: a growth study with lambs 19(1988)219
- The effect of variation in the water content of wheat straw before ammoniation on its nutritive value for beef cattle 20(1988)111
- Upgrading of straw by ammoniation 21(1988)287
- Amino acids flow to and absorption from the small intestine of sheep fed barley and sulphur dioxide-treated straw at different ratios 22(1988)147
- SO₂-treated straw as a silage additive: fermentation data on lucerne with particular reference to protein degradation 22(1988)247
- SO₂-treated straw as a silage additive: the participation of soluble and cell wall monosaccharide residues in lucerne silage fermentation 22(1988)255
- Effect of ammoniation and energy supplementation on the utilization of wheat straw by sheep 22(1988)305
- Influence of stage of forage maturity and ammoniation of wheat straw on ruminal degradation of wheat forage tissues 24(1989)201
- Ensiling characteristics of cattle waste fermented with wheat straw and green berseem (*Trifolium alexandrinum*) 25(1989)149
- Digestion rates in and outflow rates from the rumen of sheep fed untreated or calcium hydroxide-treated wheat straw 25(1989)179
- Whey**
- The replacement of skim milk by dried microbial cells and whey in milk diets for calves 2(1977)85
- The nutritive value of liquid whey, either sour or sweet, when given in restricted amounts to the growing pig 3(1978)163
- The evaluation of fresh sweet liquid whey as a protein supplement to maize for growing pigs 14(1986)171
- Observations on the feeding of non-lactating sows on graded levels of liquid whey 24(1989)327
- Wilting**
- Effect of prolonged wilting in poor conditions on the fermentation quality, metabolisability and net energy value of silage given to sheep 12(1984/1985)109
- Influence of processing methods and storage time on the cyanide potential of cassava leaf meal 17(1987)227
- Wood ash extract**
- Effects of wood ash extract treatment on the feeding value and utilization of high-tannin sorghums by broiler chicks 22(1988)131

X

Xanthophylls

Investigations into the effects of processing on the retention of the carotenoid fractions of *Leucaena leucocephala* during storage, and the effects of processing on mimosine concentration 9(1983)307

Xylose

The effect of dietary inclusion of starch, sucrose and xylose on the utilization of dietary energy in sheep 21(1988)11

Y

Yam

Raw and steam-pelleted cassava, sweet potato and yam *cayenensis* as starch sources for ruminant and chicken diets 3(1978)73

Yeast

Utilization of yeast grown on *n*-paraffins in the feeding of heavy pigs, from weaning to slaughter 1(1976)25

Nutritional evaluation, using rats, of diets based on cereals and ethanol-grown yeast 6(1981)309

The utilization of torula yeast in diets for White Leghorn birds during growth and laying periods 7(1982)185

The effect of pelleting on the utilisation of yeast grown on *n*-alkanes by broiler chicks 12(1984/1985)285

Yellow turnip

Studies on the nutritive value of roots, cabbage and grass silage for growing commercial rabbits 13(1985)299

Z

Zinc

The effect of age of regrowth on content and release of manganese, iron, zinc and copper from four tropical forages incubated in sacco in rumen of sheep 20(1988)171

